

04.01.06 Four - Dimensional (4-D) and Five - Dimensional (5-D) Fetal Ultrasound(s)

Original Effective Date: April 1999

Review Date: January 2026

Revised: January 2024

DISCLAIMER/INSTRUCTIONS FOR USE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations, or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

This Medical Policy document describes the status of medical technology at the time the document was developed. Since that time, new technology may have emerged, or new medical literature may have been published. This Medical Policy will be reviewed regularly and updated as scientific and medical literature becomes available; therefore, policies are subject to change without notice.

Summary

Description

Note: This evidence review addresses the use of 4-D, and 5-D fetal ultrasounds in maternity care. For review of 3-D fetal ultrasounds through eviCore, refer to [Wellmark's Authorization Table](#). This evidence review does not apply to ultrasound performed for non-pregnancy related conditions.

A fetal ultrasound is a test performed during pregnancy to assess for pregnancy and rule out ectopic pregnancy and confirm gestational age early on. As pregnancy advances typically in the second and third trimesters ultrasounds are utilized to assess the fetal size and position, heartbeat, congenital malformations, placental abnormalities, and measuring the volume of amniotic fluid.

4-D ultrasounds create computer generated images viewed on a video monitor that provide more detail and can produce more life-like images of the fetus.

5-D ultrasounds have been proposed to automate ultrasounds through artificial intelligence to reduce exposure time, dependency on operator skill and experience and increase reproducibility.

Summary of Evidence

For individuals who are pregnant who receive a four-dimensional (4-D) or five-dimensional (5-D) ultrasound(s) for the diagnosis of fetal cardiac abnormalities the evidence includes 2 observational studies. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. No clinical utility studies were identified. All of the studies evaluated the diagnostic accuracy of 4-D ultrasound. None of the studies evaluated 5-D ultrasound. All but one study was conducted outside of the United States. Findings from these studies are inconclusive as they were heterogenous in the types of fetal abnormalities evaluated and they did not provide adequate details on variance of effect to assess level of precision. The evidence is insufficient to determine the technology results in an improvement in the net health outcomes.

For individuals who are pregnant who receive a four-dimensional (4-D) or five-dimensional (5-D) ultrasound(s) for the diagnosis of fetal noncardiac abnormalities the evidence included 4 diagnostic accuracy studies. Relevant outcomes are diagnostic accuracy, symptoms, functional outcomes, quality of life, and treatment-related morbidity. No clinical utility studies were identified. All of the studies evaluated the diagnostic accuracy of 4-D ultrasound. None of the studies evaluated 5-D ultrasound. None were conducted in the United States. Findings from these studies are inconclusive as they were heterogenous in the types of fetal abnormalities evaluated and they did not provide adequate details on variance of effect to assess level of precision. The evidence is insufficient to determine the technology results in an improvement in the net health outcomes.

Additional Information

Not applicable

OBJECTIVE

The objective of this evidence review is to determine whether the use of four-dimensional (4-D) or five-dimensional (5-D) fetal ultrasound(s) improves the net health outcomes.

PRIOR APPROVAL

Not applicable.

POLICY

Note: This evidence review addresses the use of 4-D, and 5-D fetal ultrasounds in maternity care. For review of 3-D fetal ultrasounds through eviCore, refer to [Wellmark's Authorization Table](#). This evidence review does not apply to ultrasound performed for non-pregnancy related conditions.

The use of four-dimensional (4-D) and/or five-dimensional (5-D) fetal ultrasound(s) is considered **investigational** for all indications because the evidence is insufficient to determine the technology results in an improvement in the net health outcomes.

POLICY GUIDELINES

Coding

See the [Codes table](#) for details.

BACKGROUND

According to the Centers for Disease Control and Prevention (CDC) (CDC, 2023) in the United States, annually, there are 3% of newborns with birth defects that are a major cause of infant mortality. They contribute to 20% of all infant deaths.

Ultrasound is the transmission of high-frequency sound waves through tissues of varying densities. The echoes produced by the high-frequency sound waves at interfaces between tissues and reflect off the body to make visual images. Images created by the echoes of the sound waves are transmitted from the transducer to a CRT or television monitor. The most common frequencies of sound waves used in OB/GYN ultrasound are 2–5 Mhz. A two-dimensional (2-D) ultrasound is most widely used due to its non-invasive nature. The images created by the 2-D ultrasounds are black-and-white, flat and single-planed. 2-D ultrasounds provide a cross-sectional image which may some argue reduce the diagnostic accuracy thus, four-dimensional (4-D) and five-dimensional (5-D) ultrasounds have been proposed to be used in feal ultrasounds.

Four-Dimensional (4-D) Ultrasound

Four-dimensional (4-D) ultrasonography also known as dynamic 3-D sonography refers to real-time visualization of 3-D images. The time vector (the fourth dimension) makes it possible to perceive a rapid update of the successive individual images displayed on the monitor at very short intervals or a time-lapse which creates the impression of real-time images showing fetal movement and expressions.

Five-Dimensional (5-D) Ultrasound

Five-dimensional (5-D) ultrasonography builds upon 4-D sonography, automating the process of acquiring diagnostic images based upon volume data with a software package using artificial intelligence. This improves the detail and quality of the image, improves efficiency, and reduces human error.

The 5-D technology includes:

- 5-D Heart Color: This automatically displays nine standard fetal echocardiography views with blood flow dynamics simultaneously in a single template. The intuitive workflow can simplify examination of the fetal heart and reduce operator dependency.
- 5-D CNS+: This provides nine planes (axial, coronal, sagittal planes) of the fetal brain with anatomical landmarks and biometric measurements. The 5-D CNS+ combines clinical knowledge- based cues with pattern classification algorithms to determine the best standardization planes that are clinically significant. It complies with the International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) guideline for the fetal brain.
- 5-D Limb Vol: This technology provides an efficient way to rapidly measure fractional limb volume. This soft tissue parameter can be added to conventional 2-D ultrasound measurements of the fetal head (BPD) and abdomen (AC) to improve the precision of estimated fetal weight (EFW) and nutritional status. This computer assisted technology has clinical potential to detect and monitor malnourished fetuses with growth abnormalities.
- 5-D NT: Offers nuchal translucency measurement solutions for first trimester fetal nuchal translucency measurements.
- 5-D LB: Offers intuitive detection and measurement of fetal long bones.

5-D technology in fetal assessment is utilized in clinical practice with the following:

- Biometrics to measure biparietal diameter and crown-rump length and determine gestational age.
- Nuchal translucency - manual measurement, semi-automatic; 5-D recognizes the correct mid-sagittal plane and provides improved Herman score.

- Morphological assessment – 3-D and 4-D enhancements offer more capabilities for accurate assessment to aid diagnosis of visible anomalies, invisible anomalies and anomalies requiring analysis: cardiac, face and limbs, spina bifida.
- Diagnosis of chorionicity and aminiocity in twin pregnancies.
- Fetal risk assessment – characterizes risk that include aneuploidies, congenital heart defects, and spina bifida.

Regulatory Status

An ultrasound is a procedure and, therefore, not subject to U.S. Food and Drug Administration (FDA) regulation. However, any medical devices used as a part of this procedure may be subject to FDA regulation. Many devices for use in ultrasound are available. These devices have FDA clearance under product codes IYN, ITX, and IYO, for marketing in the United States.

The FDA recommends that health care providers consider ways to minimize exposure while maintaining diagnostic quality when using ultrasound. As with all other imaging modalities, the principles of As Low As Reasonably Achievable (ALARA) should be practiced by health care providers.

The FDA reports “the use of ultrasound solely for non-medical purposes such as obtaining fetal ‘keepsake’ videos has been discouraged.” They report. “While ultrasound is generally considered to be safe with very low risks, the risks may increase with unnecessary prolonged exposure to ultrasound energy, or when untrained users operate the device.” Refer to the following for more information: [Ultrasound Imaging | FDA](#).

Please note this section is not intended to be all-inclusive.

RATIONALE

This evidence review was created in April 1999 and has been updated regularly with searches of the PubMed database. The most recent literature update was performed through January 2026.

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function, including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Four - Dimensional (4-D) and Five - Dimensional (5-D) Ultrasound(s) for the Diagnosis of Fetal Cardiac Abnormalities

Clinical Context and Test Purpose

The purpose of four-dimensional (4-D) and five-dimensional (5-D) ultrasound for the diagnosis of fetal cardiac abnormalities is to provide an alternative to or an improvement to existing therapies such as a 2-D ultrasound.

Populations

The relevant population of interest are individuals who are pregnant.

Interventions

The therapy being considered is four-dimensional (4-D) and five-dimensional (5-D) ultrasound(s) for the diagnosis on fetal cardiac abnormalities.

Comparators

Comparators of interest is a two-dimensional (2-D) ultrasound.

Outcomes

The general outcomes of interest test accuracy related to identification of fetal abnormalities, overall survival (OS), and adverse events.

Review of Evidence

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

A January 2024 Hayes Evidence Analysis Research Brief on Four-Dimensional and Five-Dimensional Ultrasound for Diagnosis of Fetal Cardiac Abnormalities identified 2 Class I studies of Clinical Validity (Turan et al, 2014; Wang et al, 2019) that includes a comparison reference test. All evaluated the diagnostic accuracy of 4-D ultrasound. None evaluated 5-D ultrasound-US. One was conducted in the United States. Findings from these studies are inconclusive as they were heterogenous in the types of fetal abnormalities evaluated and they did not provide adequate details on variance of effect to assess level of precision.

Observational Studies

Wang et al (2019) conducted a diagnostic accuracy study on the identification of cardiac malformations of a fetus by both a 2-D ultrasound and 4-D ultrasound. A total of 206 high-risk individuals analyzed retrospectively analyzed. The two-dimensional ultrasounds identified 100 cardiac malformations. The four-dimensional ultrasound identified 120 cardiac malformations. When both 2-D and 4-D ultrasounds were used 135 cardiac malformations were diagnosed. "The sensitivity, specificity, diagnostic coincidence rate, negative predictive value, and positive predictive value of two-dimensional ultrasound diagnosis were 70.92, 78.46, 73.30, 55.43 and 87.72%, respectively; the sensitivity, specificity, diagnostic coincidence rate, negative predictive value, and positive predictive value of four-dimensional ultrasound diagnosis were 85.11, 89.23, 86.41, 73.42 and 94.49%, respectively; the sensitivity, specificity, diagnostic coincidence rate, negative predictive value and positive predictive value of two-dimensional ultrasound diagnosis combined with four-dimensional ultrasound diagnosis were 95.74, 67.69, 86.89, 88.00 and 86.54%, respectively. The sensitivity of two-dimensional ultrasound diagnosis combined with four-dimensional ultrasound diagnosis was significantly higher than that of two-dimensional ultrasound diagnosis and four-dimensional ultrasound diagnosis, the difference was statistically significant ($P<0.05$). The sensitivity of four-dimensional ultrasound diagnosis was significantly higher than that of two-

dimensional ultrasound diagnosis, the difference was statistically significant ($P < 0.05$). The specificity and positive predictive value of four-dimensional ultrasound diagnosis were significantly higher than those of two-dimensional ultrasound diagnosis and two-dimensional ultrasound diagnosis combined with four-dimensional ultrasound diagnosis, the difference was statistically significant ($P < 0.05$)." While a combination of 2-D and 4-D ultrasound may be beneficial in screening for fetal CMs, more research is indicated to analyze the diagnostic value.

Wang et al. (2025) conducted a diagnostic accuracy study evaluating the identification of fetal cardiac malformations using two-dimensional ultrasound (2D-CDU), four-dimensional ultrasound with spatiotemporal image correlation (4D-STIC), and a combined approach in 135 high-risk pregnant women at a single center in Zhejiang Province, China. High-risk factors were reported as age ≥ 35 , drinking during pregnancy, and a history of adverse pregnancy outcomes as independent risk factors for fetal cardiac malformations. A total of 40, 38, and 42 fetal cardiac malformations were detected by 2D-CDU, 4D-STIC, and the combined method, respectively. Reported sensitivity was highest for the combined approach (93.3%; 95% CI 80.7–98.3), compared with 2D-CDU (88.9%; 95% CI 75.2–95.8) and 4D-STIC (84.4%; 95% CI 69.9–93.0). Specificity was also highest for the combined approach (95.6%; 95% CI 88.3–98.6) and was similar for 4D-STIC alone (94.4%; 95% CI 86.9–97.9) compared to lower specificity for 2D-CDU alone (83.3%; 95% CI 73.7–90.1), with overlapping confidence intervals indicating no clear difference between these modalities. Limitations include small sample size, wide confidence intervals, lack of direct quantitative comparison of 4D vs 2D ultrasound, and limited generalizability due to conduct at a single non-US center that suggests the need for further research.

Turan et al (2014) conducted a diagnostic accuracy study on utilizing 4D ultrasound/echocardiogram in pregnant individuals who are at high risk of carrying a fetus with congenital heart disease. Abnormalities were detected in 20 fetuses, most commonly an atrioventricular canal defect ($n=9$). The first trimester scan missed two CSH cases. Those two cases were caught on a second trimester scan. The "first-trimester echocardiography scan showed a sensitivity of 91% (95% CI, 71–99%), a specificity of 100% (95% CI, 97–100%), a positive predictive value of 100% (95% CI, 83–100%) and a negative predictive value of 99% (95% CI, 95–100%)." A first trimester scan is limited by overall image resolution, gestational age, and identifiable overall small anatomical size and landmarks. A sonographer in a general office setting is likely to have a skill set that differs than those who were included in this study. Further research is needed to decipher how long it takes to acquire these skills, interpret the findings, or have analysis online. Bias may exist since this study was completed at a single center and the STIC operator is not blinded. While 4-Dechocardiograms may be beneficial in screening high risk individuals for a fetus effected by CDH further research is indicated to analyze the effectiveness.

Section Summary: Clinically Valid

Four-Dimensional and Five-Dimensional Ultrasound for Diagnosis of Fetal Cardiac Abnormalities identified 2 Class I studies of Clinical Validity (Wang et al., 2019; Turan et al. 2014). All evaluated the diagnostic accuracy of 4-D ultrasound. None evaluated 5-D ultrasound. One study was conducted in the United States. Findings from these studies are inconclusive as they were heterogenous in the types of fetal abnormalities evaluated and they did not provide adequate details on variance of effect to assess level of precision.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if individuals receive the correct care, more effective care, or avoid unnecessary therapy or testing.

There were no clinical utility studies that evaluated 4-D and 5-D ultrasounds for the diagnosis of fetal cardiac abnormalities identified.

A January 2024 Evidence Analysis Research Brief (EARB) by Hayes, a symplr company, regarding Four-Dimensional and Five-Dimensional Ultrasound for Diagnosis of Fetal Cardiac Abnormalities did not identify any clinical utility studies.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials (RCTs).

No randomized or nonrandomized controlled studies were identified that compared health outcomes in individuals when treatment decisions were made with and without the results of four - dimensional (4-D) and five - dimensional (5-D) ultrasound(s) for the diagnosis of fetal cardiac abnormalities.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Section Summary: Clinically Useful

Direct evidence of how 4-D and 5-D fetal ultrasounds for the diagnosis of fetal cardiac abnormalities to improve outcomes is lacking. In the absence of direct evidence for a diagnostic test, a chain of evidence can sometimes be identified to demonstrate improvement in health outcomes. However, in the case of 4-D and 5-D ultrasound(s) for the diagnosis of fetal cardiac abnormalities, the chain of evidence about clinical validity and how the test would be used in practice is uncertain; therefore, no inferences can be made about clinical utility.

Four - Dimensional (4-D) and Five - Dimensional (5-D) Ultrasound(s) for the Diagnosis of Fetal Noncardiac Abnormalities

Clinical Context and Test Purpose

The purpose of four-dimensional (4-D) and five-dimensional (5-D) ultrasound for the diagnosis of fetal noncardiac abnormalities is to provide an alternative to or an improvement to existing therapies such as a 2-D ultrasound.

Populations

The relevant population of interest are individuals who are pregnant.

Interventions

The therapy being considered is four-dimensional (4-D) and five-dimensional (5-D) ultrasound(s) for the diagnosis on fetal noncardiac abnormalities.

Noncardiac anomalies may include facial and oral deformities, genetic disorders, limb malformations, neural tube defects, spinal irregularities and others (CDC, 2023; Jabaz and Jenkins, 2023).

Comparators

Comparators of interest is a two-dimensional (2-D) ultrasound.

Outcomes

The general outcomes of interest test accuracy related to identification of fetal abnormalities, overall survival (OS), and adverse events.

Review of Evidence

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

A December 2023 Hayes Evidence Analysis Research Brief on Four-Dimensional and Five-Dimensional Ultrasound for Diagnosis of Fetal Noncardiac Abnormalities identified 4 Class I studies of Clinical Validity (Öcal et al, 2015; Wang et al, 2019; Yu et al, 2022; Zhang et al, 2023). All evaluated the diagnostic accuracy of 4-D ultrasound. None evaluated 5-D ultrasound. None were conducted in the United States. Findings from these studies are inconclusive as they were heterogenous in the types of fetal abnormalities evaluated and they did not provide adequate details on variance of effect to assess level of precision.

Observational Studies

Zhang et al (2023) conducted a diagnostic accuracy study on the GE-E10 four-dimensional (4-D) ultrasound to investigate the efficacy of GE-E10 prenatal ultrasounds in forecasting fetal abnormal weight development. 160 pregnant women were included in this study. All women had both a two-dimensional (2-D) and (4-D) ultrasound. "Sensitivity and specificity of 2-D color ultrasound in diagnosing fetal abnormal development were 78.38% and 82.60%. The sensitivity and specificity of 4-D color ultrasound in diagnosing fetal abnormal weight development were 81.15% and 83.43%. Receiver operating characteristic showed that the area under the curve (0.873) of 4-D color ultrasound was higher than that 2-D color ultrasound (0.827)." However, a limitation of this study is that the clinical and statistical significance of these findings are unclear as they were not reported. Although the results suggest the GE-E10 (4-D) ultrasound was more diagnostic than 2-D ultrasounds for value for antenatal screening of macrosomia and low birth weight. Zhang et al also referenced Eslamian et al. (2018) which reported the results had increased accuracy when the proximity of date of birth and the fetal weight date were closer together. When the timing of the date of birth was off the accuracy of the fetal weight estimate was not as reliable. Thus, the application of the 4-D ultrasound fetal weight predication "is not realistic" and providers should have careful consideration of effective timing and utilization of the 4-D fetal ultrasound in predicting fetal weight. Overall, the authors concluded the 4-D ultrasound has a "high value for antenatal screening of macrosomia and low birth weight" however, the 4-D ultrasound results were not statistically significant when compared to the 2-D ultrasound.

Yu et al (2022) evaluated the diagnostic accuracy of two-dimensional ultrasonography (2-D ultrasound) combined with four-dimensional ultrasonography (4-D ultrasound) in prenatal ultrasound screening of fetal congenital malformations (CMs) and explores the high-risk factors affecting fetal malformations. From February 2020 to October 2021, 2247, pregnant individuals completed a 2-D ultrasound. Of those 2247 those whose suspected fetal malformations were further examined with a 4-D ultrasound. "The accuracy, sensitivity, and specificity of 2D-US diagnosis were 81.40%, 43.68%, and 82.92%". For the 4-DD ultrasound the accuracy, sensitivity, and specificity of diagnosis were, "83.67%, 51.72%, and 84.95%". For 2-D and 4-D ultrasound combined accuracy, sensitivity, and specificity diagnosis was statistically higher than 2-D or 4-D ultrasound alone with 93.59%, 90.80%, and 91.70%, respectively. Although this study does highlight the combination of 2-D and 4-D ultrasound may increase the diagnosis rate of fetal malformations there are several limitations to consider. Results for 2-D and 4-D ultrasound were not stratified per the gestational week. Individuals were only studied during the second trimester. Additional

risk factors which may affect fetal congenital malformations were not evaluated such as the male counterpart's smoking or drinking history, genetic diseases, and radiation. Additionally, the clinical and statistical significance of these findings are unclear as they were not reported. While a combination of 2-D and 4-D ultrasound may be beneficial in screening for fetal CMs, more research is indicated to analyze the diagnostic value.

Wang et al (2019) evaluated the diagnostic accuracy of two-dimensional (2-D) plus four-dimensional (4-D) ultrasonography in diagnosis of fetal craniocerebral anomalies. They retrospectively reviewed 83 individuals from January 2013 to December 2017 who had been suspected of fetal craniocerebral anomalies from 2-D and 4-D ultrasound. 2-D ultrasound only was used in 56 patients, 4-D ultrasound only was used in 65 and 2-D plus 4-D ultrasound was used in 74 individuals with identified anomalies. Diagnostic accuracy of 2-D ultrasound only was 68.67%, 4-D ultrasound only was 81.93% and 2-D plus 4-D ultrasound was 95.18% ($P < 0.05$). However, a limitation of this study is that the clinical and statistical significance of these findings are unclear as they were not reported. The accuracy, sensitivity, and specificity of 2-D plus 4-D ultrasound was greater than those of 2-D ultrasound only and 4-D ultrasound only, and the accuracy of 4-D ultrasound only was higher than that of 2-D ultrasound only ($P < 0.05$). Although the results from this study indicate the 2-D plus 4-D ultrasound and the 4-D ultrasound alone is more statistically significantly diagnostic for various fetal craniocerebral anomalies when compared to the 2-D ultrasound alone, however there are limitations of this study. The study design included a small sample size with limited follow-up. Additional, well-designed studies are indicated.

Öcal et al (2015) evaluated the diagnostic accuracy of 4-D ultrasounds in the detection of fetal abnormalities. In 1,379 individuals, 2-D and 4-D fetal ultrasounds were completed in the same visit. A total of 176 pregnant individuals had 194 fetal anomalies. The authors concluded that 2-D ultrasounds were superior at detecting anomalies ($p < 0.001$). In approximately half of the cases 4-D ultrasounds identified the fetal abnormalities and 15% of cases there was enhanced image quality.

Section Summary: Clinically Valid

Four-Dimensional and Five-Dimensional Ultrasound for Diagnosis of Fetal Noncardiac Abnormalities identified 4 Class I studies of Clinical Validity (Öcal et al, 2015; Wang et al, 2019; Yu et al, 2022; Zhang et al, 2023). All evaluated the diagnostic accuracy of 4-D ultrasound. None evaluated 5-D ultrasound. None were conducted in the United States. Findings from these studies are inconclusive as they were heterogeneous in the types of fetal abnormalities evaluated and they did not provide adequate details on variance of effect to assess level of precision.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if individuals receive the correct care, more effective care, or avoid unnecessary therapy or testing.

There were no clinical utility studies that evaluated 4-D and 5-D ultrasounds for the diagnosis of fetal noncardiac abnormalities identified.

A December 2023 Evidence Analysis Research Brief (EARB) by Hayes regarding Four-Dimensional and Five-Dimensional Ultrasound for Diagnosis of Fetal Noncardiac Abnormalities did not identify any clinical utility studies.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials (RCTs).

No randomized or nonrandomized controlled studies were identified that compared health outcomes in individuals when treatment decisions were made with and without the results of four - dimensional (4-D) and five - dimensional (5-D) ultrasound(s) for the diagnosis of fetal noncardiac abnormalities.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Section Summary: Clinically Useful

Direct evidence of how 4-D and 5-D fetal ultrasounds for the diagnosis of fetal noncardiac abnormalities to improve outcomes is lacking. In the absence of direct evidence for a diagnostic test, a chain of evidence can sometimes be identified to demonstrate improvement in health outcomes. However, in the case of 4-D and 5-D ultrasound(s) for the diagnosis of fetal noncardiac abnormalities, the chain of evidence about clinical validity and how the test would be used in practice is uncertain; therefore, no inferences can be made about clinical utility.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Obstetricians and Gynecologists (ACOG)

In 2017 ACOG issued practice bulletin No. 723 Guidelines for Diagnostic Imaging During Pregnancy and Lactation which was reaffirmed 2021. This guideline does not specifically discuss the use of 4-D or 5-D ultrasound.

ACOG issued practice bulletin No. 175 Ultrasound in Pregnancy in 2016 which did not specifically discuss the use of 4-D or 5-D ultrasound.

American Heart Association (AHA)

In 2014 the AHA released a Scientific Statement on the Diagnosis and Treatment of Fetal Cardiac Disease which stated, "3-D/4-D fetal cardiac imaging is currently a research tool and is not adequate for use as an alternative to conventional fetal cardiac imaging. However, this technology may be useful to facilitate screening for congenital heart disease (CHD) or for complementary imaging in fetuses identified as having CHD" (IIb/B).

This statement does not specifically mention 5-D ultrasounds.

American Institute of Ultrasound in Medicine (AIUM)

The AIUM provides a recommendation on Keepsake Fetal Imaging in 2020 in the statement on Prudent Use and Safety of Diagnostic Ultrasound in Pregnancy which, “encourages patients to make sure that practitioners using ultrasound have received formal education and training in fetal imaging to ensure the best possible results.

The AIUM recognizes the growing pressures from patients for the performance of ultrasound examinations for bonding and reassurance purposes largely driven by advances in image quality of 3-dimensional (3-D) sonography and by more widely available information about these advances. Although there is only some scientific evidence that 3-D sonography has a positive impact on parental-fetal bonding, the AIUM recognizes that many parents may pursue scanning for this purpose. Such “keepsake imaging” currently occurs in a variety of settings, including the following:

1. Images or video clips given to parents during the course of a medically indicated ultrasound examination.
2. Images or clips given to volunteers who are scanned as part of diagnostic ultrasound education programs or demonstrations, provided that images are not used as an enticement to participate.
3. Freestanding commercial fetal imaging sites, usually without any physician review of acquired images and with no regulation of the training of the individuals obtaining the images; these sites are sometimes called “baby video studios,” and these videos are sometimes called “entertainment videos.”
4. As added-cost visits to a medical facility (office or hospital) outside the coverage of contractual arrangements between the provider and the patient’s insurance carrier. The AIUM believes that added-cost arrangements other than those for providing patients images or copies of their medical records at cost may violate the principles of medical ethics of the American Medical Association (AMA) and the American College of Obstetricians and Gynecologists (ACOG).

The AIUM, therefore, recommends that only scenarios 1 and 2 above are consistent with the ethical principles of the AIUM and those of the AMA and ACOG.

The market for keepsake images is driven in part by past medical approaches that have used medicolegal concerns as a reason not to provide images to patients. Sharing images with patients is unlikely to have a detrimental medicolegal impact. The AIUM encourages sharing images with patients as appropriate when medically indicated obstetric ultrasound examinations are performed.”

This statement does not specifically mention 4-D and 5-D ultrasounds.

American Institute of Ultrasound in Medicine (AIUM) - American College of Radiology (ACR) – American College of Obstetricians and Gynecologists (ACOG) - Society of Radiologist in Ultrasound (SRU)

In 2018 AIUM – ACR – ACOG and SRU issued a collaborative practice parameter for the performance of standard diagnostic obstetrical ultrasound examination that was revised in 2023 and does not specifically discuss the use of 4-D or 5-D ultrasound.

American Society of Echocardiography (ASE)

In 2023 the ASE provided a Guideline and Recommendation for the Performance of the Fetal Echocardiogram stated, ““Other ultrasound technologies may be used to image fetal cardiovascular structure and physiology. [STIC] captures a static or dynamic [3-D] volume data set using a specially designed ultrasound transducer and analysis software. Real-time [3-D] echocardiographic imaging with this method can be used to enhance detection of anatomic defects and quantify hemodynamics such as ventricular function and cardiac output, although [STIC] has not been validated for clinical use.”

This statement does not specifically mention 5-D ultrasounds.

International Society of Ultrasound in Obstetrics and Gynecology (ISUOG)

Performance of 11–14-week Ultrasound Scan

The ISUOG updated their Practice Guideline on the Performance of 11–14-week Ultrasound Scan in 2023; For the Role of 3-D and 4-D ultrasound they stated “3-D and 4-D ultrasound are not currently used for routine first-trimester fetal anatomical evaluation. However, in experienced hands, these methods may be helpful in evaluation of abnormalities, especially with multiplanar reconstruction of selected diagnostic planes”.

This statement does not specifically mention 5-D ultrasounds.

Performance of the Routine Mid-Trimester Fetal Ultrasound Scan

The ISUOG updated their Practice Guidelines on the Performance of the Routine Mid-Trimester Fetal Ultrasound Scan which does not specifically discuss the use of 4-D or 5-D ultrasound.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review can be located at clinicaltrials.gov.

REFERENCES

1. U.S. Food and Drug Administration. Radiation Emitting Products. Ultrasound Imaging. Updated September 28, 2020. Available at: <http://www.fda.gov/Radiation-EmittingProducts/RadiationEmittingProductsandProcedures/MedicalImaging/ucm115357.htm>. Accessed January 6, 2026.
2. U.S. Food and Drug Administration. Avoid Fetal “Keepsake” Images, Heartbeat Monitors. December 16, 2014. Available at: <https://www.fda.gov/ForConsumers/ConsumerUpdates/ucm095508.htm>
3. Whitworth M, Bricker L, Neilson JP. Et. al. Ultrasound for fetal assessment in pregnancy. Cochrane Database Syst Rev. 2010 Apr 14; (4): CD007058. PMID 20393955
4. Whitworth M, Bricker L, Mullan C. Ultrasound for fetal assessment in early pregnancy. Cochrane Database Syst Rev 2015 Jul 14;7: CD007058. PMID 2617896
5. Benacerraf BR, Shipp TD, Bromley B. Three-dimensional US of the fetus: volume imaging. Radiology 2006 Mar 238(3):988-96. PMID 16424249
6. Yagel S, Cohen SM, Messing B, et. al. Three-dimensional and four-dimensional ultrasound application in fetal medicine. Curr Opin Obstet Gynecol 2009 Apr;21(2):167-74. PMID 19996869
7. Goncalves LF, Lee W, Espinoza J, Romero R. Three and 4-dimensional ultrasound in obstetric practice: does it help? J Ultrasound Med. 2005 Dec
8. Hur H, Kim YH, Cho Hy et. al. Feasibility of three-dimensional reconstruction and automated measurement of fetal long bones using 5D long bone. Obstet Gynecol Sci 2015;58(4):268-276
9. Laban M, Alanwar AA, Etman MK, et. al. Five-dimensional long bones biometry for estimation of femur length and fetal weight at term compared to two-dimensional ultrasound: a pilot study. J Matern Fetal Neonatal Med. 2017:1-7

10. Rizzo G, Capponi A, Persico N, et al. 5D CNS+ software for automatically imaging axial, sagittal and coronal planes of normal and abnormal second trimester fetal brains. *J Ultrasound Med* 2016;35(10):2263-2272
11. Martins JG, Biggio JR, Abuhamad A. Society for Maternal-Fetal Medicine (SMFM) Consult Series #52: Diagnosis and Management of Fetal Growth Restriction. *American Journal of Obstetrics and Gynecology*. 2020. doi:10.1016/j.ajog.2020.05.010.
12. AboEllail MAM, Kanenishi K, Mori N, et al. 4D ultrasound study of fetal facial expressions in the third trimester of pregnancy. *J Matern Fetal Neonatal Med*. 2018; 31(14):1856-1864.
13. Hur H, Kim YH, Cho HYet al. Feasibility of three-dimensional reconstruction and automated measurement of fetal long bones using 5D Long Bone. *Obstet Gynecol Sci*. 2015; 58(4):268-276.
14. Kurjak A, Azumendi G, Andonotopo W, Salihagic-Kadic A. Three- and four-dimensional ultrasonography for the structural and functional evaluation of the fetal face. *Am J Obstet Gynecol*. 2007; 196(91):16-28.
15. Laban M, Alanwar AA, Etman MK, et al. Five-dimensional long bones biometry for estimation of femur length and fetal weight at term compared to two-dimensional ultrasound: a pilot study. *J Matern Fetal Neonatal Med*. 2018; 31(15):2036-2042.
16. Yagel S, Cohen SM, Messing B. 3D and 4D ultrasound in fetal cardiac scanning: a new look at the fetal heart. *Ultrasound Obstet Gynecol*. 2007; 29(1):81-95.
17. Hata T, Tanaka H, Noguchi J. 3D/4D sonographic evaluation of amniotic band syndrome in early pregnancy: A supplement to 2D ultrasound. *J Obstet Gynaecol Res*. 2011;37(6):656-660.
18. Kanenishi K, Hanaoka U, Noguchi J, et al. 4D ultrasound evaluation of fetal facial expressions during the latter stages of the second trimester. *Int J Gynaecol Obstet*. 2013;121(3):257-260.
19. Votino C, Cos T, Abu-Rustum R, et al. Use of spatiotemporal image correlation at 11-14 weeks' gestation. *Ultrasound Obstet Gynecol*. 2013;42(6):669-678.
20. Ahmed BI. The new 3D/4D based spatio-temporal imaging correlation (STIC) in fetal echocardiography: A promising tool for the future. *J Matern Fetal Neonatal Med*. 2014;27(11):1163-1168.
21. Jabaz D, Jenkins SM. Sonography 2nd Trimester Assessment, Protocols, and Interpretation. [Updated 2023 Nov 12]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK570574/> Accessed January 6, 2026.
22. Eslamian ZL, Zarean E, Moshfeghi M, et al. Evaluation of the predictive value of fetal Doppler ultrasound for neonatal outcome from the 36th week of pregnancy. *J Res Med Sci*. 2018 Feb 20;23:13. doi: 10.4103/jrms.JRMS_133_17. PMID: 29531565; PMCID: PMC5842445.
23. Zhang L, Jin TL. et al. Predictive value and accuracy of prenatal four-dimensional color ultrasound for fetal abnormal development. *Medicine (Baltimore)*. 2023 Aug 11;102(32):e34553. doi: 10.1097/MD.00000000000034553. PMID: 37565886; PMCID: PMC10419341.
24. Yu X, Liu F, Gao W, et al. Diagnostic Value and High-Risk Factors of Two-Dimensional Ultrasonography Combined with Four-Dimensional Ultrasonography in Prenatal Ultrasound

- Screening of Fetal Congenital Malformations. *Comput Math Methods Med*. 2022 Jul 12;2022:7082832. doi: 10.1155/2022/7082832. PMID: 35866037; PMCID: PMC9296308.
25. Wang Y, Chen X, Zhong S, et al. Diagnostic value of two-dimensional plus four-dimensional ultrasonography in fetal craniocerebral anomalies. *Iran J Public Health*. 2019;48(2):323-330.
 26. Kim S.H., Choi B.I., Three-dimensional and Four-dimensional Ultrasound: Techniques and Abdominal Applications, *Journal of Medical Ultrasound*, Volume 15, Issue 4, 2007, P 228-242. Available at: [https://doi.org/10.1016/S0929-6441\(08\)60040-5](https://doi.org/10.1016/S0929-6441(08)60040-5). Accessed January 6, 2026.
 27. Merz E., Abramwicz J., 3D/4D Ultrasound in Prenatal Diagnosis: Is it Time for Routine Use?. *Clinical Obstetrics and Gynecology* 55(1):p 336-351, March 2012. | DOI: 10.1097/GRF.0b013e3182446ef7.
 28. Tonni G, Grisolia G, Santana EF, et al. Assessment of fetus during second trimester ultrasonography using HDlive software: What is its real application in the obstetrics clinical practice? *World J Radiol* 2016; 8(12): 922-927 PMID: 28070244 DOI: 10.4329/wjr.v8.i12.922.
 29. Jost E, Kosian P, Jimenez Cruz J, et al. Evolving the Era of 5D Ultrasound? A Systematic Literature Review on the Applications for Artificial Intelligence Ultrasound Imaging in Obstetrics and Gynecology. *Journal of Clinical Medicine*. 2023; 12(21):6833. <https://doi.org/10.3390/jcm12216833> Accessed January 6, 2026.
 30. Inubashiri E, Fujita S, Shimakura S, et al. A new approach for quantitative assessment of fetal general movements in the early second trimester of pregnancy using four-dimensional ultrasound. *J Med Ultrason* (2001). 2021 Jul;48(3):335-344. doi: 10.1007/s10396-021-01095-1. Epub 2021 Apr 27. PMID: 33907961.
 31. Zhang D, Zhang Y, Ren W, et al. Prenatal diagnosis of fetal interrupted aortic arch type a by two-dimensional echocardiography and four-dimensional echocardiography with b-flow imaging and spatiotemporal image correlation. *Echocardiography*. 2016;33(1):90-98. doi:10.1111/echo.12996
 32. Li H, Peng F, Wu C, Kong D, Zhang Q, Zhang Z. Diagnostic value of four-dimensional ultrasonography with STIC combined with two-dimensional ultrasonography for fetal cardiac malformation and chromosomal abnormalities in early pregnancy. *Exp Ther Med*. 2020;19(2):1161-1166. doi:10.3892/etm.2019.8325
 33. Tongsong T, Tongprasert F, Srisupundit K, Luewan S, Traisrisilp K. Cardio-STIC (spatiotemporal image correlation) as genetic ultrasound of fetal down syndrome. *J Matern Fetal Neonatal Med*. 2015;28(16):1943-1949. doi:10.3109/14767058.2014.973395
 34. Turan S, Turan OM, Desai A, Harman CR, Baschat AA. First-trimester fetal cardiac examination using spatiotemporal image correlation, tomographic ultrasound and color doppler imaging for the diagnosis of complex congenital heart disease in high-risk patients. *Ultrasound Obstet Gynecol*. 2014;44(5):562-567. doi:10.1002/uog.13341
 35. Wang B, Li J, Yin J. Diagnostic value of echocardiography in fetal cardiac malformation and clinical classification. *Exp Ther Med*. 2019;18(3):1595-1600. doi:10.3892/etm.2019.7732
 36. The American College of Obstetricians and Gynecologists (ACOG) Practice Bulletin 101. *Ultrasonography in Pregnancy*. *Obstet Gynecol*. 2009; 113(2 Pt 1):451-461
 37. ACR-ACOG-AIUM-SRU Practice Parameter for the Performance of Obstetrical Ultrasound, Amended 2014.

38. AIUM, ACR, ACOG, SMFU, SRU Practice Parameter for Performance of Standard Diagnostic Obstetric Ultrasound Examinations. J Ultrasound Med 2018; 9999:1-12
39. AIUM Practice Parameter for the Performance of Detailed Second- and Third-Trimester Diagnostic Obstetric Ultrasound Examinations. Journal of Ultrasound in Medicine. 2019;38(12):3093-3100. doi:10.1002/jum.15163
40. ACOG Practice Bulletin No. 226: Screening for Fetal Chromosomal Abnormalities. Obstetrics & Gynecology. 2020;136(4):e48-e69. doi:10.1097/aog.0000000000004084.
41. ACOG Practice Bulletin No. 229: Indications for Outpatient Antenatal Fetal Surveillance. Obstetrics & Gynecology. 2021;137(6):e177-e197. doi:10.1097/aog.0000000000004407. Available at: <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2021/06/indications-for-outpatient-antenatal-fetal-surveillance> Accessed January 6, 2026.
42. American Academy of Obstetricians and Gynecologists (ACOG) Practice Bulletin No. 175 Ultrasound in Pregnancy. Obstetrics and Gynecology December 2016 Volume 128 Issue 6 p e241-e256..
43. ACOG Practice Bulletin No. 227: Fetal Growth Restriction. Obstet Gynecol. 2021;137(2):e16-e28 doi: 10.1097/AOG.0000000000004251 Galan HL. Timing Delivery of the Growth-Restricted Fetus. Seminars in Perinatology. 2011;35(5):262-269. doi:10.1053/j.semperi.2011.05.009. Available at: <https://files.medelem.com/uploads/materials/f17f05127e54e357c2accd8c6ac8ce6b.pdf> Accessed January 6, 2026.
44. Bilardo C.M., Chaoui R., Hyett A, et al. ISUOG Practice Guidelines (updated): Performance of 11-14-week ultrasound scan. International in Obstetrics & Gynecology. Ultrasound in Obstetrics & Gynecology. Published January 03, 2023. <https://doi.org/10.1002/uog.26106>. Accessed January 6, 2026.
45. National Institute of Biomedical Imaging and Bioengineering. Ultrasound. Last Reviewed December 2023. Available at: [www. https://www.nibib.nih.gov/science-education/science-topics/ultrasound](https://www.nibib.nih.gov/science-education/science-topics/ultrasound). Accessed January 6, 2026.
46. Centers for Disease Control and Prevention (CDC). Data Statistics on Birth Defects. Last reviewed November 19, 2024. Available at: <https://www.cdc.gov/birth-defects/data-research/facts-stats/index.html> Accessed January 6, 2026. .
47. AIUM-ACR-ACOG-SMFM-SRU Practice Parameter for the Performance of Standard Diagnostic Obstetric Ultrasound Examinations.” Journal of ultrasound in medicine: official journal of the American Institute of Ultrasound in Medicine vol. 37,11 (2018): Revised 2023. Available at: <https://onlinelibrary.wiley.com/doi/epdf/10.1002/jum.16406> Accessed January 6, 2026.
48. American Congress of Obstetricians and Gynecologists (ACOG). Committee on Obstetric Practice. Committee Opinion #723. Guidelines for diagnostic imaging during pregnancy and lactation. Reaffirmed 2021. Available at: <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2017/10/guidelines-for-diagnostic-imaging-during-pregnancy-and-lactation>. Accessed January 6, 2026.
49. American Institute of Ultrasound in Medicine (AIUM). Official Statements. Prudent Use and Safety of Diagnostic Ultrasound in Pregnancy. Approved on May 19, 2020. Available at: https://www.aium.org/officialStatements/79?_sw_csrfToken=67ff2ef4. Accessed January 6, 2026.

50. American Congress of Obstetricians and Gynecologists (ACOG). Committee on Obstetric Practice. Practice Bulletin #175. Ultrasound in Pregnancy. 2016. Available at: [https://journals.lww.com/greenjournal/Abstract/2016/12000/Practice Bulletin No 175 Ultrasound in Pregnancy.53.aspx](https://journals.lww.com/greenjournal/Abstract/2016/12000/Practice_Bulletin_No_175_Ultrasound_in_Pregnancy.53.aspx)
51. Monteagudo A., Timor-Tritsch I.E., Wilkins-Haug L., et al. Neural tube defects: Prenatal sonographic diagnosis. UptoDate. Review current through: December 2025. Topic last updated: November 12, 2024. Available at: www.uptodate.com. Accessed January 6, 2026.
52. Abramowicz J.S., Ahn J.T., Levine D., et al. Fetal macrosomia. UptoDate Review current through: December 2025. Topic last updated: March 26, 2024. Available at: www.uptodate.com. Accessed January 6, 2026.
53. Shipp T.D., Simpson L.L., Levine D., et al. Overview of Ultrasound Examination in Obstetrics and Gynecology. Review current through December 2025. Last updated July 16, 2025. Available at: www.uptodate.com. Accessed January 6, 2026.
54. Hayes, a symplr company. Evidence Analysis Research Brief. Four-Dimensional and Five-Dimensional Ultrasound for Diagnosis of Fetal Noncardiac Abnormalities. Hayes, Inc.; December 29, 2023. Available at: www.hayesinc.com. Accessed January 6, 2026.
55. Hayes, Inc. Evidence Analysis Research Brief. Four-Dimensional and Five-Dimensional Ultrasound for Diagnosis of Fetal Cardiovascular Abnormalities. Hayes, Inc.; January 11 2024. Available at: www.hayesinc.com. Accessed January 6, 2026.
56. American Heart Association. Donofrio M.T., Moon-Grady., Hornberger L.K., et al. Diagnosis and Treatment of Fetal Cardiac Disease. A Scientific Statement from the American Heart Association. 24 Apr 2014. Circulation. 2014;129:2183–2242. Available at: <https://www.ahajournals.org/doi/10.1161/01.cir.0000437597.44550.5d>. Accessed January 6, 2026.
57. American Society of Echocardiography. Moon-Grady A.J., Donofrio M.T., Gelehrter S., et al. Guidelines and Recommendations for Performance of the Fetal Echocardiogram: An Update from the American Society of Echocardiography. The Journal of the American Society of Echocardiography. Guidelines and Standards. Volume 36, Issue 7, P679-723. July 2023. Doi:<https://doi.org/10.1016/j.echo.2023.04.014>.
58. Öcal DF, Nas T, Güler I. The place of four-dimensional ultrasound in evaluating fetal anomalies. Ir J Med Sci. 2015 Sep;184(3):607-12. doi: 10.1007/s11845-014-1184-2. Epub 2014 Aug 21. PMID: 25142340.
59. Wang Y, Wang L, Zang H, Wang J, Wu H, Zhao W. Comparison of Different Ultrasonic Screening Methods and Analysis of High Risk Factors for Fetal Cardiac Malformation in Second Trimester of Pregnancy. Pediatr Cardiol. 2025 Apr;46(4):769-777. doi: 10.1007/s00246-024-03525-6. Epub 2024 May 29. PMID: [38809278](https://pubmed.ncbi.nlm.nih.gov/38809278/).

CODES

To report provider services, use appropriate CPT codes, HCPCS codes, Revenue codes, and/or ICD diagnosis codes.

Codes	Number	Description
CPT		
	76499	Unlisted diagnostic radiographic procedure <i>(when specified as a 4-D or 5-D rendering of a fetal ultrasound)</i>
HCPCS		
	No code(s)	
Type of Service	Medical/ Diagnostic	
Place of Service	Outpatient/ Physician's Office	

POLICY HISTORY

Date	Reason	Action
January 2026	Annual Review	Policy Renewed
January 2025	Annual Review	Policy Renewed
January 2024	Annual Review	Policy Revised
June 2023	Annual Review	Policy Revised
June 2022	Annual Review	Policy Revised
June 2021	Annual Review	Policy Renewed
June 2020	Annual Review	Policy Revised
June 2019	Annual Review	Policy Renewed
June 2018	Annual Review	Policy Revised
June 2017	Annual Review	Policy Renewed
June 2016	Annual Review	Policy Revised
July 2015	Annual Review	Policy Renewed

August 2014	Annual Review	Policy Revised
October 2013	Annual Review	Policy Revised
December 2012	Annual Review	Policy Renewed
December 2011	Annual Review	Policy Renewed

New information or technology that would be relevant for Wellmark to consider when this policy is next reviewed may be submitted to:

Wellmark Blue Cross and Blue Shield
Medical Policy Analyst
PO Box 9232
Des Moines, IA 50306-9232

*CPT® is a registered trademark of the American Medical Association.