



Wellmark Blue Cross and Blue Shield is an Independent Licensee of the Blue Cross and Blue Shield Association.

DRUG POLICY

Skyrizi (risankizumab-rzaa)

NOTICE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

BENEFIT APPLICATION

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

DESCRIPTION

The intent of the Skyrizi (risankizumab-rzaa) drug policy is to ensure appropriate selection of patients for therapy based on product labeling, clinical guidelines and clinical studies.

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

1. Treatment of moderate-to-severe plaque psoriasis (PsO) in adults who are candidates for systemic therapy or phototherapy
2. Treatment of active psoriatic arthritis (PsA) in adults
3. Treatment of moderately to severely active Crohn's disease (CD) in adults
4. Treatment of moderately to severely active ulcerative colitis (UC) in adults

POLICY

Required Documentation

Submission of the following information is necessary to initiate the prior authorization review:

- A) Plaque psoriasis (PsO)
 1. Initial requests:
 - i. Chart notes or medical record documentation of affected area(s) and body surface area (BSA) affected (if applicable).

- ii. Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
- 2. Continuation requests: Chart notes or medical record documentation of decreased body surface area (BSA) affected and/or improvement in signs and symptoms.

B) Psoriatic arthritis (PsA)

- 1. Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
- 2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response.

C) Crohn's disease (CD) and ulcerative colitis (UC)

- 1. Continuation requests: Chart notes or medical record documentation supporting positive clinical response to therapy or remission.

Prescriber Specialties (initial approvals only)

This medication must be prescribed by or in consultation with one of the following:

- A. Plaque psoriasis: dermatologist
- B. Psoriatic arthritis: rheumatologist or dermatologist
- C. Crohn's disease and ulcerative colitis: gastroenterologist

Criteria for Initial Approval

A) Moderate to severe plaque psoriasis (PsO)

- 1. Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug (e.g., Sotyktu, Otezla) indicated for the treatment of moderate to severe plaque psoriasis.
- 2. Authorization of 12 months may be granted for adult members for treatment of moderate to severe plaque psoriasis when any of the following criteria is met:
 - a. Crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
 - b. At least 10% of the body surface area (BSA) is affected
 - c. At least 3% of body surface area (BSA) is affected and the member meets either of the following criteria:
 - i. Member has had an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine or acitretin.
 - ii. Member has a clinical reason to avoid pharmacologic treatment with methotrexate, cyclosporine and acitretin (see Appendix A).

B) Active psoriatic arthritis (PsA)

- 1. Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug (e.g., Rinvoq, Otezla) indicated for active psoriatic arthritis.
- 2. Authorization of 12 months may be granted for adult members for treatment of active psoriatic arthritis when either of the following criteria is met:
 - a) Member has mild to moderate disease and meets one of the following criteria:
 - i. Member has had an inadequate response to methotrexate, leflunomide, or another conventional synthetic drug (e.g., sulfasalazine) administered at an adequate dose and duration.

- ii. Member has an intolerance or contraindication to methotrexate or leflunomide (see Appendix), or another conventional synthetic drug (e.g., sulfasalazine).
- iii. Member has enthesitis

b) Member has severe disease.

C) Crohn's disease (CD)

1. Authorization of 12 months may be granted for the treatment of moderately to severely active Crohn's disease.

D) Ulcerative Colitis (UC)

1. Authorization of 12 months may be granted for the treatment of moderately to severely active ulcerative colitis.

Continuation of Therapy

A) Moderate to severe plaque psoriasis (PsO)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderate to severe plaque psoriasis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when either of the following is met:

1. Reduction in body surface area (BSA) affected from baseline
2. Improvement in signs and symptoms from baseline (e.g., itching, redness, flaking, scaling, burning, cracking, pain)

B) Psoriatic arthritis (PsA)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for active psoriatic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of swollen joints
2. Number of tender joints
3. Dactylitis
4. Enthesitis
5. Skin and/or nail involvement
6. Functional status
7. C-reactive protein (CRP)

C) Crohn's Disease (CD)

1. Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active Crohn's disease and who achieve or maintain remission.
2. Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active Crohn's disease and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- i. Abdominal pain or tenderness
- ii. Diarrhea
- iii. Body weight
- iv. Abdominal mass
- v. Hematocrit
- vi. Appearance of the mucosa on endoscopy, computed tomography enterography, (CTE), magnetic resonance enterography (MRE), or intestinal ultrasound
Improvement on a disease activity scoring tool (e.g., Crohn's Disease Activity Index [CDAI] score)

D) Ulcerative colitis

1. Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain remission.
2. Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - i. Stool frequency
 - ii. Rectal bleeding
 - iii. Urgency of defecation
 - iv. C-reactive protein (CRP)
 - v. Fecal calprotectin (FC)
 - vi. Appearance of the mucosa on endoscopy, computed tomography enterography (CTE), magnetic resonance enterography (MRE), or intestinal ultrasound
 - vii. Improvement on a disease activity scoring tool (e.g., Ulcerative Colitis Endoscopic Index of Severity [UCEIS], Mayo score)

Other

For all indications: Member has had a documented negative tuberculosis (TB) test (which can include a tuberculosis skin test [TST] or an interferon-release assay [IGRA])* within 12 months of initiating therapy for persons who are naïve to biologic drugs or targeted synthetic drugs associated with an increased risk of TB.

* If the screening testing for TB is positive, there must be further testing to confirm there is no active disease. (e.g., a chest x-ray) Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug for the same indication.

Skyrizi (risankizumab-rzaa) is considered **not medically necessary** for members who do not meet the criteria set forth above.

Members currently receiving the requested medication as samples or via the manufacturer's patient assistance program will be required to meet the criteria for initial approval. This ensures that members are treated equally regardless of their provider's ability to access medication samples.

Dosage and Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

Quantity Limits Apply

Trade Name	Generic Name	Quantity Limit
Skyrizi™	risankizumab	<p><u>Plaque Psoriasis & Psoriatic Arthritis</u></p> <p>Initiation of therapy: 2 kits (4 x 75mg syringes) per first 28 days (4 weeks) or 2 x 150mg syringes/auto-injectors per first 28 days (4 weeks)</p> <p>Maintenance: 1 kit (2 x 75mg syringes) per 84 days (12 weeks) or 1 x 150mg syringe/auto-injector per 84 days (12 weeks)</p> <p><u>Crohn's Disease</u></p> <p>Initiation of therapy: 3 vials (600mg per vial) per lifetime</p> <p>Maintenance: 1 prefilled cartridge (360 mg/2.4mL) per 56 days (8 weeks)</p> <p><u>Ulcerative Colitis</u></p> <p>Initiation of therapy: 6 vials (600mg per vial) per lifetime</p> <p>Maintenance: 1 prefilled cartridge (360 mg/2.4mL) per 56 days (8 weeks)</p>

APPENDIX

APPENDIX A: Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine, Acitretin, or Leflunomide

1. Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease
2. Drug interaction
3. Risk of treatment-related toxicity
4. Pregnancy or currently planning pregnancy
5. Breastfeeding
6. Significant comorbidity prohibits use of systemic agents (e.g.,liver or kidney disease, blood dyscrasias, uncontrolled hypertension)
7. Hypersensitivity
8. History of intolerance or adverse event

PROCEDURES AND BILLING CODES

To report provider services, use appropriate CPT* codes, Alpha Numeric (HCPCS level 2) codes, Revenue codes, and/or ICD diagnostic codes.

- J2327 - Skyrizi Injection, risankizumab-rzaa, intravenous, 1 mg

REFERENCES

- Skyrizi [package insert]. North Chicago, IL: AbbVie Inc.; June 2024.
- Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 4: Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. *J Am Acad Dermatol.* 2009;61 (3):451-485.
- Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol.* 2011;65(1):137-174.
- Gordon KB, Strober B, Lebwohl M, et al. Efficacy and safety of risankizumab in moderate-to-severe plaque psoriasis (UltIMMA-1 and UltIMMA-2): results from two double-blind, randomised, placebo-controlled and ustekinumab-controlled phase 3 trials. *Lancet.* 2018;392(10148):650-661.
- Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019;80(4):1029-1072.
- Testing for TB Infection. Centers for Disease Control and Prevention. Retrieved on January 11, 2023 from: <https://www.cdc.gov/tb/topic/testing/tbtesttypes.htm>.
- Singh JA, Guyatt G, Oggie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheum.* 2018;71:5-32.
- Gossec L, Baraliakos X, Kerschbaumer A, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. *Ann Rheum Dis.* 2020;79(6):700-712.
- D'Haens G, Panaccione R, Baert F, et al. Risankizumab as induction therapy for Crohn's disease: results from the phase 3 ADVANCE and MOTIVATE induction trials. *Lancet.* 2022;399(10340):2015-2030.
- Lichtenstein GR, Loftus Jr EV, Isaacs KI, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol.* 2018;113:481-517.
- Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021. *Nat Rev Rheumatol.* 2022;18(8):465-479.
- Menter A, Gelfand JM, Connor C, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management of psoriasis with systemic nonbiologic therapies. *J Am Acad Dermatol.* 2020;82(6):1445-1486.
- Talley NJ, Abreu MT, Achkar J, et al. An evidence-based systematic review on medical therapies for inflammatory bowel disease. *Am J Gastroenterol.* 2011;106(Suppl 1):S2-S25.
- Rubin DT, Ananthakrishnan AN, et al. 2019 ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol.* 2019;114:384-413.
- Feuerstein JD, Isaacs KL, Schneider Y, et al. AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology.* 2020;158:1450.

*Some content reprinted from CVSHealth

POLICY HISTORY

Policy #: 05.02.74

Policy Creation: July 2019

Reviewed: April 2025

Revised: April 2025

Current Effective Date: June 6, 2025