

# 04.01.09 MRI-Guided High-Intensity Focused Ultrasound (MRgFUS) Ablation

**Original Effective Date:** August 2007

**Review Date:** September 2025

**Revised:** January 2023

## DISCLAIMER/INSTRUCTIONS FOR USE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations, or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

This Medical Policy document describes the status of medical technology at the time the document was developed. Since that time, new technology may have emerged, or new medical literature may have been published. This Medical Policy will be reviewed regularly and updated as scientific and medical literature becomes available; therefore, policies are subject to change without notice.

---

### Related Policies:

- [02.01.53 High Intensity Focused Ultrasound \(HIFU\)](#)
- [07.01.69 Treatments for Benign Prostatic Hyperplasia \(BPH\)](#)

### Summary

### Description

An integrated system providing magnetic resonance imaging (MRI)-guided focused ultrasound (MRgFUS) treatment is proposed as a noninvasive therapy for uterine fibroids and for pain palliation of bone metastases. MRgFUS is also being investigated for the treatment of other benign and malignant tumors (breast, prostate, brain and desmoid tumors as well as non-spinal osteoid osteoma), as well as essential tremor.

## Summary of Evidence

For individuals who have uterine fibroids who receive magnetic resonance-guided focused ultrasound (MRgFUS), the evidence includes, 2 randomized controlled trials (RCTs), systematic reviews, nonrandomized comparative studies, and case series. Relevant outcomes are symptoms, quality of life, resource utilization, and treatment-related morbidity. One RCT (N=20) has reported some health outcomes but its primary purpose was to determine the feasibility of a larger trial. It did not find statistically significant differences in quality of life outcomes between active and sham treatment groups but it did find lower fibroid volumes after active treatment. This trial did not have an active comparator, the clinical significance of the primary outcome was unclear, and there were no follow-up data beyond 1 year. The second RCT (N=49) had preliminary results at 6 weeks posttreatment, comparing MRgFUS with uterine artery embolization (UAE), and demonstrated that the 2 groups are comparable in medication use and symptom improvement following treatments. Patients in the MRgFUS group reported recovering significantly faster than patients in the uterine artery embolization group, as measured by time to return to work and time to normal activities. Long-term follow-up results reported lower reintervention rate and greater improvement in symptoms after UAE compared to MRgFUS. A 2021 meta-analysis reported that, comparatively, myomectomy had the lowest re-intervention rate of the 3 regimens (myomectomy vs UAE vs MRgFUS) in all time points assessed, while the MRgFUS had the highest re-intervention rate. Long-term data on the treatment effects, recurrence rates, and impact on future fertility and pregnancy are lacking. The evidence is insufficient to determine the effects of the technology on net health outcomes.

For individuals with metastatic bone cancer who have failed or are not candidates for radiotherapy who receive MRgFUS, the evidence includes a sham-controlled randomized trial, a systematic review of RCTs, observational studies, and case series. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and treatment-related morbidity. The RCT found statistically significant improvement after MRgFUS in a composite outcome comprised of reduction in pain and morphine use, and in pain reduction as a stand-alone outcome. A substantial proportion of patients in the treatment group experienced adverse events, but most events were transient and not severe. Pooled efficacy data from a systematic review reported a treatment response to MRgFUS of 79%. The evidence is sufficient to determine that the technology results in a meaningful improvement in net health outcomes.

For individuals with other tumors (e.g., breast cancer, brain cancer, prostate cancer, desmoid, nonspinal osteoid osteoma) who receive MRgFUS, the evidence includes nonrandomized, uncontrolled phase II trials and several case series. Relevant outcomes are symptoms, health status measures, and treatment-related morbidity. A nonrandomized, uncontrolled phase II trial evaluating MRgFUS for prostate cancer reported a 93% success rate at 5 months and an 86% success rate at 2-years. Another nonrandomized, phase II trial in patients with prostate cancer reported that at 24 months, 88% (78 out of 89) of patients had no evidence of grade group 2 or higher prostate cancer in the treated area. Use of MRgFUS for the treatment of nonspinal osteoid osteoma consists of several larger case series, including a propensity score-matched retrospective study that reported similar reductions in pain with radiofrequency ablation and MRgFUS. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with medicine-refractory essential tremors who receive magnetic resonance-guided focused ultrasound (MRgFUS), the evidence includes technology assessment, meta-analyses, and a double-blind, sham-controlled randomized trial. Relevant outcomes are symptoms, health status measures, and treatment-related morbidity. The assessment did not pool study results but concluded that, overall, MRgFUS decreased tremor severity and improved quality of life (QOL). One meta-analysis reported significant improvements in hand tremor scores from baseline up to 24 months post-treatment, with evidence of a diminishing treatment benefit over time. A second meta-analysis showed significant

improvements in hand tremor scores and quality of life through 36 months post treatment. A third meta-analysis found similar improvements in tremor severity with MRgFUS to unilateral DBS, but improvements in both were inferior to bilateral DBS. The sham-controlled randomized trial found significant improvements in the treatment group in tremor severity, functional improvement, and quality of life after 3 months of follow-up. The improvements in hand tremor score, function, and quality of life were maintained at the 2-year follow-up. The evidence is sufficient to determine that the technology results in a meaningful improvement in net health outcomes.

For individuals with medicine-refractory tremor dominant Parkinson's disease (PD) who receive magnetic resonance-guided focused ultrasound (MRgFUS), the evidence includes a systematic review and pilot randomized controlled trial (RCT). Relevant outcomes include symptoms, functional outcomes, quality of life, and treatment-related morbidity. The double-blind, sham-controlled, pilot randomized trial (N=27) found significant improvements in the treatment group in tremor severity after 3 months of follow-up. The systematic review, which included the RCT plus additional small observational studies, found significant improvements in tremor severity through 12 months and quality of life through 6 months post procedure. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

### Additional Information

Not applicable.

## OBJECTIVE

The objective of this evidence review is to evaluate whether magnetic resonance-guided focused ultrasound (MRgFUS) improves the net health outcome in individuals with uterine fibroids, metastatic bone cancer, other tumors, medication-refractory essential tremors, or medication-refractory tremor dominant Parkinson's disease.

## PRIOR APPROVAL

Not applicable.

## POLICY

Magnetic resonance-guided high intensity focused ultrasound (MRgFUS) ablation may be considered **medically necessary** for pain palliation in adult patients with metastatic bone cancer who have failed or are not candidates for radiotherapy.

Magnetic resonance-guided high intensity focused ultrasound (MRgFUS) may be considered **medically necessary** for the treatment of medicine-refractory essential tremors (ET).

Magnetic resonance-guided high intensity focused ultrasound (MRgFUS) ablation is considered **investigational** when the above criteria is not met and for all other indications, including but not limited to the following, because the evidence is insufficient to determine the effects of the technology on net health outcomes:

- Brain cancer
- Breast cancer
- Desmoid tumors

- Medication-refractory tremor dominant Parkinson disease
- Non-spinal osteoid osteoma
- Prostate cancer
- Uterine fibroids

**ALL** Category III will be considered **investigational** unless the code is explicitly addressed as a covered service in a Wellmark BlueCross BlueShield Medical Coverage Policy.

## POLICY GUIDELINES

*Category III codes are a set of temporary (T) codes for emerging technologies, services, and procedures that allow for data collection by the American Medical Association's (AMA). If a Category III code is available, providers must use that code instead of an unlisted or deleted Category I code. The services or procedures represented by Category III codes may not have FDA approval, may not be performed by many health care professionals across the country, and the service or procedure may not have proven clinical efficacy. Certain T codes may be addressed as a covered service in a Wellmark BlueCross BlueShield Medical Coverage Policy. But, unless there is explicit Policy criteria that specifically extends coverage to a particular Category III code, the code would generally be considered experimental, investigational, or unproven.*

### Coding

See the [Codes table](#) for details.

## BACKGROUND

### Uterine Fibroids

Uterine fibroids are one of the most common conditions affecting women in the reproductive years. African American women have a greater lifetime incidence of uterine fibroids compared to other racial groups. Symptoms of uterine fibroids include menorrhagia, pelvic pressure, or pain.

Treatment approaches currently available to treat symptomatic uterine fibroids include hysterectomy, abdominal myomectomy, laparoscopic and hysteroscopic myomectomy, hormone therapy, uterine artery embolization, and watchful waiting. Hysterectomy and various myomectomy procedures are considered the criterion standard treatments.

### Metastatic Bone Disease

Metastatic bone disease is one of the most common causes of cancer pain.

Existing treatments include conservative measures (e.g., massage, exercise) and pharmacologic agents (e.g., analgesics, bisphosphonates, corticosteroids). For patients who do not respond to these treatments, standard care is external-beam radiotherapy. However, a substantial proportion of patients have residual pain after radiotherapy, and there is a need for alternative treatments for these patients.

### Other Tumors

Other tumors (breast, prostate, brain and desmoid tumors as well as non-spinal osteoid osteoma).  
Current treatment: standard of care.

## **Essential Tremors**

Essential tremor (ET) is the most common movement disorder, with an estimated prevalence of 5% worldwide. Essential tremor most often affects the hands and arms, may affect head and voice, and rarely includes the face, legs, and trunk. Essential tremor is heterogeneous among patients, varying in frequency, amplitude, causes of exacerbation, and association with other neurologic deficits.

The neuropathology of ET is uncertain, with some evidence suggesting that ET is localized in the brainstem and cerebellum. If patients with ET experience intermittent or persistent disability due to the tremors, initial therapy is with drugs (beta-blockers or anticonvulsants). For medicine-refractory patients, surgery (deep brain stimulation or thalamotomy) may be offered, though high rates of adverse events have been observed.

## **Tremor Dominant Parkinson's Disease**

The 3 cardinal features of Parkinson disease (PD) are tremor, bradykinesia, and rigidity. The tremor in PD is a resting tremor that occurs when the body part is not engaged in purposeful activities. Major subtypes of PD include tremor-dominant, akinetic-rigid, and postural instability and gait difficulty. The progression of PD is highly variable, and patients can change subtypes as the disease progresses.

Dopaminergic therapy (i.e., levodopa or a dopamine agonist) is the first-line treatment for PD, which improves tremor. Amantadine and anticholinergics (e.g., trihexyphenidyl) can also be considered as initial treatment for tremor-dominant PD or as add-on therapy in patients who have persistent tremor despite dopaminergic therapy. For medication-refractory patients, surgery (deep brain stimulation or lesioning procedures) may be offered. Lesioning procedures include conventional unilateral thalamotomy and focused ultrasound thalamotomy. Deep brain stimulation is the most frequently performed surgical procedure for the treatment of PD.

## **Magnetic Resonance-Guided Focused Ultrasound (MRgFUS)**

Magnetic resonance-guided focused ultrasound (MRgFUS) is a non-invasive treatment that combines 2 technologies: focused ultrasound and magnetic resonance imaging (MRI). The ultrasound beam penetrates through the soft tissues and, using MRI for guidance and monitoring, the beam can be focused on targeted sites. The ultrasound causes a local increase in temperature in the target tissue, resulting in coagulation necrosis while sparing the surrounding normal structures. The ultrasound waves from each sonication are directed at a focal point which has a maximum focal volume of 20 mm in diameter and 15 mm in height/length. This causes a rapid rise in temperature (i.e., to approximately 65°C to 85°C), which is sufficient to ablate tissue at the focal point. In addition to providing guidance, the associated MRI can provide on-line thermometric imaging that provides a temperature “map” to confirm the therapeutic effect of the ablation treatment and allow for real-time adjustment of the treatment parameters.

The U.S. Food and Drug Administration (FDA) approved the ExAblate® MRgFUS system (InSightec) for 4 indications: treatment of uterine fibroids (leiomyomata), palliation of pain associated with tumors metastatic to bone, medication refractory ET, and tremor-dominant PD. The ultrasound equipment is specifically designed to be compatible with magnetic resonance magnets, and it is integrated into standard clinical MRI units; it also includes a patient table, which has a cradle that houses the focused

ultrasound transducer in water or a light oil bath. Some models have a detachable cradle; only certain cradle types can be used for palliation of pain associated with metastatic bone cancer. For treating pain associated with bone metastases, the aim of MRgFUS is to destroy nerves in the bone surface surrounding the tumor.

MRgFUS is also being investigated for the treatment of other tumors, including breast, prostate, brain, and desmoid tumors as well as nonspinal osteoid osteoma.

## Regulatory Status

In October 2004, the U.S. Food and Drug Administration (FDA) approved via the premarket application (PMA) process, the ExAblate® 2000 System (Insightec, Inc., Haifa, Israel) for “ablation of uterine fibroid tissue in pre- or perimenopausal women with symptomatic uterine fibroids who desire a uterine sparing procedure.” Treatment is indicated for women with a uterine gestational size of less than 24 weeks who have completed childbearing.

In October 2012, the U.S. Food and Drug Administration (FDA) approved the ExAblate® System, Model 2000/2100/2100 VI via the PMA process. The intended use of the device is for pain palliation in adult patients with metastatic bone cancer who failed or are not candidates for radiation therapy. The device was evaluated through an expedited review process. The FDA required a post-approval study with 70 patients to evaluate the effectiveness of the system under actual clinical conditions.

In July 2016, the U.S. Food and Drug Administration (FDA) approved the ExAblate Neuro system for use in the unilaterally thalamotomy treatment of idiopathic essential tremor patients with medication refractory tremor. Patients must be at least age 22. The designated area in the brain responsible for the movement disorder symptoms (ventralis intermedius) must be identified and accessible for targeted thermal ablation by the ExAblate device. ExAblate Neuro uses magnetic resonance images taken during the procedure to deliver focused ultrasound to destroy brain tissue in a tiny area thought to be responsible for causing tremors.

In November 2021, the FDA approved the use of the Exablate Prostate System for prostate tissue ablation through the premarket approval process.

FDA product codes: NRZ, POH, PLP.

ExAblate 4000 Type 2 MRI-guided focused ultrasound is currently being studied in the clinical setting to temporarily disrupt the blood brain barrier (BBB) with focused ultrasound allowing for the collection of circulating tumor DNA (ctDNA) from brain tumors such as glioblastoma to determine therapy and in neuro-oncology for the treatment of brain tumors in pediatric and adult individuals for enhancing cancer therapy delivery to the brain. This device currently does not have FDA approval for these indications. For current ongoing and unpublished trials that might influence this review refer to [clinicaltrials.gov](https://clinicaltrials.gov).

## RATIONALE

This evidence review was created in August 2007 and has been updated regularly with searches of the PubMed database. The most recent literature update was performed through September 4, 2025.

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to

function-including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

## **Magnetic Resonance-Guided Focused Ultrasound for Uterine Fibroids**

### ***Clinical Context and Therapy Purpose***

The purpose of magnetic resonance magnetic resonance-guided focused ultrasound (MRgFUS) in individuals with uterine fibroids is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

### ***Populations***

The relevant population of interest is individuals with uterine fibroids.

### ***Interventions***

The therapy being considered is MRgFUS, which is a thermoablative procedure to heat targeted tissue in small volume increments, under constant magnetic resonance imaging guidance.

### ***Comparators***

The comparators of interest are alternative nonsurgical treatments or surgery.

### ***Outcomes***

For uterine fibroids, the goal is to reduce or eliminate fibroid-related symptoms by reducing fibroid size. Measures to assess the effect of treatment include quality of life, change in uterine and fibroid volume, pain levels, and pain medication use. Outcome measures can be assessed at several months to several years post procedure.

## **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies;
- To assess long-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought;
- Studies with duplicative or overlapping populations were excluded.

## Review of Evidence

Evidence for the use of MRgFUS for the treatment of uterine fibroids consists of randomized controlled trials (RCTs), systematic reviews, and many observational studies. RCTs and relevant non-randomized trials not included in the systematic reviews are summarized.

## Systematic Reviews

A systematic review, published by Gizzo et al (2014), conducted a literature search through February 2013 and identified 38 uncontrolled studies with a total of 2500 patients who underwent MRgFUS for the treatment of uterine fibroids. All published studies included women 18 years or older with symptomatic uterine fibroids, and most excluded patients who desired future pregnancies. Reviewers did not pool study findings due to the heterogeneity of outcomes but concluded that, overall, MRgFUS appeared to be a safe, noninvasive option for treating uterine fibroids. Future research, particularly RCTs, were recommended to compare MRgFUS with other noninvasive procedures and to explore the fertility-sparing potential further.

A meta-analysis by Xu et al (2021) compared the reintervention rates of UAE, myomectomy, and MRgFUS in patients with uterine fibroids. There were 31 studies (N=42,103) that were included in the analysis, with 6 being RCTs and the other 25 being cohort studies. The 12-month, 24-month, 36-month, and 60-month re-intervention rates were assessed as the primary outcome. Myomectomy has the lowest re-intervention rate of the 3 regimens in all time points assessed, while the MRgFUS had the highest re-intervention rate. The estimations of the pooled rates of reintervention of MRgFUS also increased rapidly in the sixtieth month after treatment compared to myomectomy and UAE.

## Randomized Controlled Trials

Barnard et al (2017) published preliminary results from Fibroid Interventions: Reducing Symptoms Today and Tomorrow trial, a parallel RCT and cohort study comparing MRgFUS with fibroid embolization to treat uterine fibroids. For the RCT, patients were randomized to uterine artery embolization (UAE; n=22) or to MRgFUS (n=27). Patients and investigators were not blinded. Women who did not want to be randomized were enrolled in the cohort study; 16 underwent UAE and 16 underwent MRgFUS. Patients were instructed to keep diaries with the following information: medication use, return to normal activities, and symptoms. After 6 weeks of follow-up for the RCT patients, there were no differences between groups in symptoms such as fatigue, hot flashes, discomfort urinating, vaginal discharge, or constipation. Recovery was significantly faster in the MRgFUS group, as measured by the first day back to work and the first day back to normal. Medication use (i.e., opioids, nonsteroidal anti-inflammatory drugs, acetaminophen or aspirin, nausea medication, bowel medication) was also significantly lower in the MRgFUS group. Analyses combining the RCT, and cohort patients showed similar results. The MRgFUS procedure took significantly longer than the UAE procedure. A trial limitation was the inability to recruit more patients. Long-term follow-up results were reported by Laughlin-Tommaso et al (2019): Patients in both the RCT

and cohort studies had follow-up for up to 3 years. The primary outcome assessed was reintervention for uterine fibroids within 3 years; secondary outcomes included change in anti-Mullerian hormone levels and standardized measures of quality of life, pain, sexual function, and fibroid symptoms. Among the women in the MRgFUS arm (n=43), 13 (30%) had a second fibroid procedure compared to 5 (13%) women in the UAE arm (hazard ratio [HR], 2.81; 95% confidence interval [CI], 1.01 to 7.79). There was a significantly greater absolute decrease in anti-Mullerian hormone levels at 24 months in the UAE arm compared to the MRgFUS arm.

A pilot sham-controlled randomized trial evaluating MRgFUS for the treatment of uterine fibroids was published by Jacoby et al (2016). The trial included 20 premenopausal women with symptomatic uterine fibroids. Patients were randomized to MRgFUS with the ExAblate 2000 System (n=13) or to a sham treatment not using thermal energy (n=7). The sample size was calculated to assess the feasibility of a larger trial, not to provide sufficient statistical power. All patients who were assigned to the MRgFUS group and 6 of 7 in the placebo group received their allocated treatment; patients were unblinded at 3 months. The trialists concluded that a larger sham-controlled randomized trial of MRgFUS was feasible.

### **Non-randomized Trials**

In 2016, Chen et. al. evaluated 107 women undergoing MRgFUS for the treatment of uterine fibroids. Efficacy was defined as the proportion of patients with at least 10% fibroid shrinkage from baseline, as measured by magnetic resonance imaging. At the 6-month follow-up, 93% efficacy was reported.

### **Fertility Following Magnetic Resonance-Guided Focused Ultrasound for Treatment of Uterine Fibroids**

A prospective registry of pregnancies after MRgFUS had been maintained by the manufacturer of the ExAblate device. Rabinovici et al (2010) reported on 54 known pregnancies a mean of 8 months after treatment. They included 8 pregnancies from clinical trials designed for women who did not desire pregnancy, 26 pregnancies after commercial treatment, and 20 pregnancies in 17 patients from an ongoing study of MRgFUS in women trying to conceive. Twenty-two (42%) of the 54 pregnancies resulted in deliveries and 11 were ongoing beyond 20 weeks at the time the article was written. There were 14 (26%) miscarriages and 7 (13%) elective terminations. Among the 22 live births, the mean live birth weight was 3.3 kg, and the vaginal delivery rate was 64%. The article provided initial information on the impact of MRgFUS on uterine fibroids in pregnancy; findings suggested that fertility may be maintained but that the number of cases was too small to draw definitive conclusions. The study also did not address the possible impact of MRgFUS treatment on the future ability to become pregnant.

A prospective cohort study by Otonkoski et al (2023) evaluated if there was any adverse impact of MRgFUS treatment on ovarian reserve. Seventy-four premenopausal women were included who had either symptomatic uterine fibroids or adenomyosis. Ovarian reserve was estimated using serum Anti-Mullerian hormone (AMH) levels before and 3 months after treatment. The median baseline AMH level prior to treatment was 1.20 (range, 0.1 to 7.75 mcg/L) and 1.23 (range, 0.1 to 8.51 mcg/L) after treatment, and no statistically significant change was detected (p=.90). Also, none of the patients reported any symptoms that would indicate a loss of ovarian function.

### **Section Summary: Uterine Fibroids**

For the treatment of uterine fibroids, there are 2 RCTs, 1 with 49 women that compared MRgFUS with UAE and the other a feasibility trial assessing 20 women that had a sham control. Several

nonrandomized studies have also compared MRgFUS with different treatments. The sham-controlled randomized trial concluded that a larger trial would be feasible. The trial reported significantly lower fibroid volumes in the active treatment group; however, there were no statistically significant differences in quality of life between the groups. The other RCT reported no significant differences in medication use or symptoms between the MRgFUS and UAE groups. Recovery was significantly faster in the MRgFUS group than in the UAE group, however long-term follow-up results reported a lower reintervention rate and greater improvement in symptoms after UAE compared to MRgFUS. A 2014 systematic review, which identified only noncomparative studies, did not pool results due to heterogeneity in outcomes among the studies. While reviewers concluded that MRgFUS may be a safe and effective minimally invasive option for the treatment of fibroids, they noted that RCTs comparing MRgFUS with other noninvasive procedures would be informative. A 2021 meta-analysis reported that, comparatively, myomectomy had the lowest re-intervention rate of the 3 regimens (myomectomy vs UAE vs MRgFUS) in all time points assessed, while the MRgFUS had the highest re-intervention rate.

## **Magnetic Resonance-Guided Focused Ultrasound for Palliative Treatment of Bone Metastases**

### *Clinical Context and Therapy Purpose*

The purpose of magnetic resonance-guided focused ultrasound (MRgFUS) in individuals with metastatic bone cancer is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

### *Populations*

The relevant population of interest is individuals with metastatic bone cancer who have failed radiotherapy or who are not candidates for radiotherapy.

### *Interventions*

The therapy being considered is MRgFUS, which is a thermoablative procedure to heat targeted tissue in small volume increments, under constant magnetic resonance imaging guidance.

### *Comparators*

The comparator of interest for metastatic bone cancer is supportive care.

### *Outcomes*

For metastatic bone cancer, the goal is to alleviate pain. Measures to assess the effect of treatment include pain levels and pain medication use. Outcome measures can be assessed at several months to several years post-procedure.

## **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies;
- To assess long-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought;
- Studies with duplicative or overlapping populations were excluded.

## Review of Evidence

Evidence for the use of magnetic resonance-guided focused ultrasound (MRgFUS) for the treatment of painful bone metastases consists of a large RCT and a systematic review of RCTs and observational studies. Observational studies with longer-term follow-up or not included in the systematic review are summarized.

## Systematic Reviews

Baal et al (2021) conducted a systematic review of studies published between 2007 and 2019 evaluating MRgFUS treatment for painful bone metastases. A total of 33 studies were identified, comprising 3 RCTs, 6 retrospective studies, and 24 prospective studies, representing 1082 patients. Thirteen studies were available in abstract form only. The median study sample size was 21 patients (range 5 to 140) with a median follow-up period of 3 months (range, 1 to 12 months). Efficacy was assessed by treatment response (complete response or partial response [ $\geq 2$ -point improvement in pain score]) and the mean difference in pain scores (10-point VAS [visual analog scale] or NRS [numeric rating scale]) from baseline to month 1/month 3. The pooled proportion of patients with a treatment response to MRgFUS was 79% (95% CI, 73% to 83%; based on 20 studies [n=636]). The pooled 1-month and 3-month mean difference from baseline in pain score were -3.8 (95% CI, -4.3 to -3.3) and -4.4 (95% CI, -5.0 to -3.7), respectively (based on 20 studies [n=543]). Across 26 studies (n=799), 7 high-grade adverse events were observed (1 deep vein thrombosis, 2 cases of grade 3 skin burn, and 4 fractures). Approximately 11.8% of patients experienced sonication-related pain during MRgFUS treatment. The analysis was limited by a lack of a pooled comparator. Additionally, there was substantial heterogeneity of the included studies due to variable study populations (e.g., type of primary cancer), reported data, and treatment details. Many of the included studies had follow-up periods that were limited to 3 months.

## Randomized Controlled Trials

In an RCT evaluating the ExAblative system for the treatment of painful bone metastases, Hurwitz et al. (2014) evaluated patients with 3 or more months of life expectancy who had painful bone metastases despite radiotherapy, or who were unsuitable for or declined radiotherapy. Patients rated tumor pain on a 10-point scale NRS at 4 or greater. While they could have up to 5 painful lesions, only 1 lesion was treated, and it had to cause pain at least 2 points greater on the NRS than any other lesion. Also targeted tumors needed to be device accessible. Study participants were randomized 3:1 to active (n=122) or sham (n=39) MRgFUS treatment. Ten patients in the treatment group and 4 in the sham group did not receive the allocated treatment. An additional 26 patients in the treatment group and 23 in the sham group did not complete the 3-month follow-up. A larger proportion of the placebo group dropped out: 17 (49%) of 35 who were treated decided to have rescue MRgFUS treatment after a lack of response to placebo. A modified intention-to-treat analysis was used that included patients who had at least 1 MRgFUS or placebo sonication. Missing values were imputed using the last observation carried forward method. The primary efficacy endpoint, assessed at 3 months, was a composite outcome comprised of the change in baseline in worst NRS score and morphine equivalent daily dose (MEDD) intake. Patients

were considered responders if their worst NRS score decreased by at least 2 points and if their MEDD intake did not increase more than 25% from baseline to 3 months. NRS scores and MEDD intake were reported separately as secondary outcomes.

Seventy-two (64%) of 112 patients in the MRgFUS group and 7 (20%) of 35 patients in the control group were considered responders, as previously defined. The difference was statistically significant ( $p=.01$ ), favoring active treatment. When the 2 measures comprising the primary endpoint were analyzed separately, there was a statistically significant difference between groups in change in worst NRS score and a nonsignificant difference in change from baseline in pain medication. The NRS score decreased by a mean (standard deviation [SD]) of 3.6 (3.1) points in the MRgFUS group and by a mean of 0.7 (2.4) in the placebo group ( $p<.01$ ). Change in MEDD from baseline was 3.7 in the MRgFUS group and 15.3 in the placebo group. Fifty-one (46%) patients in the MRgFUS group and 1 (3%) in the placebo group experienced at least 1 adverse event. Most adverse events were transient, with the most common being sonication pain, experienced by 36 (32%) patients in the MRgFUS group. In 17 (15%) patients, sonication pain was severe; 3 patients did not complete treatment due to pain. The most clinically significant adverse events that lasted more than a week were third-degree skin burns in 1 patient (associated with noncompliance with the treatment protocol) and fracture in 2 patients (1 of which was outside the treatment location). Potential trial limitations included a nonconventional primary outcome measure and the small initial size of the sham group. Moreover, a large number of sham patients (66%) did not complete the 3-month follow-up; the trialists indicated that this low completion rate was due to a lack of response to placebo treatment.

## **Observational Studies**

Arrigoni et al (2017) evaluated the use of MRgFUS in a case series of 14 patients with intra-articular benign bone lesions who were followed for 12 months. Pain was measured by a VAS and all patients underwent computed tomography and magnetic resonance imaging. Mean pain scores significantly decreased from 7.8 pretreatments to 2.0 at 6-month follow-up to 0.6 at 12-month follow-up ( $p<.001$ ). No patients reported worse symptoms and none reported the procedure unsuccessful. Diagnostic imaging supported the clinical findings and showed calcification of the lesion, lack of contrast enhancement, and resolution of bone edema.

## **Section Summary: Palliative Treatment of Bone Metastases**

The evidence consists of a systematic review of RCTs and observational studies, a single industry sponsored RCT, and case series. The RCT found significant improvement after MRgFUS in a composite outcome comprised of a reduction in pain and morphine use, and in pain reduction as a stand-alone outcome. This trial was appropriately sham controlled. A substantial proportion of patients in the treatment group experienced adverse events but most adverse events were transient and not severe. Pooled efficacy data from a systematic review reported a treatment response to MRgFUS of 79%.

## **Magnetic Resonance-Guided Focused Ultrasound for Other Tumors**

### ***Clinical Context and Therapy Purpose***

The purpose of magnetic resonance-guided focused ultrasound (MRgFUS) in individuals with other tumors is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

## **Populations**

The relevant population of interest are individuals with other tumors (e.g., breast cancer, brain cancer, prostate cancer, desmoid, non-spinal osteoid osteoma).

## **Interventions**

The therapy being considered is MRgFUS, which is a thermoablative procedure to heat targeted tissue in small volume increments, under constant magnetic resonance imaging guidance.

## **Comparators**

The comparator of interest for other tumors is standard of care.

## **Outcomes**

For other tumors, the goal is tumor ablation. Outcomes include reductions in tumor size. Outcome measures can be assessed at several months to several years post-procedure.

## **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies;
- To assess long-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought;
- Studies with duplicative or overlapping populations were excluded.

## **Review of Evidence**

### **Non-randomized Trials**

Ghai et al (2021) conducted a phase II trial to evaluate the safety and efficacy of transrectal MRgFUS treatment for intermediate-risk prostate cancer. The primary efficacy endpoint was the presence of residual disease at the treatment site at 5 months after the procedure. Ninety-three percent of patients were free of clinically significant prostate cancer at the 5-month biopsy. No major treatment-related adverse events occurred. Ghai et al (2024) recently published the 24-month follow-up results. Treatment was successfully completed for 43 patients through month 24, although 1 participant refused biopsy. After 2-years, 39/42 participants (93%) had no clinically significant prostate cancer at the treatment site and 36/42 (86%) had no cancer in the entire prostate gland. Additionally, there was no significant decline in quality of life per the validated questionnaires at 24-months and no major adverse events were recorded.

Ehdaie et al (2022) conducted a phase II trial to evaluate whether MRgFUS could safely reduce treatment burden for patients with intermediate-risk prostate cancer. The co-primary efficacy endpoints were oncological efficacy (defined as absence of cancer that was grade group 2 or higher in the treated area on prostate biopsy) and safety (measured by adverse event reporting). At 24 months, 88% (78 out of 89) of patients had no evidence of grade group 2 or higher prostate cancer in the treated area; there was 1 grade 3 adverse event that was reported and no grade 4 or 5 adverse events.

Study characteristics and results are presented in Tables 1 and 2.

**Table 1. Summary of Key Nonrandomized Trials Characteristics**

Study	Study Type	Country	Dates	Participants	Treatment	Comparator	Follow-Up
Ghai et al (2021); Ghai et al (2024)	Prospective phase II trial	Canada	2016-2019	44 men with unifocal, intermediate-risk prostate cancer with <20 mm of MRI-visible GG2 or GG3 disease (not previously treated); 2 participants did not undergo biopsy at 2-year follow-up	Transrectal MRgFUS	None	24 months
Ehdaie et al (2022)	Prospective phase II trial	USA	2017-2022	101 men ≥50 years old, with intermediate risk prostate adenocarcinoma with no previous treatment for prostate cancer	Transrectal MRgFUS	None	24 months

GG: grade group; MRgFUS: magnetic resonance-guided focused ultrasound; MRI: magnetic resonance imaging.

**Table 2. Summary of Key Nonrandomized Trials Results**

Study	Residual Disease	Recurrence/response	PSA	Adverse Events
Ghai et al (2021); Ghai et al (2024)	7% (95% CI, 2.4 to 18.2) had residual disease at 5 months after ablation	Of the 42 other participants at 2 years, 36 (86%) were free of clinically significant disease, 3 (7%) had clinically significant disease at the treatment site, and 3 (7%) had clinically significant disease outside of the treatment zone.	Median PSA was 2.4 ng/mL (IQR, 1.1 to 5.4) at 5 months (baseline PSA was 6.4 ng/mL [IQR, 1.1 to 5.4]) and 2.7 ng/mL (IQR, 1.2 to 6.7) at 24-months.	5 months: 16 patients reported dysuria; 5 patients required antispasmodics for bladder spasm in the first week; 2 patients had urinary retention; 1 patient had severe pelvic pain 24-months: no major adverse events reported.
Ehdaie et al (2022)	NR	96 out of 101 patients (95%; 95% CI, 89 to 98) had no evidence of grade group 2 or higher prostate cancer on 6-month biopsy in the treated area; 78 out of 89 patients (88%; 95% CI, 79 to 94) had no evidence of grade group 2 or higher cancer on 24-month biopsy in the treated area	Mean decrease in PSA after treatment was -3.0 ng/mL (95% CI, -3.6 to -2.4) at 6 months and -2.6 ng/mL (95% CI, -3.3 to -2.0) at 24 months.	No serious TRAEs were reported during the study period. There was 1 grade 3 AE (UTI) that was reported. Common AEs that were reported (grade 2 or lower) included hematuria and urinary retention.

AE, adverse event; CI: confidence interval; IQR: interquartile range; MRgFUS: magnetic resonance-guided focused ultrasound; NR, not reported; PSA: prostate specific antigen; TRAE, treatment-related adverse event; UTI, urinary tract infection

## Observational Studies

Only small case series have assessed the safety and/or efficacy of MRgFUS for treating tumors related to breast cancer, and brain cancer. The most recent case series on the use of MRgFUS for breast cancer ablation was published by Merckel et al (2016). Ten patients with early-stage invasive breast cancer underwent MRgFUS prior to surgical resection. Ablation was confirmed histopathologically in 6 of these patients. The investigators concluded that MRgFUS is safe and feasible. A noted limitation is the long procedure time (average, 145 minutes), due to waiting time after contrast injection and time to find a proper magnetic resonance navigator signal.

Several case series have investigated the use of MRgFUS for non-spinal osteoid osteoma. Arrigoni et al (2021) conducted a propensity score-matched retrospective study to compare treatment with radiofrequency ablation and MRgFUS. A total of 116 patients were treated (61 with radiofrequency ablation and 55 with MRgFUS). After propensity score matching, both radiofrequency ablation and MRgFUS treatment resulted in a significant reduction in pain from baseline as measured by VAS (8.9 to 0.02 and 8.8 to 0.54, respectively). There was no statistically significant difference between the mean values of both groups after the treatment. Four cases of relapse (1 with radiofrequency ablation and 3 with MRgFUS) were observed. Arrigoni et al (2019) prospectively enrolled children into a study to evaluate MRgFUS treatment for osteoid osteoma. The primary clinical endpoint was defined as the absence of pain (evaluated on the Faces Pain Scale-Revised) at the first follow-up study 1 week after the procedure. A total of 33 children were included in the study and treated with MRgFUS. The mean pain score at baseline was 7.6; the score at week 1 after the procedure significantly improved in all children (mean score, 0.21). Complete absence of pain was reported in 32 of 33 (97%; 95% CI, 84 to 100) patients at week 1. At the 24-month follow-up visit, imaging results confirmed the complete disappearance of bone edema around all lesions. Geiger et al (2014) prospectively enrolled patients into a study to evaluate MRgFUS treatment for osteoid osteoma. Clinical success was evaluated based on pain reduction (evaluated on a VAS) through 12 months. At the 12-month follow-up, complete clinical success was achieved in 90% of the 29 patients enrolled (mean VAS, 0±0 points); partial success was achieved in the remaining patients (mean VAS, 5±0 points).

In addition, several case series have investigated the use of MRgFUS for desmoid tumors. Avedian et al (2016) used MRgFUS to treat 9 patients with desmoid tumors. Five patients were available for follow-up for at least 12 months. Mean decrease in tumor size was 36% (95% CI, 7% to 66%). Bucknor et al (2017) described the use of MRgFUS to treat 3 patients with large aggressive desmoid tumors within the posterior thigh. Each patient received multiple MRgFUS treatments. In this case series, the use of MRgFUS for desmoid tumors required different treatment parameters than those used for fibroids or bone lesions, due to differences in vascularity of the target tissue and the need for effective skin protection when using MRgFUS on extremities. Ghanouni et al (2017) used MRgFUS to treat 15 patients with extra-abdominal desmoid tumors. Treatment times ranged from 0.8 to 8 hours. Results were presented on 9 patients (3 were lost to follow-up before 6 months, 3 received additional treatments). Seven of 9 patients experienced durable clinical benefits, with a median reduction in tumor volume of 98%. Treatment-related adverse events included skin burns, nerve injury, and off-target heating.

## **Section Summary: Treatment of Other Tumors**

Evidence on the use of MRgFUS for the treatment of prostate cancer consists of a nonrandomized, uncontrolled phase II trial, which reported a 93% success rate at 5 months. Evidence on the use of MRgFUS for the treatment of non-spinal osteoid osteoma consists of several case series, including a propensity score-matched retrospective study that reported similar reductions in pain with radiofrequency ablation and MRgFUS. Currently, evidence on the use of MRgFUS for the treatment of other tumors consists of small case series, which is insufficiently robust to draw conclusions about efficacy. RCTs comparing MRgFUS with other noninvasive procedures would be informative.

## **Magnetic Resonance-Guided Focused Ultrasound for Essential Tremors**

### ***Clinical Context and Therapy Purpose***

The purpose of magnetic resonance-guided focused ultrasound (MRgFUS) in individuals with essential tremors (ET) is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

### **Populations**

The relevant population of interest is individuals with medication-refractory ET.

### **Interventions**

The therapy being considered is MRgFUS, which is a thermoablative procedure to heat targeted tissue in small volume increments, under constant magnetic resonance imaging guidance.

### **Comparators**

The comparators of interest for ET are neurosurgery or standard of care. Surgical procedures include deep brain stimulation (DBS) of the ventral intermediate nucleus of the thalamus and stereotactic thalamotomy.

### **Outcomes**

For ET, the goal is to decrease the frequency of tremors and improve the quality of life. Outcome measures can be assessed at several months to several years post-procedure.

### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies;
- To assess long-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought;
- Studies with duplicative or overlapping populations were excluded

### **Review of Evidence**

Evidence for the use of MRgFUS to treat medication-refractory ET consists of a technology assessment, meta-analyses, and a double-blind, sham-controlled randomized trial.

### **Systematic Reviews**

Mortezaei et al (2024) published a systematic review and meta-analysis evaluating the long-term safety and efficacy of MRgFUS for treating medication-refractory ET. A total of 43 observational clinical studies including 1818 patients were identified. Follow-up durations across studies ranged from 3 months to 8.4 years. Primary outcomes included changes in tremor severity as measured by the Clinical Rating Scale for Tremor (CRST) and hand tremor scores. The total CRST score showed significant reductions at 3, 6, and 12 months post-treatment (standard mean difference [SMD], -4.5 [p=.0069]; SMD, -4.9 [p=.0045]; SMD, -2.95 [p=.0039], respectively). Similarly, hand tremor scores significantly improved at 3, 6, 12, 24, and 36 months (SMD, -3.99 [p=.05]; SMD, -4.5 [p=.05]; SMD, -1.99 [p<.0001]; SMD, -2.07 [p=.0002]; SMD, -2.1 [p<.0001], respectively). Quality of life, assessed by the Quality of Life in Essential Tremor

Questionnaire (QUEST), also improved at multiple time points through 3 years post-treatment (SMD, -2.8 to -1.14 [p-values ranging from .0025 to .08]).

Miller et al. (2021) published a meta-analysis that evaluated the efficacy of magnetic resonance-guided focused ultrasound (MRgFUS) for treating medication-refractory ET with a focus on long-term trends and the durability of the response. Twenty-one studies (N=395) were included; 17 were prospective studies, 3 were retrospective, and only 1 was an RCT. Hand tremor scores decreased from a weighted mean pre-operative value of  $19.2 \pm 5.0$  to  $7.4 \pm 5.0$  after 3 months. Over time, the hand tremor score values gradually increased:  $8.3 \pm 5.3$  after 12 months and  $9.1 \pm 5.4$  after 36 months. The pooled standardized mean difference of hand tremor scores compared to pre-treatment values was 2.68 (95% CI, 1.94 to 3.41) at 3 months (5 studies), 2.44 (95% CI, 1.97 to 2.91) at the 12-month time point (7 studies), and 2.18 (95% CI, 1.50 to 2.86) at the 24-month time point (3 studies). Clinical Rating Scale for Tremor scores were only reported through 12 months. The pooled standardized mean difference in Clinical Rating Scale for Tremor scores compared to pre-treatment values was 1.86 (95% CI, 1.51 to 2.21) at the 3-month time point (8 studies) and 2.24 (95% CI, 1.55 to 2.94) at the 12-month time point (6 studies). Six studies reported Quality of Life in Essential Tremor Questionnaire (QUEST) scores as a quality-of-life measure. The pooled pre-treatment QUEST score was  $48.2 \pm 22.4$ , which improved to  $24.9 \pm 18.2$  at 3 months. Additionally, a single study detailed a mean  $23.8 \pm 19.6$  QUEST score at 36 months follow-up, an increase of 2.2 over 30 months.

Giordano et al (2020) conducted a meta-analysis to compare unilateral magnetic resonance-guided focused ultrasound (MRgFUS) to unilateral and bilateral DBS for medication-refractory ET. Forty-five studies published between 1996 and 2019 were identified. Thirty-seven studies (n=1202) evaluated DBS and 8 studies (n=477) evaluated MRgFUS. Fifteen studies had a retrospective study design, while 30 were prospectively designed. Means and standard deviations were calculated for each intervention and differences between groups were compared where appropriate. The average percentage improvement in tremor severity was significantly improved in the pooled DBS group ( $60.1\% \pm 9.7\%$ ) as compared to the MRgFUS group ( $55.6\% \pm 8.2\%$ ,  $p < .001$ ). Subgroup analyses demonstrated that the improvement in tremor severity was significantly greater with the bilateral DBS ( $61.2\% \pm 5.2\%$ ) as compared to both unilateral DBS ( $56.4\% \pm 9.7\%$ ) and MRgFUS; there was no significant difference between unilateral DBS and MRgFUS. For average percentage improvement in quality of life, MRgFUS was associated with significantly improved measures as compared to DBS ( $61.9\% \pm 7.9\%$  vs  $52.5\% \pm 16.2\%$ ,  $p < .001$ ). There were 517 complications reported in the DBS group and 484 complications reported in the MRgFUS group. The most common adverse events reported with DBS were lead-related complications (11.4%) and speech disturbances (11.1%). For MRgFUS, adverse events of sensory nature (36.7%) and gait disturbances/muscle problems (34.4%) were most common. Limitations of the review included the different scales used in studies to measure tremor severity and quality of life. There was only 1 retrospective study that directly compared DBS and MRgFUS.

The technology assessment was published by Health Quality Ontario (2018). The literature search, conducted through April 2017, identified 9 studies for inclusion: 4 single cohort studies, 2 retrospective chart reviews, 2 uncontrolled prospective studies, and an RCT. The RCT compared MRgFUS with sham treatment, and the chart reviews compared MRgFUS with DBS and radiofrequency thalamotomy. Study quality was evaluated using the GRADE system. The RCT was rated high-quality, the uncontrolled comparative studies were rated very low-quality, and the remaining studies were rated low-quality. All studies reported tremor severity as an outcome. Pooling of results was not conducted due to heterogeneity in study designs, analyses, and outcomes across the studies. Reviewers determined that, overall, MRgFUS decreased tremor severity and improved quality of life.

### **Randomized Controlled Trials**

A single high-quality study, a double-blind, sham-controlled randomized trial by Elias et al (2016) was identified by the 2 systematic reviews. Trial selection criteria included patients with moderate or severe postural or intention tremor of the hand ( $\geq 2$  on the Clinical Rating Scale for Tremor) and refractory to at least 2 medical therapies. Patients were randomized to MRgFUS thalamotomy (n=56) or sham treatment (n=20). Outcomes were tremor severity, improvement, and quality of life, measured at 3 months post-procedure. Patients in the treatment group were followed for an additional 12 months. The mean score for hand tremors improved significantly from baseline in the treatment group (47%) compared with the sham group (0.1%) at 3 months. Change in mean functional improvement score from baseline differed significantly in the MRgFUS group (62%) compared with the sham group (3%) at 3 months. Change in Quality of Life in Essential Tremor Questionnaire scores also differed significantly in the treatment group compared with the sham group, with the largest improvements experienced in the psychosocial domain. The improvements in hand tremor score, functional improvement, and quality of life were maintained at 12 months in the MRgFUS group.

Chang et. al. (2018) published results from 67 patients who participated in the open-label extension of the RCT. Because 9 patients from the original trial received additional treatment during the 2-year follow-up, they were excluded from the analysis. Improvements in tremor and disability scores were maintained at the 2-year follow-up (tremor,  $19.8 \pm 4.9$  [baseline] to  $8.8 \pm 5.0$  [at 2 years]; disability,  $16.4 \pm 4.5$  [baseline] to  $6.5 \pm 5.0$  [at 2 years]).

## **Section Summary: Essential Tremors**

Evidence for the use of MRgFUS in the treatment of medication-refractory ET consists of a technology assessment, meta-analyses, and a double-blind, sham-controlled randomized trial. The assessment did not pool results from the studies but concluded, overall, that MRgFUS decreased tremor severity and improved quality of life. One meta-analysis reported significant improvements in hand tremor scores from baseline up to 24 months post-treatment, with evidence of a diminishing treatment benefit over time. A second meta-analysis showed significant improvements in hand tremor scores and quality of life through 36 months post treatment. A third meta-analysis found similar improvements in tremor severity with MRgFUS to unilateral DBS, but improvements in both were inferior to bilateral DBS. The sham-controlled randomized trial which was considered high-quality found significant improvements in the treatment group in tremor severity, functional improvement, and quality of life after 3 months of follow-up, and these results were maintained through 2 years of follow-up.

## **Magnetic Resonance-Guided Focused Ultrasound for Tremor-Dominant Parkinson Disease**

### ***Clinical Context and Therapy Purpose***

The purpose of magnetic resonance-guided focused ultrasound (MRgFUS) in individuals with tremor-dominant Parkinson disease (PD) is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

### ***Populations***

The relevant population of interest is individuals with medication refractory tremor-dominant PD.

## **Interventions**

The therapy being considered is MRgFUS, which is a thermoablative procedure to heat targeted tissue in small volume increments, under constant magnetic resonance imaging guidance.

## **Comparators**

The comparators of interest for tremor-dominant PD are neurosurgery or standard of care. Surgical procedures include DBS and conventional unilateral thalamotomy.

## **Outcomes**

For refractory tremor associated with tremor-dominant PD, the goal is to decrease the frequency of tremors and improve quality of life. Outcome measures can be assessed at several months to several years post procedure.

## **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies;
- To assess long-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought;
- Studies with duplicative or overlapping populations were excluded.

## **Review of Evidence**

### **Systematic Reviews**

Evidence for the use of MRgFUS to treat medication-refractory tremor-dominant PD consists of a systematic review and a double-blind, sham-controlled randomized trial.

### **Systematic Review**

Monteiro et al (2024) published a systematic review with meta-analysis that evaluated the safety and efficacy of MRgFUS thalamotomy in tremor-dominant PD. Study characteristics and results are summarized in Table 3. A total of 13 studies (12 observational and 1 RCT [see Bond et al 2017 below]) involving 211 patients were analyzed. Tremor severity was assessed using the Clinical Scale Rating for Tremor (CSRT) and the Movement Disorders Society - Unified Parkinson's Disease Rating Scale (MDS-UPDRSIII), both showing significant improvements through 12 months post procedure. Quality of life also improved significantly, as demonstrated by a SMD of -0.86 (95% CI, -1.21 to -0.50) in Parkinson Disease Questionnaire (PDQ) scores from baseline to 6 months.

### **Table 3. SR & M-A Results**

Study	MDS-UPDRSIII	CRST	QOL measured by PDQ
Monteiro (2024)			
Total N	105	155	70
Pooled effect (95% CI)	1 month: MD, -8.92 (15.44 to -2.40)  6 months: MD, -7.39; (-11.47 to -3.30)  12 months: MD, -10.66 (-16.89 to -4.43)	1 month: SMD, -2.22 (-4.38 to -0.06)  3 months: -1.79 (-3.66 to -0.07)  12 months: -1.47 (-2.54 to -0.39)	6 months: MD, -0.86 (-1.21 to -0.50)

CI: confidence interval; CRST: Clinical Rating Scale for Tremor; M-A: meta-analysis; MD: mean difference; MDS-UPDRSIII: Movement Disorders Society-Unified Parkinson's Disease Rating Scale; PDQ: Parkinson Disease Questionnaire; QOL: quality of life; SMD: standardized mean difference; SR: systematic review.

### Randomized Controlled Trials

A double-blind, sham-controlled, pilot randomized trial by Bond et al (2017) assessed the safety and efficacy of unilateral MRgFUS thalamotomy in patients with tremor-dominant PD. The primary efficacy outcome evaluated was the change from baseline (on-medication state) to 3 months after the procedure in the hand tremor sub score in the Clinical Rating Scale for Tremor. Trial characteristics and results are summarized in Tables 3 and 4. After unblinding at 3 months, 6 of the 7 patients who received sham procedures crossed over to undergo open-label treatment with MRgFUS. The most common thalamotomy-related adverse events reported for all 26 patients treated were finger paresthesia (39%), ataxia (35%), and orofacial paresthesia (27%). Paresthesia and ataxia persisted to 1 year in 19% and 4% of patients, respectively. Eight severe adverse events were reported in 4 patients, and 3 were thalamotomy-related (2 patients with persistent mild hemiparesis and 1 patient had an associated persistent mild ataxia).

**Table 4. Summary of Key RCT Characteristics**

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Bond et al (2017)	US	2	2012 to 2015	27 patients with medication-refractory, severe, and disabling tremor-dominant PD	MRgFUS thalamotomy (n=20)	Sham treatment (n=7)

MRgFUS: magnetic resonance-guided ultrasound; PD: Parkinson disease; RCT: randomized controlled trial.

**Table 5. Summary of Key RCT Results**

Study	Hand Tremor Subscore	CRST
Bond et al (2017)	Percent change from baseline to month 3 (IQR)	Percent change from baseline to month 3 (IQR)
MRgFUS thalamotomy	62% (22.0 to 79.0)	44% (23.0 to 78.0)

Study	Hand Tremor Subscore	CRST
Sham treatment	22% (-11.0 to 29.0)	12% (-8.0 to 37.0)
Difference (p-value)	.04	

CRST: Clinical Rating Scale for Tremor; IQR: interquartile range; MRgFUS: magnetic resonance-guided ultrasound; RCT: randomized controlled trial.

Tables 6 and 7 summarize the relevance and conduct limitations of the RCT.

**Table 6. Study Relevance Limitations**

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Follow-Up <sup>e</sup>
Bond et al (2017)			2. Comparison to a sham treatment instead of an alternative surgical procedure		1. Efficacy evaluated through 3 months, limiting interpretation for long-term effects

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

<sup>b</sup> Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

<sup>c</sup> Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

<sup>d</sup> Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

<sup>e</sup> Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

**Table 7. Study Design and Conduct Limitations**

Study	Allocation <sup>a</sup>	Blinding <sup>b</sup>	Selective Reporting <sup>c</sup>	Data Completeness <sup>d</sup>	Power <sup>e</sup>	Statistical <sup>f</sup>
Bond et al (2017)				1. 3 of 20 patients who underwent the intervention received deep brain stimulation after 3 months in the open-label phase (additional detail not provided for these patients) 3. 6 of 7 patients receiving sham treatment crossed over after 3 months in the open-label phase	4. Study planned to enroll 30 patients, slow enrollment limited the study to 27 randomized patients	3. p-values not reported for efficacy outcomes other than hand tremor subscores

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

<sup>b</sup> Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician. 3. Blinding unclear

<sup>c</sup> Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>d</sup> Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

<sup>e</sup> Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Study did not meet conditions of power calculations

f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

## Section Summary: Tremor-Dominant Parkinson Disease

Evidence for the use of MRgFUS in the treatment of medication-refractory tremor-dominant PD consists of a double-blind, sham-controlled randomized trial and a systematic review. The sham-controlled randomized trial found significant improvements in the treatment group in tremor severity after 3 months of follow-up. Authors of the study noted that a larger study is needed to prove efficacy. The systematic review, which included the sham-controlled RCT plus additional small observational studies, found significant improvements in tremor severity through 12 months and quality of life through 6 months post procedure.

## SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

### Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

#### *American College of Radiology (ACR)*

In 2018, the American College of Radiology (ACR) issued appropriateness criteria for radiologic management of uterine leiomyomas that states: "MR [magnetic resonance]-guided high-intensity focused US [ultrasound] (MRgFUS) is another uterine-sparing option to treat focal leiomyomas. It is noninvasive, though each treatment may take several hours to complete. Its use currently is restricted to patients with fewer than six leiomyomas or leiomyoma volume < 900 cm<sup>3</sup>," and "although a reasonable alternative for patients unable or unwilling to tolerate sedation or anesthesia, long-term data and viability results are still lacking."

These appropriateness criteria were most recently updated in 2023, with evidence summaries provided for each reviewed clinical scenario. Table 8 summarizes the appropriateness category for specific populations with uterine fibroids.

**Table 8. ACR Appropriateness Criteria: Management of Uterine Fibroids**

Clinical situation	MRgFUS Appropriateness Category <sup>a</sup>
Reproductive age patient with uterine fibroids, symptomatic with heavy uterine bleeding or bulk symptoms (eg, pressure, pain, fullness, bladder, or bowel symptoms), and a desire to preserve fertility. Initial therapy.	Usually appropriate

<b>Clinical situation</b>	<b>MRgFUS Appropriateness Category<sup>a</sup></b>
Reproductive age patient with uterine fibroids, symptomatic with heavy uterine bleeding or bulk symptoms (eg, pressure, pain, fullness, bowel, or bladder symptoms), and no desire for future fertility. Initial therapy.	Usually appropriate
Reproductive age patient with uterine fibroids and concurrent adenomyosis, symptomatic with heavy uterine bleeding or bulk symptoms (eg, pressure, pain, fullness, bladder, or bowel symptoms), and no desire for future fertility. Initial therapy.	Usually not appropriate
Reproductive age patient with pedunculated submucosal uterine fibroids, symptomatic with heavy uterine bleeding. Initial therapy.	May be appropriate
Postmenopausal patient with uterine fibroids, symptomatic with heavy uterine bleeding or bulk symptoms (eg, pressure, pain, fullness, bladder, or bowel symptoms). Negative endometrial biopsy. Next step.	Usually not appropriate
Reproductive age patient with uterine fibroids desiring pregnancy and experiencing reproductive dysfunction. Initial therapy.	May be appropriate

ACR: American College of Radiology; MRgFUS: magnetic resonance-guided focused ultrasound.

<sup>a</sup>Usually appropriate: the imaging procedure or treatment is indicated in the specified clinical scenarios at a favorable risk-benefit ratio for patients; May be appropriate: The imaging procedure or treatment may be indicated in the specified clinical scenarios as an alternative to imaging procedures or treatments with a more favorable risk-benefit ratio, or the risk-benefit ratio for patients is equivocal; Usually not appropriate: The imaging procedure or treatment is unlikely to be indicated in the specified clinical scenarios, or the risk-benefit ratio for patients is likely to be unfavorable.

### ***American College of Obstetricians and Gynecologists (ACOG)***

In 2021, the American College of Obstetricians and Gynecologists (ACOG) reaffirmed in a practice bulletin on the management of symptomatic uterine leiomyoma states “while limited, low quality data suggests MRgFUS is associated with a reduction in leiomyoma and uterine size, smaller randomized comparative data suggests when compared with UAE, MRgFUS is associated with less improvement in symptoms and a higher rate of reintervention.”

### ***American Society of Radiation Oncology (ASTRO)***

In 2017, the American Society of Radiation Oncology (ASTRO) issued an updated evidence-based guideline on palliative radiation therapy for bone metastases. The updated guideline analysis confirms that radiation therapy provides excellent palliation for painful bone metastases and that retreatment is safe and effective. The guideline does not mention magnetic resonance-guided focused ultrasound. " A 2024 updated guideline, intended to replace the 2017 guideline, also does not mention MRgFUS. Recommendations for advanced radiotherapy such as stereotactic body radiotherapy (SBRT) are as follows:

- "For patients with symptomatic bone metastases treated with SBRT, 1200 to 1600 cGy in 1 fraction (nonspine) and 2400 cGy in 2 fractions (spine) are recommended."
- "For patients with symptomatic bone metastases with ECOG PS 0-2, receiving no surgical intervention, and absent neurological symptoms, SBRT is conditionally recommended over conventional palliative RT."

- "For patients with spine bone metastases that would benefit from reirradiation to the same site, treatment with SBRT is conditionally recommended."

In 2022, the American Urological Association (AUA)/ ASTRO published guidance on the management of clinically localized prostate cancer. The guidance states that "there is a lack of data to date to support the use of whole gland or focal ablation for the treatment of clinically localized prostate cancer."

### ***Congress of Neurological Surgeons***

In 2025, a guideline on the role of emerging therapies for patients with metastatic brain tumors was published by the Congress of Neurological Surgeons and endorsed by the American Association of Neurological Surgeons (AANS). The guideline states that: "For interstitial modalities and magnetic resonance imaging–guided focused ultrasound, insufficient qualifying data were identified to create recommendations." The guideline also states that: "There is insufficient evidence to make a recommendation regarding the use of HIFU [high-intensity focused ultrasound] for parenchymal and leptomeningeal brain metastases."

### ***National Comprehensive Cancer Network (NCCN)***

#### **Adult Cancer Pain Version 2.2025**

Some patients experience inadequate pain management despite pharmacologic therapy or may not tolerate an opioid titration program because of side effects. Some patients may prefer interventional therapies instead of chronic medication regimen. Interventional techniques have been demonstrated, in some cases, to eliminate or significantly reduce the level of pain, and/or may allow a significant decrease in systemic analgesics. Interventional therapies that can be useful in the relief of cancer pain include nerve blocks, vertebral augmentation, regional infusion of analgesics, image-guided ablation, and other techniques.

Ablation techniques may also be helpful for pain management in patients who receive inadequate relief from pharmacologic therapy. Prospective trials of percutaneous ablative techniques, many using thermal energy, have shown decreased patient pain from bone metastases in patients who did or did not receive prior radiation therapy.

Ablation therapy (e.g., image guided ablation, US ablation) for bone lesions can also be helpful in reducing pain.

#### **Bone Cancer Version 2.2025**

The current NCCN guideline does not mention the use of magnetic resonance-guided high intensity focused ultrasound (MRgFUS) as a treatment option for bone cancer.

#### **Breast Cancer Version 4.2025**

The current NCCN guideline does not mention the use of magnetic resonance-guided high intensity focused ultrasound (MRgFUS) as a treatment option for breast cancer.

### **Central Nervous System Cancers Version 2.2025**

The current NCCN guideline does not mention the use of magnetic resonance-guided high intensity focused ultrasound (MRgFUS) as a treatment option for central nervous system cancers (i.e., brain cancer to include glioblastoma).

### **Prostate Cancer Version 1.2026**

“Many Therapies have been investigated for the treatment of localized prostate cancer in the initial disease and recurrent settings with the goals of reducing side effects and mating the cancer control of other therapies. Cryotherapy or other local therapies are not recommended as routine primary therapy for localized prostate cancer due to lack of long-term data comparing these treatments to radiation or radical prostatectomy. At this time, the panel recommends only cryosurgery and high-intensity focused ultrasound (HIFU; category 2B) as local therapy options for RT recurrence in the absence of metastatic disease.”

### **Soft Tissue Sarcoma Version 1.2025**

The current NCCN guideline does not mention the use of magnetic resonance-guided high intensity focused ultrasound (MRgFUS) as a treatment option of soft tissue sarcomas (i.e., desmoid tumors).

### ***National Institute for Health and Care Excellence (NICE)***

In 2018, NICE issued an interventional procedure guidance (IPG617) on unilateral MRI-guided ultrasound thalamotomy for the treatment of resistant essential tremor that stated the following: The evidence on the safety of unilateral MRI-guided focused ultrasound thalamotomy for treatment-resistant essential tremor raises no major safety concerns. However, current evidence on its efficacy is limited in quantity. Therefore, this procedure should not be used unless there are special arrangements for clinical governance, consent, and audit or research.

### **Ongoing and Unpublished Clinical Trials**

Some currently ongoing and unpublished trials that might influence this review can be located at [clinicaltrials.gov](https://clinicaltrials.gov).

## **REFERENCES**

1. National Comprehensive Cancer Network (NCCN) Bone Cancer Version 2.2025. Also available at <https://www.nccn.org>
2. National Comprehensive Cancer Network (NCCN) Breast Cancer Version 4.2025. Also available at <https://www.nccn.org>
3. National Comprehensive Cancer Network (NCCN) Prostate Cancer Version 1.2026. Also available at <https://www.nccn.org>
4. National Comprehensive Cancer Network (NCCN) Central Nervous System Cancers Version 2.2025. Also available at <https://www.nccn.org>
5. National Comprehensive Cancer Network (NCCN) Adult Cancer Pain Version 2.2025. Also available at <https://www.nccn.org>
6. National Comprehensive Cancer Network (NCCN) Soft Tissue Sarcoma Version 1.2025. Also available at <https://www.nccn.org>

7. Eltoukhi HM, Modi MN, Weston M, et al. The health disparities of uterine fibroid tumors for African American women: a public health issue. *Am J Obstet Gynecol.* Mar 2014; 210(3): 194-9. PMID 23942040
8. Blue Cross Blue Shield Association Technology Evaluation Center (TEC). Magnetic resonance-guided focused ultrasound therapy for symptomatic uterine fibroids. *TEC Assessments.* 2005;Volume 20:Tab 10.
9. Barnard EP, AbdElmagied AM, Vaughan LE, et al. Periprocedural outcomes comparing fibroid embolization and focused ultrasound: a randomized controlled trial and comprehensive cohort analysis. *Am J Obstet Gynecol.* May 2017; 216(5): 500.e1-500.e11. PMID 28063909
10. Laughlin-Tommaso S, Barnard EP, AbdElmagied AM, et al. FIRSTT study: randomized controlled trial of uterine artery embolization vs focused ultrasound surgery. *Am J Obstet Gynecol.* Feb 2019; 220(2): 174.e1-174.e13. PMID 30696556
11. Jacoby VL, Kohi MP, Poder L, et al. PROMiSe trial: a pilot, randomized, placebo-controlled trial of magnetic resonance guided focused ultrasound for uterine fibroids. *Fertil Steril.* Mar 2016; 105(3): 773-780. PMID 26658133
12. Gizzo S, Saccardi C, Patrelli TS, et al. Magnetic resonance-guided focused ultrasound myomectomy: safety, efficacy, subsequent fertility and quality-of-life improvements, a systematic review. *Reprod Sci.* Apr 2014; 21(4): 465-76. PMID 23868442
13. Xu F, Deng L, Zhang L, et al. The comparison of myomectomy, UAE and MRgFUS in the treatment of uterine fibroids: a meta-analysis. *Int J Hyperthermia.* Sep 2021; 38(2): 24-29. PMID 34420449
14. Chen R, Keserci B, Bi H, et al. The safety and effectiveness of volumetric magnetic resonance-guided high-intensity focused ultrasound treatment of symptomatic uterine fibroids: early clinical experience in China. *J Ther Ultrasound.* 2016; 4: 27. PMID 27822376
15. Rabinovici J, David M, Fukunishi H, et al. Pregnancy outcome after magnetic resonance-guided focused ultrasound surgery (MRgFUS) for conservative treatment of uterine fibroids. *Fertil Steril.* Jan 2010; 93(1): 199-209. PMID 19013566
16. Otonkoski S, Sainio T, Mattila S, et al. Magnetic resonance guided high intensity focused ultrasound for uterine fibroids and adenomyosis has no effect on ovarian reserve. *Int J Hyperthermia.* 2023; 40(1): 2154575. PMID 36535925
17. Baal JD, Chen WC, Baal U, et al. Efficacy and safety of magnetic resonance-guided focused ultrasound for the treatment of painful bone metastases: a systematic review and meta-analysis. *Skeletal Radiol.* Dec 2021; 50(12): 2459-2469. PMID 34018007
18. Hurwitz MD, Ghanouni P, Kanaev SV, et al. Magnetic resonance-guided focused ultrasound for patients with painful bone metastases: phase III trial results. *J Natl Cancer Inst.* Apr 23 2014; 106(5). PMID 24760791
19. Arrigoni F, Barile A, Zugaro L, et al. Intra-articular benign bone lesions treated with Magnetic Resonance-guided Focused Ultrasound (MRgFUS): imaging follow-up and clinical results. *Med Oncol.* Apr 2017; 34(4): 55. PMID 28244018
20. Ghai S, Finelli A, Corr K, et al. MRI-guided Focused Ultrasound Ablation for Localized Intermediate-Risk Prostate Cancer: Early Results of a Phase II Trial. *Radiology.* Mar 2021; 298(3): 695-703. PMID 33529137
21. Ghai S, Finelli A, Corr K, et al. MRI-guided Focused Ultrasound Focal Therapy for Intermediate-Risk Prostate Cancer: Final Results from a 2-year Phase II Clinical Trial. *Radiology.* Mar 2024; 310(3): e231473. PMID 38441092
22. Ehdaie B, Tempany CM, Holland F, et al. MRI-guided focused ultrasound focal therapy for patients with intermediate-risk prostate cancer: a phase 2b, multicentre study. *Lancet Oncol.* Jul 2022; 23(7): 910-918. PMID 35714666

23. Zippel DB, Papa MZ. The use of MR imaging guided focused ultrasound in breast cancer patients; a preliminary phase one study and review. *Breast Cancer*. 2005; 12(1): 32-8. PMID 15657521
24. Hynynen K, Pomeroy O, Smith DN, et al. MR imaging-guided focused ultrasound surgery of fibroadenomas in the breast: a feasibility study. *Radiology*. Apr 2001; 219(1): 176-85. PMID 11274554
25. Gianfelice D, Khiat A, Amara M, et al. MR imaging-guided focused US ablation of breast cancer: histopathologic assessment of effectiveness-- initial experience. *Radiology*. Jun 2003; 227(3): 849-55. PMID 12714680
26. Gianfelice D, Khiat A, Amara M, et al. MR imaging-guided focused ultrasound surgery of breast cancer: correlation of dynamic contrast-enhanced MRI with histopathologic findings. *Breast Cancer Res Treat*. Nov 2003; 82(2): 93-101. PMID 14692653
27. Merkel LG, Knuttel FM, Deckers R, et al. First clinical experience with a dedicated MRI-guided high-intensity focused ultrasound system for breast cancer ablation. *Eur Radiol*. Nov 2016; 26(11): 4037-4046. PMID 26852219
28. McDannold N, Clement GT, Black P, et al. Transcranial magnetic resonance imaging-guided focused ultrasound surgery of brain tumors: initial findings in 3 patients. *Neurosurgery*. Feb 2010; 66(2): 323-32; discussion 332. PMID 20087132
29. Arrigoni F, Spiliopoulos S, de Cataldo C, et al. A Bicentric Propensity Score Matched Study Comparing Percutaneous Computed Tomography-Guided Radiofrequency Ablation to Magnetic Resonance-Guided Focused Ultrasound for the Treatment of Osteoid Osteoma. *J Vasc Interv Radiol*. Jul 2021; 32(7): 1044-1051. PMID 33775816
30. Arrigoni F, Napoli A, Bazzocchi A, et al. Magnetic-resonance-guided focused ultrasound treatment of non-spinal osteoid osteoma in children: multicentre experience. *Pediatr Radiol*. Aug 2019; 49(9): 1209-1216. PMID 31129699
31. Geiger D, Napoli A, Conchiglia A, et al. MR-guided focused ultrasound (MRgFUS) ablation for the treatment of nonspinal osteoid osteoma: a prospective multicenter evaluation. *J Bone Joint Surg Am*. May 07 2014; 96(9): 743-51. PMID 24806011
32. Avedian RS, Bitton R, Gold G, et al. Is MR-guided High-intensity Focused Ultrasound a Feasible Treatment Modality for Desmoid Tumors?. *Clin Orthop Relat Res*. Mar 2016; 474(3): 697-704. PMID 26040967
33. Bucknor MD, Rieke V. MRgFUS for desmoid tumors within the thigh: early clinical experiences. *J Ther Ultrasound*. 2017; 5: 4. PMID 28174660
34. Ghanouni P, Dobrotwir A, Bazzocchi A, et al. Magnetic resonance-guided focused ultrasound treatment of extra-abdominal desmoid tumors: a retrospective multicenter study. *Eur Radiol*. Feb 2017; 27(2): 732-740. PMID 27147222
35. Miller WK, Becker KN, Caras AJ, et al. Magnetic resonance-guided focused ultrasound treatment for essential tremor shows sustained efficacy: a meta-analysis. *Neurosurg Rev*. Feb 2022; 45(1): 533-544. PMID 33978922
36. Elias WJ, Lipsman N, Ondo WG, et al. A Randomized Trial of Focused Ultrasound Thalamotomy for Essential Tremor. *N Engl J Med*. Aug 25 2016; 375(8): 730-9. PMID 27557301
37. Giordano M, Caccavella VM, Zaed I, et al. Comparison between deep brain stimulation and magnetic resonance-guided focused ultrasound in the treatment of essential tremor: a systematic review and pooled analysis of functional outcomes. *J Neurol Neurosurg Psychiatry*. Dec 2020; 91(12): 1270-1278. PMID 33055140
38. Schaink A, Li C, Gajic-Veljanoski O, et al. Magnetic Resonance-Guided Focused Ultrasound Neurosurgery for Essential Tremor: A Health Technology Assessment. *Ont Health Technol Assess Ser*. 2018; 18(4): 1-141. PMID 29805721

39. Chang JW, Park CK, Lipsman N, et al. A prospective trial of magnetic resonance-guided focused ultrasound thalamotomy for essential tremor: Results at the 2-year follow-up. *Ann Neurol.* Jan 2018; 83(1): 107-114. PMID 29265546
40. Bond AE, Shah BB, Huss DS, et al. Safety and Efficacy of Focused Ultrasound Thalamotomy for Patients With Medication-Refractory, Tremor-Dominant Parkinson Disease: A Randomized Clinical Trial. *JAMA Neurol.* Dec 01 2017; 74(12): 1412-1418. PMID 29084313
41. Knuttinen MG, Stark G, Hohenwarter EJ, et al. ACR Appropriateness Criteria @ Radiologic Management of Uterine Leiomyomas. *J Am Coll Radiol.* May 2018; 15(5S): S160-S170. PMID 29724419
42. American College of Radiology. ACR Appropriateness Criteria: Management of Uterine Fibroids. Updated 2023. <https://acsearch.acr.org/docs/69508/Narrative/>. Accessed September 5, 2025.
43. Lutz S, Balboni T, Jones J, et al. Palliative radiation therapy for bone metastases: Update of an ASTRO Evidence-Based Guideline. *Pract Radiat Oncol.* 2017; 7(1): 4-12. PMID 27663933
44. Eastham JA, Boorjian SA, Kirkby E. Clinically Localized Prostate Cancer: AUA/ASTRO Guideline. *J Urol.* Sep 2022; 208(3): 505-507. PMID 35830561
45. UpToDate. Image guided ablation of skeletal metastases. Anil Nicholas Kurup M.D., Matthew R Callstrom M.D., PhD. Topic last updated May 2025. Also available at <https://www.uptodate.com>
46. UpToDate. Surgical treatment of essential tremor. Kelvin L Chou M.D., Topic last updated August 2025. Also available at <https://www.uptodate.com>
47. UpToDate. Uterine fibroids (leiomyomas): Treatment overview. Elizabeth Stewart M.D., Topic last updated August 2024. Also available at <https://www.uptodate.com>
48. Hayes, a symplr company. Health Technology Assessment Magnetic Resonance-Guided Focused Ultrasound Therapy for Treatment of Uterine Fibroids. September 2022
49. Hayes, a symplr company. Health Technology Assessment Magnetic Resonance-Guided Focused Ultrasound Unilateral Thalamotomy for Essential Tremor. February 2025
50. Mortezaei A, Essibayi MA, Mirahmadi Eraghi M, et al. Magnetic resonance-guided focused ultrasound in the treatment of refractory essential tremor: a systematic review and meta-analysis. *Neurosurg Focus.* Sep 01 2024; 57(3): E2. PMID 39217634
51. Monteiro JDS, E Silva BB, de Oliveira RR, et al. Magnetic resonance-guided focused ultrasound ventral intermediate thalamotomy for Tremor-Dominant Parkinson's disease: a systematic review and meta-analysis. *Neurosurg Rev.* Sep 27 2024; 47(1): 701. PMID 39331247

## CODES

To report provider services, use appropriate CPT codes, HCPCS codes, Revenue codes, and/or ICD diagnosis codes.

Codes	Number	Description
CPT		

<b>Codes</b>	<b>Number</b>	<b>Description</b>
	0071T	Focused ultrasound ablation of uterine leiomyomata, including MR guidance; total leiomyomata volume less than 200 cc of tissue
	0072T	Focused ultrasound of uterine leiomyomata, including MR guidance, total; leiomyomata volume greater or equal to 200 cc or tissue
	0947T	Magnetic resonance image guided low intensity focused ultrasound (MRgFUS), stereotactic blood-brain barrier disruption using microbubble resonators to increase the concentration of blood-based biomarkers of target, intracranial, including stereotactic navigation and frame placement, when performed
	61715	Magnetic resonance image guided high intensity focused ultrasound (MRgFUS), stereotactic ablation of target, intracranial, including stereotactic navigation and frame placement, when performed
HCPCS		
	None	
Type of Service	Medicine	
Place of Service	Inpatient/Outpatient	

## POLICY HISTORY

<b>Date</b>	<b>Action</b>	<b>Action</b>
September 2025	Annual Review	Policy Renewed
September 2024	Annual Review	Policy Renewed
September 2023	Annual Review	Policy Renewed
January 2023	Annual Review	Policy Revised
January 2022	Annual Review	Policy Renewed
January 2021	Annual Review	Policy Renewed
January 2020	Annual Review	Policy Renewed
January 2019	Annual Review	Policy Revised

<b>Date</b>	<b>Action</b>	<b>Action</b>
January 2018	Annual Review	Policy Revised
January 2017	Annual Review	Policy Renewed
January 2016	Annual Review	Policy Revised
February 2015	Annual Review	Policy Revised
March 2014	Annual Review	Policy Revised
May 2013	Annual Review	Policy Renewed
May 2012	Annual Review	Policy Renewed
August 2011	Annual Review	Policy Renewed

New information or technology that would be relevant for Wellmark to consider when this policy is next reviewed may be submitted to:

Wellmark Blue Cross and Blue Shield  
 Medical Policy Analyst  
 PO Box 9232  
 Des Moines, IA 50306-9232

\*CPT® is a registered trademark of the American Medical Association.