

09.03.02 Vision Therapy/Orthoptic Training

Original Effective Date: March 1988

Review Date: June 2025

Revised: May 2023

DISCLAIMER/INSTRUCTIONS FOR USE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations, or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

This Medical Policy document describes the status of medical technology at the time the document was developed. Since that time, new technology may have emerged, or new medical literature may have been published. This Medical Policy will be reviewed regularly and updated as scientific and medical literature becomes available; therefore, policies are subject to change without notice.

Related Policies:

- None

Summary

Description

Orthoptics training refers to techniques designed to correct accommodative and convergence insufficiency (or convergence dysfunction). Regimens may include push-up exercises using an accommodative target of letters, numbers, or pictures; push-up exercises with additional base-out prisms; jump-to-near convergence exercises; stereogram convergence exercises; and/or recession from a target. In addition to its use to treat convergence insufficiency, orthoptic training has been investigated for treating attention deficit disorders, dyslexia, and dysphasia.

This evidence review addresses office-based orthoptic training. It does not address standard vision therapy with lenses, prisms, filters, or occlusion (i.e., for treatment of amblyopia or acquired esotropia prior to surgical intervention).

Summary of Evidence

For individuals who have convergence insufficiency who receive office-based orthoptic training, the evidence includes a TEC Assessment, systematic reviews, several randomized controlled trials (RCTs), and nonrandomized comparative studies. Relevant outcomes are symptoms and functional outcomes. The most direct evidence on office-based orthoptic training comes from a 2008 RCT that demonstrated that office-based vision or orthoptic training improves symptoms of convergence insufficiency in a greater percentage of patients than a home-based vision exercise program consisting of pencil push-ups or home computer vision exercises. Subgroup analyses of this RCT demonstrated improvements in accommodative vision, parental perception of academic behavior, and specific convergence insufficiency-related symptoms. However, in this trial, as in others, the home-based regimen did not include the full range of home-based therapies, which may have biased results in favor of the orthoptic training. Another RCT published in 2019 did not find a difference in symptoms of convergence insufficiency between office-based orthoptic training plus home exercises and office-based placebo therapy plus home exercises, possibly due to notable improvements in symptoms in the placebo group. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome. However, even though there is a paucity in the peer reviewed scientific evidence in showing consistent evidence regarding the effectiveness of orthoptic training/vision therapy for convergence insufficiency, for certain carefully-selected individuals this treatment is widely considered to be in accordance with generally accepted standards of medical practice in the United States and will be considered medically necessary when the criteria below are met, see [Policy](#).

For individuals who have learning disabilities who receive office-based orthoptic training, the evidence includes nonrandomized comparative and noncomparative studies. Relevant outcomes are functional outcomes. Studies have not directly demonstrated improvements in reading or learning outcomes with orthoptic training. At least 2 earlier studies that addressed other types of vision therapies have reported mixed improvements in reading. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Additional Information

Clinical input was sought to help determine whether the use of orthoptic training for individuals with convergence insufficiency or learning disabilities would provide an improvement in net health outcome and whether the use is consistent with generally accepted medical practice. In response to requests, input was received from 4 physician specialty societies (5 reviewers) and 3 academic medical centers while this policy was under review in 2011.

Clinical input supported the use of office-based orthoptic training when home-based therapy has failed. Therefore, orthoptic training may be considered medically necessary in patients with convergence insufficiency whose symptoms have failed to improve with a home-based treatment trial of at least 12 weeks. Home-based therapy should include push-up exercises using an accommodative target, push-up exercises with additional base-out prisms, jump-to-near convergence exercises, stereogram convergence exercises, recession from a target, and maintaining convergence for 30 to 40 seconds.

OBJECTIVE

The objective of this evidence review is to evaluate whether the use of office-based orthoptic training improves the net health outcome for individuals with vision or learning disabilities.

PRIOR APPROVAL

Not applicable.

POLICY

Note: See also *Wellmark Provider Guide to Billing for Eye Care: Chapter 3: Noncovered Eye Care Services - Vision Therapies*.

Medically Necessary

- Office based orthoptic training/vision therapy to establish a home program may be considered **medically necessary** for the treatment of symptomatic convergence insufficiency up to 12 visits regardless of age.
- Office based orthoptics training/vision therapy for the treatment of convergence insufficiency regardless of age exceeding 12 visits will be considered **not medically necessary**.

Investigational

Orthoptics training/vision therapy is considered **investigational** for all other conditions, including but not limited to the following because the evidence is insufficient in determining that the technology results in improved net health outcomes:

- Visual disorders other than convergence insufficiency
- Dyslexia
- Learning disabilities
- Reading disabilities

POLICY GUIDELINES

Published clinical studies show that a limited number of office visits, usually less than twelve, are required for resolution of convergence insufficiency. Office visits are only necessary to help establish or support a home program.

A diagnosis of convergence insufficiency is based on asthenopic symptoms (sensations of visual or ocular discomfort) at near point combined with difficulty sustaining convergence.

Convergence insufficiency and stereoacuity are documented by:

- Exodeviation at near vision at least 4 prism diopters greater than at far vision AND
- Insufficient positive fusional vergence at near (positive fusional vergence <15 prism diopters blur or break) on positive fusional vergence testing using a prism bar AND
- Near point of convergence break of more than 6 cm AND
- Appreciation by the patient of at least 500 seconds of arc on stereoacuity testing.

Coding

See the [Codes table](#) for details.

BACKGROUND

Treatment

Orthoptic training refers to techniques designed to correct accommodative and convergence insufficiency (or convergence dysfunction), which may include push-up exercises using an accommodative target of letters, numbers, or pictures; push-up exercises with additional base-out prisms; jump-to-near convergence exercises; stereogram convergence exercises; and recession from a target. A related but distinct training technique is behavioral or perceptual vision therapy, in which eye movement and eye-hand coordination training techniques are used to improve learning efficiency by optimizing visual processing skills.

In addition to its use in the treatment of accommodative and convergence dysfunction, orthoptic training is being investigated for the treatment of attention deficit disorders, dyslexia, dysphasia, and reading disorders.

Regulatory Status

Not applicable.

RATIONALE

This evidence review was created in March 1988 and has been updated regularly with searches of the PubMed database. The most recent literature update was performed through June 2025.

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to individuals and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and

conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice. The following is a summary of the key literature to date.

Orthoptic Training for Convergence Insufficiency

Clinical Context and Therapy Purpose

Convergence insufficiency is a binocular vision disorder associated with defects in the eyes' ability to turn inward toward each other (e.g., when looking at near objects). The diagnosis of convergence insufficiency is made when individuals have a remote near point of convergence or difficulty in sustaining convergence in conjunction with sensations of visual or ocular discomfort at near vision. Symptoms of this common condition may include eyestrain, headaches, blurred vision, diplopia, sleepiness, difficulty concentrating, movement of print, and loss of comprehension after short periods of reading or performing close activities. Prism reading glasses, home therapy with pencil push-ups, and office-based vision therapy and orthoptics have been evaluated for the treatment of convergence insufficiency.

The purpose of orthoptic training in individuals who have convergence insufficiency is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with convergence insufficiency.

Interventions

The treatment being considered is in-office orthoptic training. Orthoptic training refers to techniques designed to correct accommodative and convergence insufficiency (or convergence dysfunction).

Comparators

The comparator of interest is standard management of convergence insufficiency with at-home vision training exercises.

Outcomes

The general outcomes of interest are symptoms and functional outcomes.

Timing of intervention is approximately 12 weeks of in-office training, followed by 6 months of at-home training. Follow-up at 1 year or more is preferable.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

At least 2 systematic reviews have addressed the role of orthoptic training for convergence insufficiency. A systematic review by Rawston et al (2005) assessing the applicability and efficacy of eye exercises found that small controlled trials and a large number of cases supported their use in the treatment of convergence insufficiency. Scheiman et al (2020) conducted a systematic review and network meta-analysis of RCTs that evaluated nonsurgical treatments for convergence insufficiency. Six trials in children (n=968) were analyzed. When treatment success was defined as a composite of normal clinical convergence parameters and a prespecified magnitude of improvement, office-based vergence/accommodative (orthoptic) training with home reinforcement was more likely to lead to a successful outcome than home-based computer training (risk ratio, 1.96; 95% confidence interval [CI], 1.32 to 2.94) and home-based pencil/target push-up training (risk ratio, 2.86; 95% CI, 1.82 to 4.35). An analysis that defined treatment success as a composite of both improved convergence parameters and improved symptoms found that office-based training with home reinforcement was more effective than home-based computer training (risk ratio, 4.65; 95% CI, 1.23 to 17.54) or home-based pencil push-up training (risk ratio, 4.41; 95% CI, 1.26 to 15.38); however, these findings were based on low-certainty evidence. Six RCTs in adults were included, but none compared office-based and home-based orthoptic training. Three trials in adults compared office-based training to placebo; results were limited, and the authors concluded that the benefit of orthoptic training in adults was less clear overall than in children.

Randomized Controlled Trials

In 2008, the Convergence Insufficiency Treatment Trial Study Group reported on an RCT of 221 children with symptomatic convergence insufficiency. Symptoms were evaluated by the Convergence Insufficiency Symptom Survey, a 15-item survey with a final score ranging from 0 (least symptomatic) to 60 (most symptomatic). Scores of less than 16 were considered “asymptomatic,” and a decrease of 10 or more points was considered “improved.” On blinded evaluation after 12 weeks of treatment (99% completion rate), 73% of patients treated with office-based therapy were considered to be successful or improved on the composite outcome of Convergence Insufficiency Symptom Survey, near point convergence, and positive fusional vergency, as defined above, compared with 43%, 33%, and 35% of those treated with home pencil push-ups, home computer exercise, or placebo, respectively. At 1-year follow-up, 88% of the 32 children who were asymptomatic at the completion of the 12-week office-based treatment program remained successful or improved; 67% of the home-based pencil push-up group remained successful or improved. A limitation of this RCT is that near point exercises generally consisted of multiple therapies, making it difficult to correlate outcomes with specific modalities.

Following the publication of the main results of the Convergence Insufficiency Treatment Trial, a number of reanalysis were performed. The effectiveness of these forms of vision therapy (pencil push-ups, home computer exercises, office-based vision therapy) in improving accommodative amplitude in 164 (74%) of the 221 children who had coexisting accommodative dysfunction with convergence insufficiency was

reported by the Convergence Insufficiency Treatment Trial Study Group in 2011. Of the 164 children with accommodative dysfunction, 63 (29%) had a decreased amplitude of accommodation, 43 (19%) had decreased accommodative facility (latency and speed of the accommodative response), and 58 (26%) had both. After 12 weeks of treatment, increases in amplitude of accommodation were significantly greater in the 3 active groups (range, 5.8-9.9 diopters) compared with office-based placebo therapy (2.2 diopters). The percentage of children who no longer showed decreased amplitude of accommodation was 91.4% for office-based therapy, 79.3% for home computer therapy, 74.1% for home pencil push-ups, and 35.7% for placebo treatment. Accommodative facility improved by 9.4 cycles per minute for office-based therapy, 7.0 cycles per minute for home computer-based therapy, 5.0 cycles per minute for home pencil push-ups, and 5.5 cycles per minute for office-based placebo therapy; only the office-based therapy showed significantly greater improvement than office-based placebo therapy. One year after completion of therapy, recurrence of decreased accommodative amplitude was found in 5 (11%) of 44 children and in 4 (12.5%) of 32 children who did not undergo subsequent treatment.

The effect of successful treatment for convergence insufficiency on parents' perception of academic behavior in the 218 children who completed this trial was also reported by the Convergence Insufficiency Treatment Trial group (2012). Participants were classified as successful (n=42), improved (n=60), or nonresponder (n=116) after 12 weeks of treatment. This study used the Academic Behavior Survey, a 6-item questionnaire (scoring range, 0-24 points) developed by the Convergence Insufficiency Treatment Trial Study Group to quantify parents' perceptions of the frequency of adverse behaviors exhibited by children when reading or performing school work (5 questions) and overall parental concern about the child's academic performance (1 question). Mean Academic Behavior Survey score at baseline was 12.85 points, which improved by 4.0, 2.9, and 1.3 points in children classified as successful, improved, and nonresponder, respectively. Improvements in Academic Behavior Survey scores correlated with reductions in symptom level ($r=0.29$), but not changes in measures of convergence. Although the Academic Behavior Survey has not been validated outside of this study, the effect sizes in the successful and improved groups were 0.9 and 0.7, representing a clinically meaningful change.

In 2012, the Convergence Insufficiency Treatment Trial Study Group reported on a post hoc analysis of this RCT evaluating the effect of convergence insufficiency treatment on specific types of symptoms. Outcomes were measures on the Convergence Insufficiency Symptom Survey, which has 2 subscales: a performance-related subscale consisting of 6 symptoms related to visual efficiency when reading or performing near work (e.g., loss of place with reading) and an eye-related subscale consisting of 9 symptoms specific to visual function or asthenopic-type complaints (e.g., eye pain). Those with a "treatment response" (improvement of at least 8 points) on the overall Convergence Insufficiency Symptom Survey score demonstrated improvements in both the performance-related subscale and the eye-related subscale (mean, 1.1 points). Further research is needed to determine whether the treatment-related improvement in performance-related symptoms seen with orthoptics training translates into improvements in reading performance and attention.

In 2019, results of the Convergence Insufficiency Treatment Trial - Attention & Reading Trial (CITT-ART) were published. Children with convergence insufficiency were randomized to 16 weeks of weekly office-based vergence/accommodative therapy or office-based placebo therapy. Both groups performed home exercises 15 minutes per day, 5 days per week. The study outcomes for convergence ability and symptoms were the same as the outcomes in the Convergence Insufficiency Treatment Trial. After 16 weeks, mean Convergence Insufficiency Symptom Survey scores had decreased from baseline by -11.8 (95% CI, -13.4 to -10.3) and -10.4 (95% CI, -12.4 to -8.4) in the therapy and placebo groups, respectively, which was statistically similar between groups. There was no difference in the proportion of patients in each group that achieved normal or improved symptoms. Significantly more patients in the therapy group versus the placebo group met the criteria for normal or improved near point of convergence ($p<.001$) and

positive fusional vergence ($p < .001$). Several composite outcomes for treatment success found significant improvements with therapy versus placebo. Interpretation of the symptom comparisons in this trial may be limited by the clinically relevant improvement in symptoms in the placebo group. Results for accommodation were published separately by Chen et al (2020).¹⁰ Among the 288 children in the CITT-ART study with decreased accommodative amplitude or facility, normal amplitude (69% vs. 32%; $p < .0001$) and facility (85% vs. 49%; $p < .0001$) were achieved by significantly more patients in the therapy group compared to the placebo group, respectively. In a separate publication, results for improvement in reading comprehension were not significantly different between the therapy and placebo groups. Reading comprehension subtest scores of the Wechsler Individual Achievement Test, Third Edition (WIAT-III) increased by 3.68 points in the therapy group and 3.8 points in the placebo group (difference -0.12; 95% CI, -1.89 to 1.66). All other reading outcome measures were also similar between groups.

Singh et. al. (2021) published results of a randomized controlled trial (RCT) in 176 children and young adults (aged 9 to 30 years, mean 19 years) with symptomatic convergence insufficiency. Patients were randomized to 6 weeks of office-based orthoptic therapy (3 times per week) or home-based pencil push-up exercises (15 minutes per day). At study end, there was no difference between groups in near point of convergence or Convergence Insufficiency Symptom Survey scores, but there was a significantly greater improvement in positive fusional vergence with office-based therapy compared to home-based exercises ($p < 0.001$). Limitations of this study include lack of blinding, a wide range of patient ages, short duration compared to other studies, 20% to 30% loss to follow-up leading to a lack of power, and the study was conducted at a single center in India.

In 2020, Alvarez et. al., conducted the Convergence Insufficiency Neuro-mechanism in Adult Population Study, a small RCT (N=50) that compared 6 weeks of twice weekly office-based vergence/accommodation therapy and office-based placebo therapy in young adults (aged 18 to 35 years) with symptomatic convergence insufficiency.¹³ All patients performed home-based computer exercises 10 minutes per day, 3 days per week. Outcomes included change in near point of convergence, positive fusional vergence, and Convergence Insufficiency Symptom Survey scores. Both near point of convergence ($p < 0.01$) and positive fusional vergence ($p < 0.001$) were significantly improved with office-based therapy compared to placebo, but there was no difference between groups in symptom scores (2.3 points; 95% CI, -8.3 to 4.6; $p = 0.6$).

Tables 1 and 2 summarize the key RTCs in individuals with convergence insufficiency.

Table 1. Summary of Key Randomized Controlled Trial Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Convergence Insufficiency Treatment Trial Study Group (2008)	US	9	2005-2006	Children aged 9 to 17 years, exodeviation at or near at least 4Δ greater than at distance, insufficient positive fusional convergence, receded near point of convergence of ≥6 cm break, best corrected visual acuity of 20/25 in both eyes at distance and near, Convergence Insufficiency Symptom Survey score ≥16, not previously treated	Office-based vergence therapy with home reinforcement; weekly visits and 15 minutes/day, 5 days/week for 12 weeks; n=60	Home-based pencil push-up therapy; 15 minutes/day, 5 days/week for 12 weeks; n=54 Home-based computer vergence therapy with pencil push-ups;

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
				with pencil push-up or vergence orthoptic therapy		20 minutes/day, 5 days/week for 12 weeks; n=53 Office-based placebo therapy with home reinforcement; weekly visits and 15 minutes/day, 5 days/week for 12 weeks; n=54
CITT-ART Investigator Group (2019); CITT-ART	US	9	2014-2017	Children aged 9 to 14 years with exodeviation at or near at least 4Δ greater than at distance, insufficient positive fusional convergence, receded near point of convergence of ≥6 cm break, best corrected visual acuity of 20/25 in both eyes at distance and near, Convergence Insufficiency Symptom Survey score ≥16, not previously treated with office-based or home-based vergence therapy	Office-based vergence therapy with home reinforcement; weekly visits and 15 minutes/day, 5 days/week for 16 weeks; n=206	Office-based placebo therapy with home reinforcement; weekly visits and 15 minutes/day, 5 days/week for 16 weeks; n=104

CITT-ART: Convergence Insufficiency Treatment Trial – Attention & Reading Trial; RCT: randomized controlled trial.

Table 2. Summary of Key Randomized Controlled Trial Results

Study	Change in Convergence Insufficiency Survey Symptom score ^a (mean, 95% CI)	Symptoms resolved or improved at end of study ^b	Convergence ability normal or improved at end of study ^c
Convergence Insufficiency Treatment Trial Study Group (2008)	N=221	N=218	N=218
Office-based vergence therapy	-14.8 (-17.2 to -12.4)	72.9%	72.9-95.0%
Home-based pencil push-up therapy	-7.1 (-9.6 to -4.5)	47.1%	56.6-77.4%
Home-based computer vergence therapy with pencil push-ups	-6.0 (-8.6 to -3.4)	38.5%	59.6-77.0%
Office-based placebo therapy	-7.8 (-10.4 to -5.3)	42.6%	44.5-59.3%
Mean difference (95% CI); p-value	Office-based vergence therapy vs. home-based pencil push-ups: 7.9 (4.4 to 11.4); p<0.001 Office-based vergence therapy vs. home-based computer vergence therapy with pencil push-ups: 8.4 (4.9 to 11.9); p<0.001	Office-based vergence therapy vs. home-based pencil push-ups: p=0.008 Office-based vergence therapy vs. home-based computer vergence therapy with pencil push-ups: p=0.006	Office-based vergence therapy vs. home-based pencil push-ups: p<0.05 Office-based vergence therapy vs. home-based computer vergence therapy with pencil push-ups: p<0.05

Study	Change in Convergence Insufficiency Survey Symptom score ^a (mean, 95% CI)	Symptoms resolved or improved at end of study ^b	Convergence ability normal or improved at end of study ^c
CITT-ART Investigator Group (2019); CITT-ART	N=311	N=311	N=311
Office-based vergence therapy	-11.8	61.8%	92.4-95.5%
Office-based placebo therapy	-10.4	58.7%	50.0-67.3%
Mean difference (95% CI); p-value	1.5 (-3.8 to 0.8); p=0.21	p=NS	p<0.001

CI: confidence interval; CITT-ART: Convergence Insufficiency Treatment Trial – Attention & Reading Trial; NS: not significant.

^a Based on the Convergence Insufficiency Symptom Survey, a 15-item survey with scores ranging from 0 (least symptomatic) to 60 (most symptomatic). A score of <16 is considered asymptomatic, and a decrease of ≥10 points is considered improved.

^b Asymptomatic (Convergence Insufficiency Symptom Score <16) or improved (change in Convergence Insufficiency Symptom Score ≥10 points).

^c Based on near point convergence and positive fusional vergency. A “normal” near point convergence was defined as <6 cm, and an improved near point convergence was defined as an improvement (decrease) of >4 cm from baseline to follow-up. To be classified as having normal positive fusional vergency, a patient had to pass Sheard’s criteria (ie, positive fusional vergency blur, or if no blur, then a break value at least twice the near phoria magnitude) and have a positive fusional vergency blur/break of >15 prism diopters. Improvement in positive fusional vergency was defined as an increase of ≥10 prism diopters from baseline to follow-up.

Table 3 and 4 display notable limitations identified in each study.

Table 3. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Convergence Insufficiency Treatment Trial Study Group (2008)				3 - harms briefly described in text	
CITT-ART Investigator Group (2019); CITT-ART	3 - patients with learning or developmental disabilities were not excluded		2 - placebo involved some eye exercises and may have had a therapeutic effect		

CITT-ART: Convergence Insufficiency Treatment Trial - Attention & Reading Trial

The evidence limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 4. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Convergence Insufficiency Treatment Trial Study Group (2008)		1 - office vs. home therapy				
CITT-ART Investigator Group (2019); CITT-ART						

CITT-ART: Convergence Insufficiency Treatment Trial - Attention & Reading Trial

The evidence limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Nonrandomized Comparative Studies

Shin et al (2011) reported on a nonrandomized comparative study of office-based vision therapy. Fifty-seven children with symptomatic convergence insufficiency or combined convergence insufficiency and accommodative insufficiency were divided into a treatment and a sham control group, matched by age and sex. Vision therapy was performed in the school clinic 2 times a week with instructions for home exercises to be performed for 15 to 25 minutes a day during the week. After 12 weeks of office-based vision therapy, the mean College of Optometrists in Vision Development-Quality of Life questionnaire score decreased from 27.07 to 10.40, and near point convergence improved from 8.67 to 3.20 in the children with convergence insufficiency. Mean positive fusional vergency improved from 13.93 to 26.80. Sixty-seven percent of the children were considered to have been cured, and 82% were improved. There were no significant changes between baseline and 12-week follow-up for the control group. Of the 20 children in the treatment group who completed a 1-year follow-up, 3 (15%) showed recurrence.

Dusek et al (2011) reported on a nonrandomized comparative study of 134 children with convergence insufficiency who had been referred to a tertiary care center in Austria for reading difficulties. Thirty-two participants refused all treatment offered (control group); the remaining children were given base-in prism reading glasses (n=51) or computerized home vision therapy (n=51) based on preference. Parents were instructed to ensure that their child carried out the procedure correctly; compliance was verified weekly. All participants were examined for total reading time, reading error score, the amplitude of accommodation, and binocular accommodative facility at baseline and after 4 weeks. Prismatic reading glasses were not worn during testing. Significant improvements were found in the prism glasses and computer exercise groups for total reading time, reading error score, the amplitude of accommodation, binocular accommodative facility, and vergence facility. For example, reading speed improved by 21 seconds in the reading glasses group, by 12 seconds in the computer exercise group, and by 4 seconds in the control group. Mean amplitude of accommodation improved by 1.4 diopters in the reading glasses group, by 1.0 diopters in the computer exercise group, and by 0.3 diopters in the control group. The only significant improvement for the control group was vergence facility. Although this nonrandomized study had the potential for selection and performance bias, the results suggested that base-in prism reading glasses might be an effective treatment for convergence insufficiency and associated reading problems in children.

Lee et al (2014) reported on results from a small nonrandomized, controlled trial of vision therapy in children with vergence insufficiency and symptomatic ADHD. Of 1,123 children (age range, 8-13 years) who were screened for ADHD, 81 were identified as having symptomatic ADHD; of those, 16 were identified as having accommodative dysfunction on binocular function testing. Eight subjects received vision therapy, and the remainder acted as a control group; eligibility criteria for vision therapy included: high exophoria at near vision (≥ 6 prism diopters), exophoria at near vision at least 4 prism diopters greater than at distant vision, a receded near point of convergence break (≥ 6 cm), or insufficient positive fusional vergency at near vision, failing Sheard's criterion (positive fusional vergency less than twice the near phorias), or a minimum positive fusional vergency of 15 prism diopters or less base-out blur or break. Vision therapy included progressive home- and office-based convergence and accommodative exercises

over 12 weeks. At the 12-week follow-up, intervention group subjects demonstrated improvements in near point convergence (11.50 to 4.38 cm; $p < 0.05$), breakpoint of near positive fusional vergency (11.88 to 32.38 cm; $p < 0.01$), recovery point of near positive fusional vergency (6.38 to 19.75 cm; $p < 0.01$), and near exophoria (12.00 to 7.81 cm; $p < 0.05$). ADHD symptoms, as measured by the parent-reported Korea-ADHD Rating Scale, improved from 23.25 at baseline to 17.13 ($p < 0.05$) after vision therapy. Only within-group comparisons were reported. Control group subjects did not demonstrate improvements in vision metrics or Korea-ADHD Rating Scale scores.

In a small randomized comparative study, Momeni-Moghaddam et al (2015) compared the effectiveness of pencil push-up therapy with office-based vision therapy in 60 individuals who had convergence insufficiency (mean age, 21.3 years). Subjects received either pencil push-up therapy or office-based therapy without home intervention and underwent reevaluation at 4 and 8 weeks after the start of treatment. With a single exception, the 2 groups did not differ significantly regarding the near point convergence, phoria, and positive fusional vergency. After 4 and 8 weeks of follow-up, positive fusional vergency was significantly more improved in the pencil push-up therapy group ($p = 0.001$). Study authors suggested that pencil push-up therapy and office-based vision therapy were largely comparable for treatment of convergence insufficiency.

Noncomparative Studies

Borsting et al (2016) published the results of a single-arm multicenter study, the Convergence Insufficiency Treatment Trial Reading Study. Investigators evaluated parent-reported behavioral and emotional problems at baseline among children with symptomatic convergence insufficiency and after 16 weeks of office-based vergence accommodative therapy. The intervention was consistent with that administered in the Convergence Insufficiency Treatment Trial. Parent-reported ADHD symptoms were assessed with the Conners 3 ADHD Index and behavioral and emotional symptoms with the 120-item Child Behavior Checklist. Of the 53 children enrolled, 48 consented to office-based therapy and 44 completed therapy and provided posttreatment data. After completion of therapy, there were significant within-subject improvements in Convergence Insufficiency Symptom Survey scores and Conners 3 ADHD Index scores ($d = 0.58$, significantly different from zero). Subjects also demonstrated statistically significant improvements in the Child Behavior Checklist competency-related subscale related to school performance but not to social- or activities-related performance. On Child Behavior Checklist's symptom-related subscales, there were statistically significant improvements in the anxious/depressed, somatic complaints, and internalizing problems subscales. This study provided some evidence that ADHD-like and emotional and behavior problems may improve among children with symptomatic convergence insufficiency after office-based vision therapies. However, the study's small size and lack of a control group preclude drawing definitive conclusions about the efficacy of this treatment.

Section Summary: Orthoptic Training for Convergence Insufficiency

At least 2 systematic reviews support the efficacy of orthoptic training for convergence insufficiency, especially in children. The most direct evidence on office-based orthoptic training comes from a 2008 RCT that demonstrated that office-based vision training improves symptoms of convergence insufficiency in a greater percentage of patients than a home-based vision exercise program. Subgroup analyses of this RCT demonstrated improvements in accommodative vision, parental perception of academic behavior, and specific convergence insufficiency-related symptoms. However, in this trial, as in others, the home-based regimen did not include the full range of home-based therapies, which may have biased results in favor of the orthoptic training. Another RCT published in 2019 did not find a difference in symptoms of convergence insufficiency between office-based orthoptic training plus home exercises and

office-based placebo therapy plus home exercises, possibly due to notable improvements in symptoms in the placebo group.

Orthoptic Training for Learning Disabilities

Clinical Context and Test Purpose

Some learning disabilities, particularly those in which reading is impaired, have been associated with deficits in eye movements and/or visual tracking. For example, many dyslexic individuals may have an unstable binocular vision and report that letters appear to move around, causing visual confusion.

The purpose of orthoptic training in individuals who have learning disabilities is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with learning disabilities, including attention deficit disorders, dyslexia, dysphasia, and reading disorders. Diagnosis of learning disabilities should be conducted by a qualified, licensed professional. Attention deficit disorder can be diagnosed by professionals qualified and licensed to do so, as well as by psychiatrists and physicians, although only medical doctors can prescribe medication.

Interventions

The treatment being considered is office-based orthoptic training for learning disabilities.

Comparators

The comparator of interest is standard management of learning disabilities. The practices currently being used to treat learning disabilities vary depending on the type of disability, but they could include receiving special services at school such as individualized education programs and accommodations.

Outcomes

The general outcome of interest is functional outcomes.

The limited available literature showed that approximately 12 sessions over 5 weeks are needed to assess results. Longer-term follow-up was not indicated.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.

- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Review

A 2005 systematic review evaluating the applicability and efficacy of eye exercises found no clear scientific evidence to support the use of eye exercises for other disorders (e.g., learning disabilities, dyslexia), except convergence insufficiency.

Randomized Controlled Trials

Two studies focused on the use of tinted lenses and eye patching as a technique to steady binocular vision for dyslexia. Stein et al (2000) reported on results of a randomized trial in which 143 dyslexic children were instructed to wear yellow-tinted glasses with or without the left lens occluded. Children were instructed to wear these glasses when reading or writing. Significantly more children given occluded glasses gained stable binocular vision in the first 3 months (59%) compared with children given unoccluded glasses (36%). Christenson et al (2001), however, found no difference in reading ability of children with dyslexia and abnormal binocular vision tested with and without occluded, blue-tinted lenses.

Nonrandomized Comparative Studies

Ramsay et al (2014) reported on results from a non-RCT assessing a computerized vergence training program in 13- to 14-year-old patients with dyslexia. Twelve subjects with dyslexia were treated with the computerized vergence training program, receiving an average of 11.75 sessions over 5 weeks; 12 control students included were not treated. All subjects underwent vision testing and were not diagnosed with convergence insufficiency. The computerized training program involved the generation of a computerized stereogram, which appears in 3 dimensions with convergent vision. For the intervention groups, reading speed improved from 87.83 to 95.58 words read per minute from baseline to follow-up ($p < 0.006$); reading speed was unchanged from baseline to follow-up for the control group (85.00 words per minute at baseline to 89.37 words per minute at follow-up; $p < 0.123$). Mean improvement in reading speed from baseline to follow-up did not differ significantly between groups ($p < 0.123$).

Several studies have reported that poor reading in children with dyslexia or attention deficits may be related to impairments in accommodation or convergence, suggesting the need for an ophthalmologic and orthoptic evaluation.

Section Summary: Orthoptic Training for Learning Disabilities and Miscellaneous Indications

Peer-reviewed studies have not directly demonstrated improvements in reading or learning outcomes with orthoptic training. At least 2 earlier studies that addressed other types of vision therapies reported mixed improvements in reading.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Clinical Input from Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

In response to requests, input was received from 4 physician specialty societies (5 reviewers) and 3 academic medical centers while this policy was under review in 2011. Although input supported the use of office-based orthoptic training when home-based therapy had failed, some reviewers indicated that home-based therapy would typically include more exercises than pencil push-ups. Recommended were push-up exercises using an accommodative target, push-up exercises with additional base-out prisms, jump-to-near convergence exercises, stereogram convergence exercises, recession from a target, and maintaining convergence for 30 to 40 seconds.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Academy of Ophthalmology

Convergence Insufficiency

“Orthoptic training (vision therapy) is the primary treatment modality used by most eye care professionals for the treatment of convergence insufficiency (CI). The plasticity of the fusional convergence reflex system allows patients to improve their convergence amplitudes with simple exercises. There are numerous different types of eye exercises; however, the primary treatment modalities for CI include home-based exercise, in-office exercises, computer vergence exercises or a combination of these. Some studies have suggested that performing both the home-based exercises and the computer program is more effective than either modality performed on its own.

Eye care professionals will sometimes prescribe both in-office and home-based CI exercises. Occasionally patients will require additional treatment strategies such as anti-suppression or extra time and assistance with the exercises and will require in-office treatment.

Conventional Convergence Exercises:

- Gradual convergence exercises
- Convergence cards
- Stereograms
- Vergence facility exercises
- Base out prism exercises
- Computer based convergence exercises.

Idiopathic convergence insufficiency responds very well to convergence exercises and has a very high reported success rate. Published success rates vary between 70 to 80% depending on the patient population and study size.”

American Optometric Association

In 2010, the American Optometric Association issued an optometric clinical practice guideline for the care of the patient with accommodative and vergence dysfunction that stated: “Patients with convergence insufficiency (CI) can be treated by various strategies, depending on the severity of symptoms. Numerous studies have shown that vision therapy is the treatment of choice for CI. The recommended treatment includes in-office therapy and supplemental home therapy. Home therapy alone, which is less effective, may be prescribed when in-office therapy is not possible. To ensure its success, home therapy should be closely monitored for patient compliance and make adjustment when needed. For the patient who cannot participate in vision therapy, prescribed prisms may reduce the load on the vergence system, however, prisms do not always alleviate the patient’s symptoms.”

American Academy of Pediatrics (AAP), American Academy of Ophthalmology (AAO), American Association for Pediatric Ophthalmology and Strabismus (AAPOS), and American Association of Certified Orthoptists (AAO)

In 2009 (reaffirmed in 2014), the American Academy of Pediatrics, American Academy of Ophthalmology, American Association for Pediatric Ophthalmology and Strabismus, and the American Association of Certified Orthoptists issued a joint statement concerning pediatric learning disabilities, dyslexia and vision. For vision therapy, the statement concluded:

“Currently, there is no adequate scientific evidence to support the view that subtle eye or visual problems cause learning disabilities. Furthermore, the evidence does not support the concept that vision therapy or tinted lenses or filters are effective, directly or indirectly, in the treatment of learning disabilities. Thus, the claim that vision therapy improves visual efficiency cannot be substantiated. Diagnostic and treatment approaches that lack scientific evidence of efficacy are not endorsed or recommended.”

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review can be located at clinicaltrials.gov.

REFERENCES

1. Handler SM, Fierson WM, Section on Ophthalmology SM. Learning disabilities, dyslexia, and vision. *Pediatrics*. Mar 2011; 127(3): e818-56. PMID 21357342
2. Rawstron JA, Burley CD, Elder MJ. A systematic review of the applicability and efficacy of eye exercises. *J Pediatr Ophthalmol Strabismus*. 2005; 42(2): 82-8. PMID 15825744
3. Convergence Insufficiency Treatment Trial Study Group. Randomized clinical trial of treatments for symptomatic convergence insufficiency in children. *Arch Ophthalmol*. Oct 2008; 126(10): 1336-49. PMID 18852411

4. Scheiman M, Kulp M, Cotter S, et al. Long-term effectiveness of treatments for symptomatic convergence insufficiency in children. *Optom Vis Sci.* Sep 2009; 86(9): 1096-103. PMID 19668097
5. Scheiman M, Cotter S, Kulp MT, et al. Treatment of accommodative dysfunction in children: results from a randomized clinical trial. *Optom Vis Sci.* Nov 2011; 88(11): 1343-52. PMID 21873922
6. Borsting E, Mitchell GL, Kulp MT, et al. Improvement in academic behaviors after successful treatment of convergence insufficiency. *Optom Vis Sci.* Jan 2012; 89(1): 12-8. PMID 22080400
7. Barnhardt C, Cotter SA, Mitchell GL, et al. Symptoms in children with convergence insufficiency: before and after treatment. *Optom Vis Sci.* Oct 2012; 89(10): 1512-20. PMID 22922781
8. Institute For Clinical Systems Improvement's Technical Assessment Report TA #68; Vision Therapy, January 2003
9. Momeni-Moghaddam H, Kundart J, Azimi A, et al. The effectiveness of home-based pencil push-up therapy versus office-based therapy for the treatment of symptomatic convergence insufficiency in young adults. *Middle East Afr J Ophthalmol.* 2015; 22(1): 97-102. PMID 25624682
10. Grisham D, Powers M, Riles P. Visual skills of poor readers in high school. *Optometry.* Oct 2007; 78(10): 542-9. PMID 17904495
11. Palomo-Alvarez C, Puell MC. Accommodative function in school children with reading difficulties. *Graefes Arch Clin Exp Ophthalmol.* Dec 2008; 246(12): 1769-74. PMID 18751994
12. Ponsonby AL, Williamson E, Smith K, et al. Children with low literacy and poor stereoacuity: an evaluation of complex interventions in a community-based randomized trial. *Ophthalmic Epidemiol.* 2009; 16(5): 311-21. PMID 19874111
13. National Guideline Clearinghouse; Esotropia and exotropia. [NGC:2770] March 2003
14. Ziring PR, et al. Learning Disabilities, Dyslexia, and Vision: A Subject Review (RE9825). American Academy of Pediatrics Policy Statement Volume 102, No 5 November 1998, 1217-1219.
15. Robinson GL, Foreman PJ. Scotopic sensitivity/Irlen Syndrome and the use of coloured filters: a long-term placebo controlled and masked study of reading achievement and perception of ability. *Perceptual and Motor Skills* 1999 August; 89(1): 83-113.
16. Simmers AJ, Bex PJ, Smith FK, Wilkins AJ. Spatiotemporal visual function in tinted lens wearers. *Investigative Ophthalmology and Visual Sciences* 2001 March; 42(3): 879-84.
17. Scheiman M, Mitchell GL, Cotter S, Cooper J, Kulp M, Rouse M, Borsting E, London R, Wensveen J; Convergence Insufficiency Treatment Trial Study Group. A randomized clinical trial of treatments for convergence insufficiency in children. *Arch Ophthalmol.* 2005 Jan; 123(1): 14-24.
18. American Academy of Pediatrics. Learning Disabilities, Dyslexia, and Vision: A Subject Review. *Pediatrics* 1998; 102(5): 1217-1219. A statement of reaffirmation for this policy was published on August 1, 2008.
19. Convergence Insufficiency Treatment Trial Study Group. Randomized clinical trial of treatments for symptomatic convergence insufficiency in children. *Arch Ophthalmol.* 2008 Oct; 126(10): 1336-49.
20. Borsting E, Mitchell GL, Kulp MT, Scheiman M, et al. Improvement in academic behaviors after successful treatment of convergence insufficiency. *Optom Vis Sci.* 2012; 89(1): 12-18.
21. Henderson M. Questioning the benefits that coloured overlays can have for reading in students with and without dyslexia. *Journal of Research in Special Educational Needs.* 2012 June.
22. American Academy of Pediatrics, Learning Disabilities, Dyslexia and Vision: A Joint Technical Report. *Pediatrics* Vol. 127 No. 3 March 1, 2011.
23. American Optometric Association. Fact Sheets on Optometric Vision Therapy, Reimbursement Packet. January 2011.
24. National Guideline Clearing House. Care of the Patient with Accommodative and Vergence Dysfunction.

25. American Association for Pediatric and Ophthalmology and Strabismus: Convergence Insufficiency; Vision Therapy.
26. Borsting E, Mitchell GL, Arnold LE, et al. Behavioral and Emotional Problems Associated with Convergence Insufficiency in Children: An Open Trial. *J Atten Disord*. Nov 22 2013
27. Handler SM, Fierson WM, Section on Ophthalmology and Council on Children with Disabilities AAoO, American Association for Pediatric Ophthalmology and Strabismus, and American Association of Certified Orthoptists,. Learning disabilities, dyslexia, and vision. *Pediatrics*. Mar 2011;127(3):e818-856. PMID 21357342
28. Scheiman M, Gwiazda J, Li T. Non-surgical interventions for convergence insufficiency. *Cochrane Database Syst Rev*. 2011(3):CD006768. PMID 21412896
29. Convergence Insufficiency Treatment Trial Study Group. Randomized clinical trial of treatments for symptomatic convergence insufficiency in children. *Arch Ophthalmol*. Oct 2008;126(10):1336-1349. PMID 18852411
30. Convergence Insufficiency Treatment Trial Study Group. Long-term effectiveness of treatments for symptomatic convergence insufficiency in children. *Optom Vis Sci*. Sep 2009;86(9):1096-1103. PMID 19668097
31. Scheiman M, Cotter S, Kulp MT, et al. Treatment of accommodative dysfunction in children: results from a randomized clinical trial. *Optom Vis Sci*. Nov 2011;88(11):1343-1352. PMID 21873922
32. Borsting E, Mitchell GL, Kulp MT, et al. Improvement in Academic Behaviors After Successful Treatment of Convergence Insufficiency. *Optom Vis Sci*. Nov 10 2012. Jan;89(1):12-8
33. Barnhardt C, Cotter SA, Mitchell GL, et al. Symptoms in children with convergence insufficiency: before and after treatment. *Optom Vis Sci*. Oct 2012;89(10):1512-1520. PMID 22922781
34. Scheiman M, Rouse M, Kulp MT, et al. Treatment of convergence insufficiency in childhood: a current perspective. *Optom Vis Sci*. May 2009;86(5):420-428. PMID 19319008
35. Shin HS, Park SC, Maples WC. Effectiveness of vision therapy for convergence dysfunctions and long-term stability after vision therapy. *Ophthalmic Physiol Opt*. Mar 2011;31(2):180-189. PMID 21309805
36. Dusek WA, Pierscionek BK, McClelland JF. An evaluation of clinical treatment of convergence insufficiency for children with reading difficulties. *BMC Ophthalmol*. 2011;11:21. PMID 21835034
37. Borsting E, Mitchell GL, Arnold LE, et al. Behavioral and Emotional Problems Associated With Convergence Insufficiency in Children: An Open Trial. *J Atten Disord*. Nov 22 2013. PMID 24271946
38. Lee SH, Moon BY, Cho HG. Improvement of Vergence Movements by Vision Therapy Decreases K-ARS Scores of Symptomatic ADHD Children. *J Phys Ther Sci*. Feb 2014;26(2):223-227. PMID 24648636
39. Ramsay MW, Davidson C, Ljungblad M, et al. Can vergence training improve reading in dyslexics? *Strabismus*. Dec 2014;22(4):147-151. PMID 25333204
40. United States Preventative Services Task Force (USPSTF) Visual Impairment in Children Ages 1-5, January 2011.
41. American Academy of Ophthalmology. Convergence Insufficiency. EyeWiki.
42. UpToDate. Causes of Horizontal Strabismus in Children. Topic last updated January 2023. Also available at <https://www.uptodate.com>
43. UpToDate. Sequelae of Mild Traumatic Brain Injury. Topic last updated September 2024. Also available at <https://www.uptodate.com>
44. Lavrich JB. Convergence Insufficiency and its current treatment. *Curr Opin Ophthalmol* 2010 Sep;21(5):356-60
45. Lambert J. Vision therapy and computer orthoptics: evidence-based approach to use in your practice. *Am Orthopt J* 2013;63:32-35
46. Whitecross S. Vision therapy: are you kidding me? Problems with current studies. *Am Orthopt J* 2013;63:36-40

47. Dusek WA, Pierscionek BK, McClelland JF. An evaluation of clinical treatment of convergence insufficiency for children with reading difficulties. *BMC Ophthalmol.* 2011;11:12
48. National Eye Institute. More Effective Treatment Identified for Common Childhood Vision Disorder.
49. Cooper J, Burns C, Cotter S, et. al. Optometric Clinical Practice Guideline Care of the Patient with Accommodative and Vergence Dysfunction, revised 2010.
50. United States Preventative Services Task Force (USPSTF), Visual Impairment in Children Ages 1-5: Screening, issued January 2011.
51. Hayes, a symplr company. Technology Brief. Vision Therapy for Convergence Insufficiency and Accommodative Dysfunction in Children, December 17, 2013.
52. Ramsay MW, Davidson C, Ljungblad M, et. al. Can vergence training improve reading in dyslexics? *Strabismus* Dec 2014;2294):147-151. PMID 25333204
53. Borsting E, Mitchell GL, Arnold LE, et. al. Behavioral and emotional problems associated with convergence insufficiency in children: an open trial. *J Atten Disord* Oct 2016;20910):836-844. PMID 24271946
54. Alvarez TL, Scheiman M, Santos EM, et al. Convergence Insufficiency Neuro-mechanism in Adult Population Study Randomized Clinical Trial: Clinical Outcome Results. *Optom Vis Sci.* Dec 2020; 97(12): 1061-1069. PMID 33186192
55. Singh A, Saxena V, Yadav S, et al. Comparison of home-based pencil push-up therapy and office-based orthoptic therapy in symptomatic patients of convergence insufficiency: a randomized controlled trial. *Int Ophthalmol.* Jan 04 2021. PMID 33392946
56. CITT-ART Investigator Group. Effect of Vergence/Accommodative Therapy on Reading in Children with Convergence Insufficiency: A Randomized Clinical Trial. *Optom Vis Sci.* Nov 2019; 96(11): 836-849. PMID 31651592
57. Chen AM, Roberts TL, Cotter SA, et al. Effectiveness of vergence/accommodative therapy for accommodative dysfunction in children with convergence insufficiency. *Ophthalmic Physiol Opt.* Jan 2021; 41(1): 21-32. PMID 33119180
58. CITT-ART Investigator Group. Treatment of Symptomatic Convergence Insufficiency in Children Enrolled in the Convergence Insufficiency Treatment Trial-Attention Reading Trial: A Randomized Clinical Trial. *Optom Vis Sci.* Nov 2019; 96(11): 825-835. PMID 31651593
59. Scheiman M, Kulp MT, Cotter SA, et al. Interventions for convergence insufficiency: a network meta-analysis. *Cochrane Database Syst Rev.* Dec 02 2020; 12: CD006768. PMID 33263359
60. Hayes, a symplr company. Evidence Analysis Research Brief. Vision Therapy for Treatment for Convergence Insufficiency in Adults. August 16, 2023
61. Hayes, a symplr company. Evolving Evidence Review. RevitalVision Perceptual Training Program (Taishir Medical Technologies LTD) for Treatment of Amblyopia. May 8, 2024
62. Jang JU, Jang JY, Tai-Hyung K. et. al. Effectiveness of vision therapy in school children with symptomatic convergence insufficiency. *J Ophthalmic Vis Res.* 2017 Apr-Jun;12(2):187-192. PMID 28540011

CODES

To report provider services, use appropriate CPT codes, HCPCS codes, Revenue codes, and/or ICD diagnosis codes.

Codes	Number	Description
CPT		
	92065	Orthoptic training; performed by a physician or other qualified health care professional
	92066	Orthoptic training; under supervision of a physician or other qualified health care professional
HCPCS		
	None	
Type of Service	Vision	
Place of Service	Physician's Office	

POLICY HISTORY

Date	Action	Action
June 2025	Annual Review	Policy Renewed
May 2024	Annual Review	Policy Renewed
May 2023	Annual Review	Policy Revised
March 2022	Annual Review	Policy Renewed
March 2021	Annual Review	Policy Renewed
March 2020	Annual Review	Policy Renewed
March 2019	Annual Review	Policy Renewed
March 2018	Annual Review	Policy Revised
March 2017	Annual Review	Policy Revised
October 2016	Interim Review	Policy Revised
March 2016	Annual Review	Policy Revised
March 2015	Annual Review	Policy Renewed
April 2014	Annual Review	Policy Revised
June 2013	Annual Review	Policy Renewed

Date	Action	Action
July 2012	Annual Review	Policy Renewed
August 2011	Annual Review	Policy Renewed

New information or technology that would be relevant for Wellmark to consider when this policy is next reviewed may be submitted to:

Wellmark Blue Cross and Blue Shield
 Medical Policy Analyst
 PO Box 9232
 Des Moines, IA 50306-9232

*CPT® is a registered trademark of the American Medical Association.