



Wellmark Blue Cross and Blue Shield is an Independent Licensee of the Blue Cross and Blue Shield Association.

DRUG POLICY

Neuromuscular Blocking Agents

This Prior Authorization request will be reviewed for medical necessity only. Benefits are subject to the terms and conditions of the patient's contract. Please contact Wellmark customer service at the number on the patient's card with benefit questions.

NOTICE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

BENEFIT APPLICATION

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

DESCRIPTION

The intent of the Neuromuscular Blocking Agent policy is to ensure appropriate selection of patients for therapy based on product labeling, clinical guidelines and clinical studies while steering utilization to the most cost-effective medication within the therapeutic class. For this program, Botox (onabotulinumtoxinA), Daxxify (daxibotulinumtoxinA), and Xeomin (incobotulinumtoxinA) are the preferred products. The criteria will require the use of the health plan's preferred products before the use of targeted products, Myobloc (rimabotulinumtoxinB) and Dysport (abobotulinumtoxinA), for an indication that is FDA-approved for the preferred product unless there are clinical circumstances that exclude the use of the preferred products and may be based on previous use of a product. The coverage review process will ascertain situations where a clinical exception can be made.

Botulinum toxin is a protein produced by the bacterium *Clostridium Botulinum*. There are seven distinct serotypes designated as type, A, B, C-1, D, E, F and G. Only Type A and Type B preparations are currently available in the United States. When administered intramuscularly, all botulinum toxins reduce muscle tone by interfering with the release of acetylcholine from nerve endings.

U.S. Food and Drug Administration (FDA)-approved labeled indications are few, but botulinum toxin has been used for a wide variety of off-label indications.

Exclusions

Coverage will not be provided for cosmetic use.

Botox (onabotulinum toxin A)

FDA Approved Indications

- Treatment of overactive bladder with symptoms of urge urinary incontinence, urgency, and frequency, in adults who have an inadequate response to or are intolerant of an anticholinergic medication
- Treatment of urinary incontinence due to detrusor overactivity associated with a neurologic condition (e.g., spinal cord injury, multiple sclerosis) in adults or pediatric patients 5 years of age or older who have an inadequate response to or are intolerant of an anticholinergic medication
- Prophylaxis of headaches in adult patients with chronic migraine (≥ 15 days per month with headache lasting 4 hours a day or longer)
- Treatment of spasticity in patients 2 years of age and older
- Treatment of cervical dystonia in adults, to reduce the severity of abnormal head position and neck pain
- Treatment of severe primary axillary hyperhidrosis that is inadequately managed with topical agents. Safety and effectiveness have not been established in patients under age 18.
- Treatment of strabismus and blepharospasm associated with dystonia, including benign essential blepharospasm or VII nerve disorders in patients 12 years of age and older

Compendial Uses

- Achalasia
- Chronic anal fissures
- Essential tremor
- Excessive salivation (ptyalism)
- Hemifacial spasm
- Spasmodic dysphonia (laryngeal dystonia)
- Oromandibular dystonia
- Myofascial pain syndrome
- Focal hand dystonia
- Facial myokymia
- Hirschsprung disease with internal sphincter achalasia
- Orofacial tardive dyskinesia
- Painful bruxism
- Palatal myoclonus
- First bite syndrome
- Palmar or gustatory (Frey's syndrome) hyperhidrosis

Myobloc (rimabotulinumtoxin B)

FDA Approved Indications

- Treatment of cervical dystonia in adults to reduce the severity of abnormal head position and neck pain associated with cervical dystonia
- Treatment of chronic sialorrhea in adults

Compendial Uses

- Primary axillary and palmar hyperhidrosis
- Upper limb spasticity

Dysport (abobotulinumtoxin A)

FDA Approved Indications

- Treatment of cervical dystonia in adults

- Treatment of spasticity in patients 2 years of age and older

Compensatory Uses

- Blepharospasm
- Hemifacial spasm
- Chronic anal fissures
- Excessive salivation
- Primary axillary hyperhidrosis

Xeomin (incobotulinumtoxin A)

FDA Approved Indications

- Treatment of chronic sialorrhea in patients 2 years of age and older
- Treatment of upper limb spasticity in adult patients
- Treatment of upper limb spasticity in pediatric patients 2 to 17 years of age, excluding spasticity caused by cerebral palsy
- Treatment of cervical dystonia in adult patients
- Treatment of blepharospasm in adult patients

Daxxify (daxibotulinumtoxin A)

FDA Approved Indications

- Treatment of cervical dystonia in adult patients

Preferred Drug Plan Design

Must meet BOTH the Preferred Drug Plan Design and Criteria for Initial Approval/Continuation of Therapy when both are applicable.

Criteria for initial approval for Myobloc and Dysport will only apply when one of the following criteria are met:

1. The patient has had an inadequate response to treatment, intolerable adverse event, or has a contraindication to therapy with ALL of the preferred products, Botox, Daxxify, and Xeomin, for an indication that is FDA-approved for the preferred product(s).
2. Member is currently receiving treatment with the requested product through insurance (excludes obtainment as samples or via manufacturer's patient assistance programs) and experiencing a positive therapeutic outcome.

Table. Botulinum Toxins

Medication	Generic Name
Preferred Products:	
Botox	onabotulinumtoxinA
Daxxify	daxibotulinumtoxinA
Xeomin	incobotulinumtoxinA
Targeted Product(s):	
Myobloc	rimabotulinumtoxinB
Dysport	abobotulinumtoxinA

POLICY

Prescriber Specialty

The medication must be prescribed by, or in consultation with one of the following:

- Blepharospasm, Strabismus: neurologist or ophthalmologist
- Overactive bladder, urinary incontinence: neurologist, urologist, or gynecologist
- Spasticity, cervical dystonia, hemifacial spasm, myofascial pain syndrome, upper limb spasticity focal hand dystonia, facial myokymia: neurologist, orthopedist, otolaryngologist or physiatrist
- Hyperhidrosis: neurologist or dermatologist
- Tremor, orofacial tardive dyskinesia: neurologist or physiatrist
- Achalasia, Hirschsprung disease: gastroenterologist, proctologist, or colorectal surgeon
- Excessive salivation, spasmodic dystonia, oromandibular dystonia, bruxism, palatal myoclonus: neurologist or otolaryngologist
- First bite syndrome: neurologist or oncologist
- Chronic sialorrhea: neurologist or otolaryngologist
- Primary axillary and palmar hyperhidrosis: neurologist or dermatologist
- Chronic anal fissures: gastroenterologist, proctologist, colorectal surgeon, or general surgeon with experience treating anal fissures

Criteria for Initial Approval

Botox

A. Blepharospasm

Authorization of 12 months may be granted for treatment of blepharospasm when all of the following are met:

1. Member is 12 years of age or older
2. Member is diagnosed with blepharospasm including blepharospasm associated with dystonia, benign essential blepharospasm or VII nerve disorder.

B. Cervical dystonia

Authorization of 12 months may be granted for the treatment of adults with cervical dystonia (e.g., torticollis) when all of the following are met:

1. There is abnormal placement of the head with limited range of motion in the neck
2. Member is 18 years of age or older.

C. Chronic migraine prophylaxis

Authorization of 6 months (two injection cycles) may be granted for treatment of chronic migraine prophylaxis when all of the following criteria are met:

1. Member experiences headaches 15 days or more per month.
2. Member experiences headaches lasting 4 hours or longer on at least 8 days per month.
3. Member completed an adequate trial of (or has a contraindication to) two oral migraine preventative therapies coming from at least 2 of the following classes with a trial of each medication at least 60 days in duration:
 - a. Antidepressants (e.g., amitriptyline, venlafaxine)
 - b. Antiepileptic drugs (AEDs) (e.g., divalproex sodium, topiramate, valproate sodium)
 - c. Beta-adrenergic blocking agents (e.g., metoprolol, propranolol, timolol, atenolol, nadolol)
4. Member has signs and symptoms consistent with chronic migraine diagnostic criteria as defined by the International Headache Society (IHS).
5. Member is 18 years of age or older

D. Overactive bladder with urinary incontinence

Authorization of 12 months may be granted for treatment of overactive bladder with urinary incontinence, urgency, and frequency when all of the following criteria are met:

1. The member has tried and failed behavioral therapy.
2. The member has had an inadequate response or experienced intolerance to two agents from either of the following classes:
 - a. Anticholinergic medication (e.g., Vesicare [solifenacin], Enablex [darifenacin], Toviaz [fesoterodine], Detrol/Detrol LA [tolterodine], Sanctura/Sanctura XR [trospium], Ditropan XL [oxybutynin]).
 - b. Beta-3 adrenergic agonist (e.g., Myrbetriq [miraberon], Gemtesa [vibegron]).
3. Member is 18 years of age or older.

E. Primary axillary, palmar, and gustatory (Frey's syndrome) hyperhidrosis

Authorization of 12 months may be granted for treatment of primary axillary, palmar, or gustatory (Frey's syndrome) hyperhidrosis when all of the following criteria are met:

1. Significant disruption of professional and/or social life has occurred because of excessive sweating; and
2. Topical aluminum chloride or other extra-strength antiperspirants are ineffective or result in a severe rash.
3. Member is 18 years of age or older.

F. Strabismus

Authorization of 12 months may be granted for treatment of strabismus when all of the following are met:

1. Strabismus interference with normal visual system development is likely to occur and spontaneous recovery is unlikely.
2. Member is 12 years of age or older.

Note: Strabismus repair is considered cosmetic in adults with uncorrected congenital strabismus and no binocular fusion.

G. Upper or lower limb spasticity

Authorization of 12 months may be granted for treatment of upper or lower limb spasticity when all of the following are met:

1. Member is 2 years of age or older
2. Member has a primary diagnosis of upper or lower limb spasticity or as a symptom of a condition causing limb spasticity (including focal spasticity or equinus gait due to cerebral palsy).

H. Urinary incontinence associated with a neurologic condition (e.g., spinal cord injury, multiple sclerosis)

Authorization of 12 months may be granted for treatment of urinary incontinence associated with a neurologic condition (e.g., spinal cord injury, multiple sclerosis) when all of the following criteria are met:

1. The member has tried and failed behavioral therapy
2. The member has had an inadequate response or experienced intolerance to one agent from either of the following classes:
 - a. Anticholinergic medication (e.g., Vesicare [solifenacin], Enablex [darifenacin], Toviaz [fesoterodine], Detrol/Detrol LA [tolterodine], Sanctura/Sanctura XR [trospium], Ditropan XL [oxybutynin]).
 - b. Beta-3 adrenergic agonist (e.g., Myrbetriq [miraberon])
3. Member is 5 years of age or older.

I. Achalasia

Authorization of 12 months may be granted for treatment of achalasia when the member has tried and failed or is a poor candidate for conventional therapy such as pneumatic dilation and surgical myotomy.

J. Chronic anal fissures

Authorization of 12 months may be granted for treatment of chronic anal fissures when the member has not responded to first line therapy such as topical calcium channel blockers or topical nitrates.

K. Essential tremor

Authorization of 12 months may be granted for treatment of essential tremor.

L. Excessive salivation

Authorization of 12 months may be granted for treatment of excessive salivation (chronic sialorrhea or ptyalism) when the member has been refractory to pharmacotherapy (e.g., anticholinergics).

M. Hemifacial Spasm

Authorization of 12 months may be granted for treatment of hemifacial spasm.

N. Spasmodic dysphonia (laryngeal dystonia)

Authorization of 12 months may be granted for treatment of spasmodic dysphonia (laryngeal dystonia).

O. Oromandibular dystonia

Authorization of 12 months may be granted for treatment of oromandibular dystonia.

P. Myofascial Pain Syndrome

Authorization of 12 months may be granted for treatment of myofascial pain syndrome when the member has tried and failed all of the following:

1. Physical therapy
2. Injection of local anesthetics into trigger points
3. Injection of corticosteroids into trigger points

Q. Focal hand dystonia

Authorization of 12 months may be granted for the treatment of focal hand dystonias.

R. Facial myokymia

Authorization of 12 months may be granted for the treatment of facial myokymia.

S. Hirschsprung disease with internal sphincter achalasia

Authorization of 12 months may be granted for the treatment of Hirschsprung's disease with internal sphincter achalasia following endorectal pull through and the member is refractory to laxative therapy.

T. Orofacial tardive dyskinesia

Authorization of 12 months may be granted for the treatment of orofacial tardive dyskinesia when conventional therapies have been tried and failed (e.g., benzodiazepines, clozapine, or tetrabenazine).

U. Painful bruxism

Authorization of 12 months may be granted for the treatment of painful bruxism when the member has had an inadequate response to a night guard and has had an inadequate response to pharmacologic therapy such as diazepam.

V. Palatal myoclonus

Authorization of 12 months may be granted for the treatment of palatal myoclonus when the member has disabling symptoms (e.g., intrusive clicking tinnitus) who had an inadequate response to clonazepam, lamotrigine, carbamazepine or valproate.

W. First bite syndrome

Authorization of 12 months may be granted for the treatment of first bite syndrome when the member has failed relief from analgesics, antidepressants or anticonvulsants.

Myobloc**A. Cervical dystonia**

Authorization of 12 months may be granted for treatment of adults with cervical dystonia (e.g., torticollis) when all of the following are met:

1. Member is 18 years of age or older
2. Member has abnormal placement of the head with limited range of motion in the neck

B. Chronic Sialorrhea (excessive salivation)

Authorization of 12 months may be granted for treatment of excessive salivation (chronic sialorrhea) when all of the following are met:

1. Member is 18 years of age or older
2. Member is refractory to pharmacotherapy (e.g., anticholinergics)

C. Primary axillary and palmar hyperhidrosis

Authorization of 12 months may be granted for treatment of primary axillary or palmar hyperhidrosis when all of the following criteria are met:

1. Significant disruption of professional and/or social life has occurred because of excessive sweating; and
2. Topical aluminum chloride or other extra-strength antiperspirants are ineffective or result in a severe rash.

D. Upper limb spasticity

Authorization of 12 months may be granted for treatment of upper limb spasticity either as a primary diagnosis or as a symptom of a condition causing limb spasticity.

Dysport**A. Cervical dystonia**

Authorization of 12 months may be granted for treatment of adults with cervical dystonia (e.g., torticollis) when all of the following are met:

1. Member is 18 years of age or older
2. Member has abnormal placement of the head with limited range of motion in the neck.

B. Upper or lower limb spasticity

Authorization of 12 months may be granted for treatment of upper or lower limb spasticity when all of the following are met:

1. Member is 2 years of age or older
2. Member has a primary diagnosis of upper or lower limb spasticity or as a symptom of a condition (including focal spasticity or equinus gait due to cerebral palsy)

C. Blepharospasm

Authorization of 12 months may be granted for treatment of blepharospasm, including blepharospasm associated with dystonia and benign essential blepharospasm.

D. Hemifacial spasm

Authorization of 12 months may be granted for treatment of hemifacial spasm.

E. Chronic anal fissures

Authorization of 12 months may be granted for treatment of chronic anal fissures when the member has not responded to first-line therapy such as topical calcium channel blockers or topical nitrates.

F. Excessive salivation

Authorization of 12 months may be granted for treatment of excessive salivation (chronic sialorrhea) when the member has been refractory to pharmacotherapy (e.g., anticholinergics).

G. Primary axillary hyperhidrosis

Authorization of 12 months may be granted for treatment of primary axillary hyperhidrosis when all of the following criteria are met:

1. Significant disruption of professional and/or social life has occurred because of excessive sweating; and
2. Topical aluminum chloride or other extra-strength antiperspirants are ineffective or result in a severe rash.

Xeomin

A. Chronic Sialorrhea (Excessive Salivation)

Authorization of 12 months may be granted for treatment of chronic sialorrhea (excessive salivation) when all of the following criteria are met:

1. Member is 2 years of age or older
2. Member has been refractory to pharmacotherapy (e.g., anticholinergics)

B. Cervical dystonia

Authorization of 12 months may be granted for treatment of adults with cervical dystonia (e.g., torticollis) when all of the following criteria are met:

1. Member is 18 years of age or older
2. There is abnormal placement of the head with limited range of motion in the neck.

C. Blepharospasm

Authorization of 12 months may be granted for treatment of blepharospasm when all of the following criteria are met:

1. Member has a diagnosis of blepharospasm including benign essential blepharospasm or blepharospasm associated with dystonia
2. Member is 18 years of age or older.

D. Upper limb spasticity

Authorization of 12 months may be granted for the treatment of upper limb spasticity when all of the following are met:

1. Member has a diagnosis of upper limb spasticity either as a primary diagnosis or as a symptom of a condition causing limb spasticity
2. Member meets one of the following criteria:

- a. Member is 18 years of age or older
- b. Member is 2 to 17 years of age and the spasticity is not caused by cerebral palsy.

Daxxify

A. Cervical dystonia

Authorization of 12 months may be granted for the treatment of adults with cervical dystonia (e.g., torticollis) when all of the following criteria are met:

1. Member is 18 years of age or older.
2. Member has abnormal placement of the head with limited range of motion in the neck.

Continuation of Therapy

- A.** All members (including new members) requesting authorization for continuation of therapy for all approvable conditions other than chronic migraine prophylaxis must meet ALL initial authorization criteria and be experiencing benefit from therapy.
- B.** Authorization of 12 months may be granted for treatment of chronic migraine prophylaxis when the member has achieved or maintained a reduction in monthly headache frequency since starting therapy with Botox.

Botox, Daxxify, Dysport, Myobloc, and Xeomin are considered **not medically necessary** for members who do not meet the criteria set forth above.

Dosage and Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

Quantity Limits

Botox 400 units; Myobloc 10,000 units; Dysport 1,500 units; Xeomin 400 units; Daxxify 1,000 units

PROCEDURES AND BILLING CODES

To report provider services, use appropriate CPT* codes, Alpha Numeric (HCPCS level 2) codes, Revenue codes, and/or ICD diagnostic codes.

- J0585 Injection, onabotulinumtoxinA (Botox), 1 unit
- J0586 Injection, abobotulinumtoxinA (Dysport), 5 units
- J0587 Injection, rimabotulinumtoxinB (Myobloc), 100 units
- J0588 Injection, incobotulinumtoxinA (Xeomin), 1 unit
- J0589 Injection, daxibotulinumtoxinA-lanm (Daxxify), 1 unit (effective 4/1/2024)
- C9160 Injection, daxibotulinumtoxinA-lanm (Daxxify), 1 unit (cancelled 3/31/2024)

REFERENCES

- Botox [package insert]. Madison, NJ: Allergan USA, Inc; August 2023.
- Dysport [package insert]. Wrexham, UK; Ipsen Biopharm, Ltd ; September 2023.
- Myobloc [package insert]. South San Francisco, CA: Solstice Neurosciences, Inc; March 2021.
- Xeomin [package insert]. Raleigh, NC: Merz Pharmaceuticals LLC; September 2023.
- Daxxify [package insert]. Newark, CA: Revance Therapeutics, Inc; August 2023.
- DRUGDEX® System (electronic version). Truven Health Analytics, Ann Arbor, MI. Available at <http://www.micromedexsolutions.com>. Accessed August 1, 2022.
- AHFS Drug Information. <http://online.lexi.com/lco>. Accessed August 1, 2022.

- Silberstein SD, Holland S, Freitag F, et al. Evidence-based guideline update: pharmacologic treatment for episodic migraine prevention in adults. Report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Headache Society. *Neurology*. 2012;78(17):1337-45.
- Snow V, Weiss K, Wall EM, et al. Pharmacologic management of acute attacks of migraine and prevention of migraine headache. *Ann Intern Med*. 2002;137(10):840-849.
- Evers S, Afra J, Frese A, et al. EFNS guideline on the drug treatment of migraine--revised report of an EFNS task force. *Eur J Neurol*. 2009;16(9):968-981.
- Pringsheim T, Davenport W, Mackie G, et al. Canadian Headache Society guideline for migraine prophylaxis. *Can J Neurol Sci*. 2012;39(2 Suppl 2):S1-59.
- Clinical Consult: CVS Caremark Clinical Programs Review. Focus on Internal Medicine Programs; November 2015.
- Zesiewicz TA, Elble RJ, Louis ED et al. Evidence-based guideline update: Treatment of essential tremor: Report of the Quality Standards Subcommittee of the American Academy of Neurology: *Neurology* 2011; 77: 1752-1755.
- Restivo D, Panebianco M, Casabona A et al. Botulinum Toxin A for Sialorrhea Associated with Neurological Disorders: Evaluation of the Relationship between Effect of Treatment and the Number of Glands Treated. *Toxins* 2018;55:1-10.
- Lakraj AA, Moghimi N, Jabbari B. Sialorrhea: Anatomy, Pathophysiology and Treatment with
- Emphasis on the Role of Botulinum Toxins. *Toxins* 2013, 5, 1010-1031
- Zoons E, Dijkgraaf M, Dijk J et al. Botulinum toxin as treatment for focal dystonia: a systematic review of the pharmaco-therapeutic and pharmaco-economic value. *J Neurol* 2012;259: 2519-2526.
- Conrin L, Karp BI, Alter K et al. Long Term Follow-up Botulinum Toxin Therapy for Focal Hand Dystonia: Outcome at 10 or More Years: *Mov Disord*. 2011 Mar; 26(4): 750–753.
- Daele D, Finnegan E, Rodnitzky R et al. Head and Neck Muscle Spasm After Radiotherapy. *Otolaryngol Head and Neck Surg* 2002;128:956-959.
- Han-Guerts I, Hendrix V, Blaauw I et al. Outcome after Anal Intraspinal Botox Injection in Children with Surgically Treated Hirschsprung Disease. *JPGN* 2014; 59: 604-607.
- Asutay F, Atalay Y, Asutay H, Huseyin Acar A. The Evaluation of the clinical effects of Botulinum Toxin on Nocturnal Bruxism. *Pain Research and Management* 2017;1-5.
- Gosain A, Frykman PK, Cowles RA, et al. Guidelines for the diagnosis and management of Hirschsprung-associated enterocolitis. *Pediatr Surg Int*. 2017; 33(5):517–521.
- Thuruthiath N, Arayamparambil R. Essential palatal myoclonus: A rare cause of objective tinnitus. *J Adv Med Health Res* 2016;3:1-3.
- Laccourreye O, Werner A, Garcia D, Malinvaud D, Tran Ba Huy P, Bonfils P. First bite syndrome. *Euro Ann Otolaryngol Head Neck Diseases*. 2013; 130:269-273.
- Linkov G et al. First bite syndrome: incidence, risk factors, treatment, and outcomes. *The Laryngoscope*. 2012; 122: 1773-1778.
- Slotema CW;van Harten PN;Bruggeman R;Hoek HW. Botulinum toxin in the treatment of orofacial tardive dyskinesia: a single blind study. *Prog Neuropsychopharmacol Biol Psychiatry* 2008;32(2):507-509.
- Odderson IR. Hyperhidrosis treated by botulinum A exotoxin. *Dermatol Surg*. 1998;24(11):1237-1241.
- Solomon BA, Hayman R. Botulinum toxin type A therapy for palmar and digital hyperhidrosis. *J Am Acad Dermatol*. 2000;42(6):1026-1029.
- Naver H, Swartling C, Aquilonius SM. Palmar and axillary hyperhidrosis treated with botulinum toxin: One-year clinical follow-up. *Eur J Neurol*. 2000;7(1):55-62.
- Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders: 3rd edition. *Cephalalgia*. 2018;Vol. 38(1) 1–211.

- Glader L, Delsing C, Hughes A et al. Sialorrhea in cerebral palsy. American Academy for Cerebral Palsy and Developmental Medicine Care Pathways. <https://www.aacpdm.org/publications/care-pathways/sialorrhea>. Accessed August 3, 2021.
- Garuti G, Rao F, Ribuffo V et al. Sialorrhea in patients with ALS: current treatment options. *Degener Neurol Neuromuscul Dis*. 2019; 9: 19–26.
- Digre KB for the American Headache Society. The American Headache Society Position Statement On Integrating New Migraine Treatments Into Clinical Practice. *Headache* 2019;59:1-18. doi: 10.1111/head.13456.
- Lightner DJ, Gomelsky A, Souter L, Vasavada SP. Diagnosis and Treatment of Overactive Bladder (Non-Neurogenic) in Adults: AUA/SUFU Guideline Amendment 2019. *J Urol* 2019; 202:558.
- Smith C, Pariser D. Primary Focal Hyperhidrosis. Waltham, MA. UpToDate. Last Modified January 2, 2020. <https://www.uptodate.com/contents/primary-focal-hyperhidrosis>. Accessed August 15, 2022
- Lexi-Drugs. Hudson, OH: Lexicomp, 2019. <http://online.lexi.com/>. Accessed July 28, 2022.
- Simpson DM, Hallett M, Ashman EJ et al. Practice guideline update summary: Botulinum neurotoxin for the treatment of blepharospasm, cervical dystonia, adult spasticity, and headache: Report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology* 2016; 86 (19) 1818-1826.
- Dashtipour K, Chen JJ, Frei K, et al. Systemic Literature Review of AbobotulinumtoxinA in Clinical Trials for Blepharospasm and Hemifacial Spasm. *Tremor Other Hyperkinet Mov (NY)*. 2015;5:338.
- Clinical Pharmacology. Tampa (FL): Elsevier. 2019. Available from: <http://www.clinicalpharmacology-ip.com>. Accessed August 2, 2022.

POLICY HISTORY

Policy #: 05.01.02

Policy Creation: May 1991

Reviewed: June 2026

Revised: January 2026

Current Effective Date: March 28, 2026