

02.01.71 Biofeedback as a Treatment of Chronic Pain

Original Effective Date: July 2001

Review Date: December 2025

Revised: December 2025

DISCLAIMER/INSTRUCTIONS FOR USE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations, or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

This Medical Policy document describes the status of medical technology at the time the document was developed. Since that time, new technology may have emerged, or new medical literature may have been published. This Medical Policy will be reviewed regularly and updated as scientific and medical literature becomes available; therefore, policies are subject to change without notice.

Related Policies:

- [02.01.65 Neurofeedback](#)
- [02.01.70 Biofeedback as a Treatment of Anorectal Disorders](#)
- [02.01.72 Biofeedback as a Treatment of Headache](#)
- [02.01.73 Biofeedback as a Treatment of Urinary Incontinence](#)
- [02.01.74 Biofeedback as a Treatment of Miscellaneous Indications](#)

Summary

Description

Biofeedback is a technique intended to teach patients self-regulation of certain physiologic processes not normally considered to be under voluntary control. Electromyography biofeedback has been evaluated as a method to reduce chronic or recurrent pain of musculoskeletal or psychosomatic origin.

Summary of Evidence

For individuals who have cancer pain who receive biofeedback (BFB) the evidence includes systematic reviews, randomized controlled trials (RCTs), observational and case studies. Relevant outcomes are symptoms, functional outcomes, quality of life (QOL), and medication use. Spada et al (2022) completed a scope review of the available literature regarding heart rate variability (HRV) BFB utilized to modulate autonomic balance, cancer symptom management and quality of life (QOL) in cancer survivors or terminal cancer patients with thyroid, lung, brain, colon, and hematologic cancers. This review of the available studies found HRV BFB to be beneficial in improving cancer pain rates, fatigue scores, sleep and a reduction in depression and anxiety, however, not all studies achieved improvement in QOL and distress. While the evidence continues to accumulate regarding biofeedback in the treatment of cancer pain, this complementary approach using BFB is safe, non-invasive and absent of side effects. The evidence is considered insufficient to determine that the technology results in an improvement in the net health outcome, however, based on the current NCCN guidelines biofeedback is recommended as an optional intervention to reduce pain, see [Practice Guideline and Position Statements](#). Therefore, biofeedback will be considered medically necessary for the management of cancer pain when criteria below are met, see [Policy](#).

For individuals who have chronic pain (including low back, knee, neck and shoulder, orofacial, and abdominal pain as well as fibromyalgia, osteoarthritis, systemic lupus erythematosus, and vulvar vestibulitis) who receive biofeedback, the evidence includes multiple RCTs for different pain syndromes. Relevant outcomes are symptoms, functional outcomes, QOL, and medication use. The results of these RCTs, some of which were sham controlled, did not consistently report a benefit for biofeedback. Some RCTs reported improved outcomes with biofeedback, but these improvements were often of uncertain clinical significance or were not durable. Many other RCTs have found that biofeedback did not provide a significantly greater benefit in outcomes when it was used instead of or in addition to other conservative interventions such as exercise. Overall, the available RCTs were limited by small sample sizes and high dropout rates. This evidence base does not permit conclusions about the specific effects of biofeedback beyond the nonspecific effects of sham interventions, nor does it permit conclusions about the contribution of biofeedback beyond that of other conservative treatments for pain. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have chronic pain (including low back, knee, neck and shoulder, orofacial and abdominal pain as well as fibromyalgia, osteoarthritis, systemic lupus erythematosus, and vulvar vestibulitis) who receive various biofeedback devices for home use, no studies were identified. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

OBJECTIVE

The objective of this evidence review is to determine whether the use of biofeedback improves the net health outcome in individuals with chronic pain.

PRIOR APPROVAL

Not applicable.

POLICY

Biofeedback therapy training sessions by qualified provider (see [Policy Guidelines](#)) may be considered **medically necessary** up to 20 sessions as part of the treatment plan for the management of cancer pain.

Biofeedback therapy training sessions when the above criteria is not met and treatment of all other chronic pain conditions, including but not limited to the following, is considered **investigational** because the evidence is insufficient to determine that the technology results in an improvement in the net health outcome:

- Abdominal pain
- Fibromyalgia pain
- Knee pain
- Low back pain
- Orofacial pain
- Osteoarthritis
- Shoulder pain
- Systemic lupus erythematosus
- Vulvar vestibulitis

Home biofeedback is considered **investigational** for all indications, including but not limited to the following devices because the evidence is insufficient to determine the effects of the technology on net health outcomes:

- FreeSpira
- Heart Rate Variability
- InTandem
- StressEraser

POLICY GUIDELINES

Guidance on biofeedback protocol:

Biofeedback may require up to 20 office-based sessions of 30 to 60 minutes each.

Covered Provider

Biofeedback supervised by a physician or licensed qualified non-physician practitioner (a licensed qualified non-physician may include the following: physical therapists, Nurse Practitioners, Physician Assistants, or Clinical Nurse Specialists practitioner).

Required Documentation

The medical record documentation must support the medical necessity of the services as directed in this policy. Documentation in the individual's medical record must include the following:

- There should be a written treatment plan which includes **ALL** the following information:

- The specific diagnosis/conditions to be treated; **and**
- Indicate the types of biofeedback training to be utilized; **and**
- The time frame and the frequency of treatment in which the goals and objectives will be achieved.

Note: *If the provider or licensed qualified non-physician practitioner (a licensed qualified non-physician practitioner may include the following: physical therapists, Nurse Practitioners, Physician Assistants, or Clinical Nurse Specialists practitioner) has determined that the individual does not appear to be benefiting from biofeedback training or moving forward toward individual treatment goals after 20 sessions the use of biofeedback should be re-evaluated and the documentation provided should include information to support the request for further biofeedback therapy training sessions or the provider should consider suggesting an alternative treatment plan.*

Coding

See the [Codes](#) table for details.

BACKGROUND

Biofeedback for Chronic Pain

Biofeedback is intended to teach individuals self-regulation of certain physiological processes not normally considered to be under voluntary control. The various forms of biofeedback mainly differ in the nature of the disease or disorder under treatment, the biologic variable that the subject attempts to control, and the information that is fed back to the subject.

Biofeedback techniques include peripheral skin temperature feedback, blood-volume-pulse feedback (vasoconstriction and dilation), vasoconstriction training (temporalis artery), and electromyographic biofeedback; they may be used alone or in conjunction with other therapies (eg, relaxation, behavioral management, medication). Biofeedback training is done either in individual or group sessions. A typical program consists of 10 to 20 training sessions of 30 minutes each.

Biofeedback has been proposed as a treatment for a variety of diseases and disorders, including anxiety, headaches, hypertension, movement disorders, incontinence, pain, asthma, Raynaud disease, and insomnia.

Treatment for chronic pain is often multimodal and typically includes psychological therapy. Psychological techniques vary but may include cognitive therapy, which teaches subjects the ability to cope with stressful stimuli by attempting to alter negative thought patterns and dysfunctional attitudes, and behavioral approaches to reduce muscle tension and break the pain cycle. Relaxation, using any of a variety of techniques including meditation or mental imagery, is considered a behavioral therapy that may be used alone or as a component of a cognitive-behavioral therapy program. Electromyography biofeedback has also been used for the treatment of chronic pain, on the assumption that the ability to reduce muscle tension will be improved through the feedback of data to the patient regarding the degree of muscle tension. While some consider electromyography biofeedback to be a method used to obtain relaxation, others consider biofeedback to be distinct from other relaxation techniques.

Regulatory Status

A variety of biofeedback devices have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. These devices are designated by the FDA as class II with special controls and are exempt from premarket notification requirements. The FDA defines a biofeedback device as “an instrument that provides a visual or auditory signal corresponding to the status of 1 or more of a patient’s physiological parameters (e.g., brain alpha wave activity, muscle activity, skin temperature) so that the patient can control voluntarily these physiological parameters.”

RATIONALE

The evidence review was created in July 2001 and has been updated regularly with searches of the PubMed database. The most recent literature update was performed through December 2025.

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Psychological treatments involve both nonspecific and specific therapeutic effects. Nonspecific effects (sometimes called placebo effects) occur as a result of contact with the therapist, positive expectations on the part of the patient and therapist, and other beneficial effects that occur as a result of the patient being in a therapeutic environment. Specific effects are those that occur only because of the active treatment, beyond any nonspecific effects that may be present. This literature review focuses on identifying evidence that the effects of biofeedback are distinct from nonspecific placebo effects. Because establishing an ideal placebo control is problematic with psychological treatments and because treatment of chronic pain is typically multimodal, isolating the specific contribution of biofeedback is challenging.

Biofeedback

Clinical Context and Therapy Purpose

The purpose of electromyography (EMG) biofeedback in individuals who have chronic pain is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with chronic pain, including cancer pain, low back, knee, neck and shoulder, orofacial, and abdominal pain as well as fibromyalgia, osteoarthritis, systemic lupus erythematosus, and vulvar vestibulitis.

Interventions

The therapy being considered is EMG biofeedback.

Comparators

The following therapies are currently being used to treat chronic pain: pharmacologic and nonpharmacologic therapy. For chronic pain management, a multimodal, multidisciplinary approach that is individualized to the patient is recommended. A multimodal approach to pain management consists of using treatments (i.e., nonpharmacologic and pharmacologic) from one or more clinical disciplines incorporated into an overall treatment plan. This allows for different avenues to address the pain condition, often enabling a synergistic approach that impacts various aspects of pain, including functionality. The efficacy of such a coordinated, integrated approach has been documented to reduce pain severity, improve mood and overall quality of life, and increase function.

Outcomes

The general outcomes of interest are reductions in symptoms and medication usage and improvements in functional outcomes.

The Initiative on Methods, Measurement, and Pain Assessment in Clinical Trials (IMMPACT) recommends that chronic pain trials should consider assessing outcomes representing 6 core domains: pain, physical functioning, emotional functioning, participant ratings of improvement and satisfaction with treatment, symptoms and adverse events, and participant disposition. Table 1 summarizes provisional benchmarks for interpreting changes in chronic pain clinical trial outcome measures per IMMPACT.

Table 1. Benchmarks for Interpreting Changes in Chronic Pain Outcome Measures

Outcome Domain and Measure	Type of Improvement	Change
Pain intensity 0 to 10 numeric rating scale	Minimally important Moderately important Substantial	10 to 20% decrease ≥30% decrease ≥50% decrease
Physical functioning Multidimensional Pain Inventory Interference Scale	Clinically important	≥0.6-point decrease
Brief Pain Inventory Interference Scale	Minimally important	1 point decrease
Emotional functioning Beck Depression Inventory Profile of Mood States	Clinically important	≥5 point decrease

Outcome Domain and Measure	Type of Improvement	Change
Total Mood Disturbance Specific Subscales	Clinically important Clinically important	≥10 to 15 point decrease ≥2 to 12 point change
Global Rating of Improvement Patient Global Impression of Change	Minimally important Moderately important Substantial	Minimally improved Much improved Very much improved

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Cancer Pain

Spada et al (2022) completed a scope review of the available literature regarding heart rate variability (HRV) biofeedback (BFB) utilized to modulate autonomic balance, cancer symptom management and quality of life (QOL) in cancer survivors or terminal cancer patients with thyroid, lung, brain, colon, and hematologic cancers. Nineteen studies were included in this review which consisted of systematic reviews, RCTs, observational and case studies with sample size of up to 272 cancer patients (adults, adolescent and child). The outcomes reported were pain, fatigue, sleep quality, depression, anxiety, HVR coherence, frequency HCR indexes, stress and QOL. This review found there was an overall positive effect observed regarding the following variables utilizing HRV BFB: cancer pain rates and fatigue scores lowered based on self-report scoring; increased sleep efficiency, sleep duration and a general reduction of sleep symptoms to include insomnia, and daytime impairments were improved; and a significant reduction in the levels of depression and anxiety compared to before and after HRV BFB training were reported. However, not all studies reached improvements such as QOL and distress. The HRV measurements and coherence scores (i.e., a measure of the degree of coherence in the heart rhythm pattern) were increased, following the BFB and respiration exercises in several studies, which mirrored a better autonomic balance compared to the individual’s pre-treatment condition. Limitations of this review included medium-to-high dropout rates and missing data. While the evidence collected may have found HRV BFB to be beneficial in the treatment of these cancer patients, further high-quality studies are needed.

General Chronic Pain

Systematic Reviews

Several meta-analyses have reviewed RCTs assessing psychological therapies for a variety of non-headache chronic pain conditions. Calderone et al (2025) conducted a systematic review of biofeedback in the setting of rehabilitation for chronic pain. A total of 25 studies were included in the review; 15 studies

were RCTs and 10 studies were nonrandomized. The overall risk of bias was moderate for almost all of the included studies. A meta-analysis was not performed, but the authors concluded that the data supports the efficacy of biofeedback as adjunctive therapy for chronic pain. A limitation of this analysis is that some studies used biofeedback techniques other than EMG biofeedback.

A Cochrane review by Williams et al (2020) focused on chronic pain in adults. Two RCTs were identified that compared behavioral therapy with an active control designed to change behavior (i.e., exercise or instruction). Three RCTs had sufficient follow-up to be included in a comparison of behavioral therapy and usual treatment. Reviewers found no evidence that behavioral therapy had any effect on pain compared to active control or usual treatment. Additionally, there was no evidence of a difference between behavioral therapy and active control or usual treatment in terms of disability at the end of treatment.

Another Cochrane review by Fisher et al (2018) focused on children and adolescents with chronic and recurrent pain. Although psychological therapies were found to improve pain, only 1 study evaluated biofeedback in non-headache pain. Biofeedback did not improve abdominal pain more than cognitive-behavioral therapy (CBT) in this trial; see the section on Abdominal Pain. Palermo et al (2010) published a meta-analysis of studies on psychological therapies for the management of chronic pain in children and adolescents. These authors did not identify any additional RCTs on biofeedback for managing non-headache pain.

Low Back Pain

Systematic Reviews

A Cochrane review by Henschke et al (2010) assessed behavioral treatments for chronic low back pain and conducted a meta-analysis of 3 small, randomized trials that compared EMG biofeedback with a waiting-list control group. In the pooled analysis, there were a total of 34 patients in the intervention group and 30 patients in the control group. The standardized mean difference (SMD) in short-term pain was -0.80 (95% confidence interval [CI], -1.32 to -0.28); this difference was statistically significant favoring the biofeedback group. Reviewers did not conduct meta-analyses of trials comparing biofeedback with sham biofeedback and therefore were unable to control for any nonspecific effects of treatment.

Randomized Controlled Trials

Yelden et al (2024) compared biofeedback to physiotherapist feedback in an RCT in 40 patients with chronic nonspecific low back pain. All patients received 12 sessions of the designated therapy (3 sessions weekly for 4 weeks) and a core stabilization activity program. The primary outcome, disability as measured by the Revised Oswestry Disability Index scale was not significantly different between groups at the end of treatment. Secondary measures of visual analogue scale pain scores, muscle activity, and quality of life were also not different between groups.

At least one RCT has compared biofeedback with a sham intervention for the treatment of low back pain. Kapitza et al (2010) compared the efficacy of respiratory biofeedback with sham biofeedback in 42 patients with low back pain. Both groups showed a reduction in pain levels on a 10-point visual analog scale (VAS) at the end of the intervention period and at 3-month follow-up. Between-group differences were not statistically significant. For example, 3 months after the intervention, mean change in pain with activity decreased by 1.12 points in the intervention group and 0.96 points in the sham control group ($p > .05$); mean change in pain at rest decreased by 0.79 points in the intervention group and 0.49 points in the control group ($p > .05$).

Lazaridou et al (2023) conducted a prospective, single-center RCT to assess the impact of surface EMG biofeedback versus continued care (no intervention) on chronic lower back pain in adults. Sixty-six patients were randomized 2:1 to receive EMG biofeedback or no additional intervention for 8 weeks and included in analysis. Compared to usual care, patients receiving EMG biofeedback reported lower pain intensity on the Brief Pain Inventory (BPI) questionnaire after 8 weeks (mean difference [MD], 0.9; 95% CI, -1.07 to -0.32; $p \leq .01$). Compared to baseline scores, individuals in the EMG biofeedback group demonstrated statistically significant reductions in pain interference (MD, 1.3; 95% CI, 0.42 to 2.1; $p \leq .01$), disability (MD, 4.32; 95% CI, 1.2 to 7.3; $p \leq .01$), and significant increases in low back pain thresholds (MD, 0.5; 95% CI, -0.87 to -0.05; $p \leq .01$). Significant changes were also observed in muscle tension for the lower back muscles in the EMG biofeedback group ($p < .001$).

Several trials with active comparison groups have not found that biofeedback is superior to alternative treatments. More recently, Tan et al (2015) evaluated 3 self-hypnosis interventions and included EMG biofeedback as a control intervention. This RCT enrolled 100 patients with chronic low back pain. After the 8-week intervention, reported reductions in pain intensity were significantly higher in the combined hypnosis groups than in the biofeedback group ($p = .042$).

A trial published by Glombiewski et al (2010) assessed whether the addition of EMG biofeedback to CBT improved outcomes in 128 patients with low back pain. Patients were randomized to one of 3 groups: CBT, CBT plus biofeedback, or waiting-list control. Both treatments improved outcomes including pain intensity compared with the waiting-list control (moderate effect size of 0.66 for pain intensity in the CBT plus biofeedback group). However, the addition of biofeedback did not improve outcomes over CBT alone.

Chronic Knee Pain

Systematic Reviews

Ananias et al (2024) conducted a systematic review and meta-analysis of 8 RCTs that compared the efficacy of biofeedback and standard rehabilitation in patients undergoing anterior cruciate ligament reconstruction surgery. Four of the RCTs were included in the meta-analysis. Two RCTs showed a significant effect of biofeedback on quadriceps strength, 2 studies reported a significant difference in pain scores, 2 studies found a significant difference in knee extension deficit, and one study reported a significant difference in balance. The heterogeneity of outcomes assessed limits the interpretation of these results in this subset of studies.

Karaborklu Argut et al (2022) conducted a systematic review of 8 RCTs of patients who had undergone orthopedic knee surgery. Therapeutic EMG biofeedback during rehabilitation was more effective for improving muscle strength and activation compared to home exercise, standard rehabilitation, or electrical stimulation. There were no clear trends in the effect of EMG biofeedback on pain or knee range of motion.

Collins et al (2012) conducted a systematic review and meta-analysis of RCTs on nonsurgical interventions for anterior knee pain. In a pooled analysis of data from 2 trials, there was no significant benefit of adding EMG biofeedback to an exercise-only intervention at 8 to 12 weeks (SMD, -0.22; 95% CI, -0.65 to 0.20).

Chronic Neck and Shoulder Pain

Systematic Reviews

Campo et al (2021) published a systematic review and meta-analysis that evaluated the effectiveness of biofeedback for improving pain, disability, and work ability in adults with neck pain. The review included 15 RCTs with 8 studies utilizing EMG biofeedback and 7 studies of pressure biofeedback (Table 2). There was no restriction on the control intervention (e.g., no treatment, placebo, active treatment) or co-intervention, provided the independent effects of biofeedback could be elucidated. An overview of the characteristics and results is presented in Tables 3 and 4. Results suggest that biofeedback has a moderate effect on reducing short-term disability and a small effect on reducing intermediate-term disability with no effect on pain or work ability in the short- and intermediate-term. Of note, there were a variety of control interventions across included studies (e.g., exercise, electroacupuncture, electrotherapy, education) with few studies directly comparing biofeedback to no treatment or placebo.

Kamonseki et al (2021) completed a systematic review and meta-analysis of 5 RCTs that examined the effects of EMG biofeedback for shoulder pain and function. Study characteristics and results are presented in Tables 3 and 4. Overall, the evidence did not support the use of EMG biofeedback for reducing shoulder pain and improving shoulder function.

Table 2. Comparison of Studies Included in Systematic Reviews and Meta-Analyses

Study	Campo et al (2021)	Kamonseki et al (2021)
Juul-Kristensen et al (2019)		●
Kosterink et al (2010)	●	●
Ma et al (2011)	●	●
Middaugh et al (2013)		●
Sandsjo et al (2010)	●	●
Arami et al (2012)	●	
Bissett et al (1985)	●	
Bobos et al (2016)	●	
Delive et al (2011)	●	
Ehrenborg et al (2010)	●	
Eslamian et al (2020)	●	
Iqbal et al (2013)	●	
Jull et al (2002)	●	
Jull et al (2007)	●	
Nezamuddin et al (2013)	●	
Voerman et al (2007)	●	
Wani et al (2013)	●	

Table 3. Systematic Reviews and Meta-Analysis Characteristics

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Campo et al (2021)	To Sept 2020	15	Adults with neck pain including pain associated with radiculopathy, cervicogenic headaches, whiplash, shoulder pain, and work-related injuries administered biofeedback (EMG or pressure) on at least 2 occasions	990 (27 to 200)	RCT (8 studies EMG; 7 pressure)	8 days to 6 weeks (duration of interventions)
Kamonseki et al (2021)	To Dec 2020	5	Adults with shoulder pain	272 (15 to 72)	RCT (all EMG)	4 weeks to 6 months (follow-up period)

EMG: electromyography; RCT: randomized controlled trial.

Table 4. Systematic Reviews and Meta-Analysis Results

Study	Pain (short-term: 4 to 6 weeks)	Pain (intermediate-term: 8 to 12 weeks)	Disability (short-term: 4 to 6 weeks)	Disability (intermediate-term: 8 to 12 weeks)	Work ability (short-term: 4 to 6 weeks)	Work ability (intermediate-term: 8 to 12 weeks)
Campo et al (2021)						
Total N	602 (11 RCTs)	383 (6 RCTs)	627 (9 RCTs)	458 (5 RCTs)	190 (3 RCTs)	190 (3 RCTs)
Between-group difference in SMC (95% CI)	-0.26 (-0.77 to 0.24)	-0.15 (-0.34 to 0.05)	-0.42 (-0.59 to -0.26)	-0.30 (-0.53 to -0.06)	-0.01 (-0.26 to 0.28)	-0.03 (-0.26 to 0.31)
Certainty of Evidence ^a	Moderate	Low	Moderate	Moderate	Low	Low
Kamonseki et al (2021)						
	Shoulder pain intensity	Shoulder function				
Total N	250 (5 RCTs)	175 (3 RCTs)				
SMD (95% CI)	-0.21 (-0.67 to 0.34)	-0.11 (-0.41 to 0.19)				
p value (I ²)	.36 (65%)	.48 (0%)				
Quality of Evidence ^a	Very low	Very low				

CI: confidence interval; RCT: randomized controlled trial; SMC: standardized mean change; SMD: standardized mean difference.

^a High certainty: we are very confident that the true effect lies close to that of the estimate of the effect; moderate certainty: we are moderately confident in the effect estimate.; low certainty: our confidence in the effect estimate is limited; very low certainty: we have very little confidence in the effect estimate.

Randomized Controlled Trial

de Oliveira et al (2022) conducted an RCT in 24 patients with subacromial pain syndrome who received exercise or exercise plus EMG biofeedback for 8 weeks. The primary outcomes were pain and shoulder

function. At 8 weeks, pain was better in the exercise-only group (mean numeric pain rating, 0.5 vs 2 with exercise plus biofeedback; $p=.01$); however, this outcome was not different between groups at other time points. The only other significant finding was forward rotation of the scapula, which was better in the biofeedback group at 12 weeks ($p=.006$). All other outcomes were similar between groups.

Ribeiro and Silva (2019) published an RCT assessing whether visual feedback improves range of motion in patients with chronic idiopathic neck pain. Forty-two patients from a single Portuguese clinic were included in the study and randomly assigned to either the visual feedback group ($n=21$) or the control group ($n=21$). There was no effect of time and intervention on pain intensity ($p=.729$), but there was a significant interaction between time and intervention in neck flexion ($p<.001$). The study was limited by its small sample size, short duration of intervention, and by the researcher assessing patients not being blinded.

Orofacial Pain

Systematic Reviews

A Cochrane review by Aggarwal et al (2011) identified 17 trials evaluating nonpharmacologic psychological interventions for adults with chronic orofacial pain (e.g., temporomandibular joint disorder). For studies reporting on short-term pain relief (≤ 3 months), a significantly greater reduction in pain was found for interventions that combined CBT plus biofeedback compared with usual care (2 studies; SMD, 0.46; 95% CI, 0.02 to 0.90). However, when reviewers assessed results from studies reporting on long-term pain relief (≥ 6 months), no significant benefit was found with a combined intervention of CBT plus biofeedback, and there were no studies that compared CBT alone with CBT plus biofeedback. For studies reporting on biofeedback-only interventions, a pooled analysis of 2 studies on short-term pain relief did not find a significant benefit compared with usual care (SMD, -0.41; 95% CI, -1.06 to 0.25). Only 1 study reported long-term pain relief after a biofeedback-only intervention, so a pooled analysis could not be done. Reviewers concluded that there was weak evidence to support psychosocial interventions for managing chronic orofacial pain and the most promising evidence was for CBT, with or without biofeedback. The authors noted that the trials comprising the review were few in number and had a high-risk of bias.

The conclusions drawn from this Cochrane review are similar to those of earlier systematic reviews on the treatment of temporomandibular joint disorder. These older reviews also concluded that there was weak evidence that psychosocial/physical therapy interventions (including biofeedback) are beneficial for treating temporomandibular joint disorder and that, of the few studies available, they tended to be of poor methodologic quality.

Abdominal Pain

Systematic Reviews

In a systematic review of therapies for recurrent abdominal pain in children by Weydert et al (2003), the behavioral interventions of CBT and biofeedback had a generally positive effect on nonspecific recurrent abdominal pain and were deemed safe. The specific effects of biofeedback were not isolated in this systematic review and therefore cannot be assessed.

Randomized Controlled Trials

In a study by Humphreys and Gevirtz (2000), 64 children and teenagers diagnosed with recurrent abdominal pain were randomized to groups treated with increased dietary fiber; fiber and biofeedback; fiber, biofeedback, and CBT; or fiber, biofeedback, CBT, and parental support. The similar nature of the 3 multicomponent treatment groups was associated with greater pain reduction than the fiber-only group. This trial did not address placebo effects.

Fibromyalgia

Systematic Reviews

Glombiewski et al (2013) published a systemic review and meta-analysis of RCTs reporting data on the efficacy of EMG and electroencephalography (EEG) biofeedback (i.e., neurofeedback) for treating patients with fibromyalgia. Reviewers identified 7 RCTs that compared biofeedback with a control method in patients with fibromyalgia. Studies in which biofeedback was evaluated only as part of multicomponent interventions were excluded. Three studies used EEG biofeedback and 4 used EMG biofeedback (N=321 patients). A sham intervention was used as a control condition in 4 studies, 2 using EEG biofeedback and 2 using EMG biofeedback. In a pooled analysis of the studies using EMG biofeedback, a significant reduction in pain intensity was found compared with a different intervention (effect size, Hedges $g=0.86$; 95% CI, 0.11 to 0.62). A pooled analysis of studies on EEG biofeedback did not find a significant benefit in pain reduction compared with control methods. Pooled analyses of studies of EMG and EEG biofeedback did not find a significant benefit of either intervention on other outcomes such as sleep problems, depression, and health-related quality of life. None of the studies reviewed were of high quality, with the risk of bias assessed as unclear or high for all included studies. In addition, all studies reported short-term outcomes, resulting in a lack of evidence on whether longer-term outcomes improved with these interventions.

Randomized Controlled Trials

In a small, double-blind RCT from Asia, Babu et al (2007) compared actual and sham biofeedback for effects on pain, fitness, function, and tender points in 30 patients with fibromyalgia. Pain reduction, as assessed on a VAS, did not differ significantly between groups. The trialists calculated that a sample size of 15 patients could detect a difference of 5 cm (on a 10-cm scale) on a VAS, suggesting that the trial lacked adequate power.

A larger unblinded RCT by van Santen et al (2002) evaluated 143 women with fibromyalgia, and compared EMG biofeedback with fitness training and usual care. The primary outcome was pain measured on a VAS. Compared with usual care, the investigators reported no clear improvements in objective or subjective patient outcomes with biofeedback (or fitness training).

Another RCT on EMG biofeedback for fibromyalgia was conducted by Buckelew et al (1998), and enrolled 119 patients; however, the trial did not follow a double-blind design. Patients were randomized to 1 of 4 treatment groups: (1) biofeedback/relaxation training, (2) exercise training, (3) combination treatment, and (4) an educational/attention control program. While the combination treatment group had better tender point index scores than other treatment groups, this trial did not address placebo effects or the impact of adding biofeedback to relaxation therapy.

Osteoarthritis

Systematic Reviews

A systematic review by Macfarlane et al (2012) evaluated practitioner-based complementary and alternative medicine treatments (defined as any treatment not taken orally or applied topically) for osteoarthritis and identified 2 trials on biofeedback. One was an RCT by Yilmaz et al (2010) that assessed whether the addition of EMG biofeedback to strengthening exercises improved outcomes in 40 patients with knee osteoarthritis.⁴⁴ After a 3-week treatment period, no significant differences between the 2 treatments regarding pain or quality of life were found. The other RCT, by Durmus et al (2007), compared electrical stimulation with biofeedback-assisted exercise in 50 women with knee osteoarthritis. After 4 weeks of treatment, there were no statistically significant differences between groups in pain and functioning scores.

Systematic Lupus Erythematosus

Randomized Controlled Trials

In an RCT by Greco et al (2004), of 92 patients with systemic lupus erythematosus, those treated with 6 sessions of biofeedback-assisted CBT for stress reduction had statistically greater reductions in pain posttreatment than a symptom-monitoring support group ($p=.044$) and a group receiving usual care ($p=.028$). However, these reductions in pain were not sustained at a 9-month follow-up.

Vulvar Vestibulitis

Randomized Controlled Trials

A randomized study by Bergeron et al (2001) of 78 patients with dyspareunia resulting from vulvar vestibulitis compared treatment with EMG biofeedback, surgery, or CBT.⁴⁷ Patients who underwent surgery had significantly lower pain scores than patients who received biofeedback or CBT. No placebo treatment was used.

Home Biofeedback

No relevant studies on the clinical effectiveness of biofeedback using home equipment were identified.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Occupational and Environmental Medicine

In 2020, the American College of Occupational and Environmental Medicine updated their guideline on noninvasive and minimally invasive management of low back disorders. The role of biofeedback is not addressed in this updated guideline.

American Society of Anesthesiologists and American Society of Regional Anesthesia and Pain Medicine

In 2010, practice guidelines from the American Society of Anesthesiologists and the American Society of Regional Anesthesia and Pain Medicine suggested that "cognitive behavioral therapy, biofeedback, or relaxation training....may be used as part of a multimodal strategy for patients with low back pain, as well as for other chronic pain conditions."

National Comprehensive Cancer Network (NCCN)

Adult Cancer Pain Version 2.2025

The current guideline for National Comprehensive Cancer Network (NCCN) on the management of adult cancer pain, biofeedback is recommended as an optional component of an integrative intervention to reduce pain (2A recommendation).

"Cognitive interventions are aimed at enhancing a sense of control over the pain or underlying disease. Cognitive behavioral therapy (CBT), cognitive restructuring, mindfulness-based stress reduction (MBSR), breathing exercises, relaxation, yoga, imagery, hypnosis, biofeedback, music and other behavioral therapies can be very useful. Patient-based educational interventions have a significant impact in providing pain relief. Skills training helps modify the patient's experience of pain and helps patients acquire techniques of pain management such as deep muscle relaxation. Patients who may benefit from skills training may be referred to a licensed mental health professional trained in CBT, hypnosis, biofeedback, or MBSR. Education provides patients and family/caregivers with the knowledge to use analgesics correctly and to address side effects or unrelieved pain."

North American Spine Society

In 2020, the North American Spine Society published a guideline for the diagnosis and treatment of low back pain. Although nonpharmacologic therapies are addressed in this guideline, the specific role of biofeedback for low back pain is not addressed.

U.S. Department of Veterans Affairs and U.S. Department of Defense

In 2022, the U.S. Department of Veterans Affairs and U.S. Department of Defense updated their guideline on the diagnosis and treatment of low back pain. The guideline recommends several nonpharmacologic therapies for chronic low back pain (e.g., cognitive-behavioral therapy [CBT] and/or mindfulness-based stress reduction, progressive relaxation, exercise including yoga, pilates, and tai chi) but does not address the role of biofeedback.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review can be located at clinicaltrials.gov.

REFERENCES

1. U.S. Department of Health and Human Services. Pain management best practices. May 2019. <https://www.hhs.gov/sites/default/files/pain-mgmt-best-practices-draft-final-report-05062019.pdf>.
2. Dworkin RH, Turk DC, Farrar JT, et al. Core outcome measures for chronic pain clinical trials: IMMPACT recommendations. *Pain*. Jan 2005; 113(1-2): 9-19. PMID 15621359
3. Dworkin RH, Turk DC, Wyrwich KW, et al. Interpreting the clinical importance of treatment outcomes in chronic pain clinical trials: IMMPACT recommendations. *J Pain*. Feb 2008; 9(2): 105-21. PMID 18055266
4. Williams ACC, Fisher E, Hearn L, et al. Psychological therapies for the management of chronic pain (excluding headache) in adults. *Cochrane Database Syst Rev*. Aug 12 2020; 8: CD007407. PMID 32794606
5. Fisher E, Law E, Dudeney J, et al. Psychological therapies for the management of chronic and recurrent pain in children and adolescents. *Cochrane Database Syst Rev*. Sep 29 2018; 9: CD003968. PMID 30270423
6. Humphreys PA, Gevirtz RN. Treatment of recurrent abdominal pain: components analysis of four treatment protocols. *J Pediatr Gastroenterol Nutr*. Jul 2000; 31(1): 47-51. PMID 10896070
7. Palermo TM, Eccleston C, Lewandowski AS, et al. Randomized controlled trials of psychological therapies for management of chronic pain in children and adolescents: an updated meta-analytic review. *Pain*. Mar 2010; 148(3): 387-397. PMID 19910118
8. Henschke N, Ostelo RW, van Tulder MW, et al. Behavioural treatment for chronic low-back pain. *Cochrane Database Syst Rev*. Jul 07 2010; (7): CD002014. PMID 20614428
9. Kapitza KP, Passie T, Bernateck M, et al. First non-contingent respiratory biofeedback placebo versus contingent biofeedback in patients with chronic low back pain: a randomized, controlled, double-blind trial. *Appl Psychophysiol Biofeedback*. Sep 2010; 35(3): 207-17. PMID 20237953
10. Tan G, Rintala DH, Jensen MP, et al. A randomized controlled trial of hypnosis compared with biofeedback for adults with chronic low back pain. *Eur J Pain*. Feb 2015; 19(2): 271-80. PMID 24934738
11. Glombiewski JA, Hartwich-Tersek J, Rief W. Two psychological interventions are effective in severely disabled, chronic back pain patients: a randomized controlled trial. *Int J Behav Med*. Jun 2010; 17(2): 97-107. PMID 19967572
12. Karaborklu Argut S, Celik D, Yasaci Z. Effectiveness of therapeutic electromyographic biofeedback after orthopedic knee surgeries: a systematic review. *Disabil Rehabil*. Jul 2022; 44(14): 3364-3372. PMID 33417500

13. Collins NJ, Bisset LM, Crossley KM, et al. Efficacy of nonsurgical interventions for anterior knee pain: systematic review and meta-analysis of randomized trials. *Sports Med.* Jan 01 2012; 42(1): 31-49. PMID 22149696
14. Campo M, Zadro JR, Pappas E, et al. The effectiveness of biofeedback for improving pain, disability and work ability in adults with neck pain: A systematic review and meta-analysis. *Musculoskelet Sci Pract.* Apr 2021; 52: 102317. PMID 33461043
15. Kamonseki DH, Calixtre LB, Barreto RPG, et al. Effects of electromyographic biofeedback interventions for shoulder pain and function: Systematic review and meta-analysis. *Clin Rehabil.* Jul 2021; 35(7): 952-963. PMID 33517777
16. Juul-Kristensen B, Larsen CM, Eshoj H, et al. Positive effects of neuromuscular shoulder exercises with or without EMG-biofeedback, on pain and function in participants with subacromial pain syndrome - A randomised controlled trial. *J Electromyogr Kinesiol.* Oct 2019; 48: 161-168. PMID 31394380
17. Kosterink SM, Huis in 't Veld RM, Cagnie B, et al. The clinical effectiveness of a myofeedback-based teletreatment service in patients with non-specific neck and shoulder pain: a randomized controlled trial. *J Telemed Telecare.* 2010; 16(6): 316-21. PMID 20798425
18. Ma C, Szeto GP, Yan T, et al. Comparing biofeedback with active exercise and passive treatment for the management of work-related neck and shoulder pain: a randomized controlled trial. *Arch Phys Med Rehabil.* Jun 2011; 92(6): 849-58. PMID 21621660
19. Middaugh S, Thomas KJ, Smith AR, et al. EMG Biofeedback and Exercise for Treatment of Cervical and Shoulder Pain in Individuals with a Spinal Cord Injury: A Pilot Study. *Top Spinal Cord Inj Rehabil.* 2013; 19(4): 311-23. PMID 24244096
20. Sandsjo L, Larsman P, Huis in 't Veld RM, et al. Clinical evaluation of a myofeedback-based teletreatment service applied in the workplace: a randomized controlled trial. *J Telemed Telecare.* 2010; 16(6): 329-35. PMID 20798427
21. Arami J, Rezasoltani A, Khalkhali Z, et al. The effect of two exercise therapy programs (proprioceptive and endurance training) to treat patients with chronic non-specific neck pain. *JBUMS.* 2012;14(1):77-84.
22. Bissett A, Mitchell KR, Major G. The cervico-brachial pain syndrome: muscle activity and pain relief. *Behav Change.* 1985;2(2):129-132.
23. Bobos P, Billis E, Papanikolaou DT, et al. Does Deep Cervical Flexor Muscle Training Affect Pain Pressure Thresholds of Myofascial Trigger Points in Patients with Chronic Neck Pain? A Prospective Randomized Controlled Trial. *Rehabil Res Pract.* 2016; 2016: 6480826. PMID 27990302
24. Dellve L, Ahlstrom L, Jonsson A, et al. Myofeedback training and intensive muscular strength training to decrease pain and improve work ability among female workers on long-term sick leave with neck pain: a randomized controlled trial. *Int Arch Occup Environ Health.* Mar 2011; 84(3): 335-46. PMID 20803028
25. Ehrenborg C, Archenholtz B. Is surface EMG biofeedback an effective training method for persons with neck and shoulder complaints after whiplash-associated disorders concerning activities of daily living and pain -- a randomized controlled trial. *Clin Rehabil.* Aug 2010; 24(8): 715-26. PMID 20562165
26. Eslamian F, Jahanjoo F, Dolatkah N, et al. Relative Effectiveness of Electroacupuncture and Biofeedback in the Treatment of Neck and Upper Back Myofascial Pain: A Randomized Clinical Trial. *Arch Phys Med Rehabil.* May 2020; 101(5): 770-780. PMID 31954696
27. Iqbal ZA, Rajan R, Khan SA, et al. Effect of deep cervical flexor muscles training using pressure biofeedback on pain and disability of school teachers with neck pain. *J Phys Ther Sci.* Jun 2013; 25(6): 657-61. PMID 24259822
28. Jull G, Trott P, Potter H, et al. A randomized controlled trial of exercise and manipulative therapy for cervicogenic headache. *Spine (Phila Pa 1976).* Sep 01 2002; 27(17): 1835-43; discussion 1843. PMID 12221344

29. Jull G, Falla D, Treleaven J, et al. Retraining cervical joint position sense: the effect of two exercise regimes. *J Orthop Res*. Mar 2007; 25(3): 404-12. PMID 17143898
30. Nezamuddin M, Answer S, Khan SA, et al. Efficacy of pressure-biofeedback guided deep cervical flexor training on neck pain and muscle performance in visual display terminal operators. *J Musculoskelet Res*. 2013;16(3):1350011
31. Voerman GE, Sandsjo L, Vollenbroek-Hutten MM, et al. Effects of ambulant myofeedback training and ergonomic counselling in female computer workers with work-related neck-shoulder complaints: a randomized controlled trial. *J Occup Rehabil*. Mar 2007; 17(1): 137-52. PMID 17260162
32. Wani S, Raka N, Jethwa J, et al. Comparative efficacy of cervical retraction exercises (McKenzie) with and without using pressure biofeedback in cervical spondylosis. *Int J Ther Rehabil*. 2013;20(10):501-508.
33. de Oliveira AKA, da Costa KSA, de Lucena GL, et al. Comparing exercises with and without electromyographic biofeedback in subacromial pain syndrome: A randomized controlled trial. *Clin Biomech (Bristol, Avon)*. Mar 2022; 93: 105596. PMID 35183878
34. Ribeiro D, Silva AG. A single session of visual feedback improves range of motion in patients with chronic idiopathic neck pain: A randomized and controlled study. *Musculoskeletal Care*. Mar 2019; 17(1): 72-78. PMID 30378756
35. Aggarwal VR, Lovell K, Peters S, et al. Psychosocial interventions for the management of chronic orofacial pain. *Cochrane Database Syst Rev*. Nov 09 2011; (11): CD008456. PMID 22071849
36. McNeely ML, Armijo Olivo S, Magee DJ. A systematic review of the effectiveness of physical therapy interventions for temporomandibular disorders. *Phys Ther*. May 2006; 86(5): 710-25. PMID 16649894
37. Medlicott MS, Harris SR. A systematic review of the effectiveness of exercise, manual therapy, electrotherapy, relaxation training, and biofeedback in the management of temporomandibular disorder. *Phys Ther*. Jul 2006; 86(7): 955-73. PMID 16813476
38. Weydert JA, Ball TM, Davis MF. Systematic review of treatments for recurrent abdominal pain. *Pediatrics*. Jan 2003; 111(1): e1-11. PMID 12509588
39. Glombiewski JA, Bernardy K, Hauser W. Efficacy of EMG- and EEG-Biofeedback in Fibromyalgia Syndrome: A Meta-Analysis and a Systematic Review of Randomized Controlled Trials. *Evid Based Complement Alternat Med*. 2013; 2013: 962741. PMID 24082911
40. Babu AS, Mathew E, Danda D, et al. Management of patients with fibromyalgia using biofeedback: a randomized control trial. *Indian J Med Sci*. Aug 2007; 61(8): 455-61. PMID 17679735
41. van Santen M, Bolwijn P, Verstappen F, et al. A randomized clinical trial comparing fitness and biofeedback training versus basic treatment in patients with fibromyalgia. *J Rheumatol*. Mar 2002; 29(3): 575-81. PMID 11908576
42. Buckelew SP, Conway R, Parker J, et al. Biofeedback/relaxation training and exercise interventions for fibromyalgia: a prospective trial. *Arthritis Care Res*. Jun 1998; 11(3): 196-209. PMID 9782811
43. Macfarlane GJ, Paudyal P, Doherty M, et al. A systematic review of evidence for the effectiveness of practitioner-based complementary and alternative therapies in the management of rheumatic diseases: osteoarthritis. *Rheumatology (Oxford)*. Dec 2012; 51(12): 2224-33. PMID 22923762
44. Yilmaz OO, Senocak O, Sahin E, et al. Efficacy of EMG-biofeedback in knee osteoarthritis. *Rheumatol Int*. May 2010; 30(7): 887-92. PMID 19693508
45. Durmus D, Alayli G, Canturk F. Effects of quadriceps electrical stimulation program on clinical parameters in the patients with knee osteoarthritis. *Clin Rheumatol*. May 2007; 26(5): 674-8. PMID 16897119

46. Greco CM, Rudy TE, Manzi S. Effects of a stress-reduction program on psychological function, pain, and physical function of systemic lupus erythematosus patients: a randomized controlled trial. *Arthritis Rheum.* Aug 15 2004; 51(4): 625-34. PMID 15334437
47. Bergeron S, Binik YM, Khalife S, et al. A randomized comparison of group cognitive-behavioral therapy, surface electromyographic biofeedback, and vestibulectomy in the treatment of dyspareunia resulting from vulvar vestibulitis. *Pain.* Apr 2001; 91(3): 297-306. PMID 11275387
48. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med.* Apr 04 2017; 166(7): 514-530. PMID 28192789
49. Hegmann KT, Travis R, Andersson GBJ, et al. Non-Invasive and Minimally Invasive Management of Low Back Disorders. *J Occup Environ Med.* Mar 2020; 62(3): e111-e138. PMID 31977923
50. Benzon HT, Connis RT, De Leon-Casasola OA, et al. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology.* Apr 2010; 112(4): 810-33. PMID 20124882
51. VA/DoD Clinical Practice Guideline. (2022). The Diagnosis and Treatment of Low Back Pain. Washington, DC: U.S. Government Printing Office. Version 3.0.
<https://www.healthquality.va.gov/guidelines/Pain/lbp/VADoDLBPCPGFinal508.pdf>.
52. Kreiner DS, Matz P, Bono CM, et al. Guideline summary review: an evidence-based clinical guideline for the diagnosis and treatment of low back pain. *Spine J.* Jul 2020; 20(7): 998-1024. PMID 32333996
53. National Comprehensive Cancer Network (NCCN) Adult Cancer Pain Version 3.2024. Also available at <https://www.nccn.org>
54. UpToDate. Approach to the management of chronic non-cancer pain in adults. Topic last updated November 2025. Also available at <https://www.uptodate.com>
55. UpToDate. Integrative pediatric health care. Topic last updated September 2025. Also available at <https://www.uptodate.com>
56. UpToDate. Complementary and integrative health in pediatrics. Kathi J. Kemper M.D., MPH. Topic last updated November 2022. Also available at <https://www.uptodate.com>
57. UpToDate. Rehabilitative and integrative therapies for pain in patients with cancer. Topic last updated April 2024. Also available at <https://www.uptodate.com>
58. UpToDate. Pain in children: Approach to pain assessment and overview of management principles. Topic last updated August 2025. Also available at <https://www.uptodate.com>
59. UpToDate. Management of knee osteoarthritis. Topic last updated October 2025. Also available at <https://www.uptodate.com>
60. UpToDate. Overview of the management of osteoarthritis. Topic last updated November 2025. Also available at <https://www.uptodate.com>
61. UpToDate. Subacute and chronic low back pain: Management. Topic last updated November 2025. Also available at <https://www.uptodate.com>
62. UpToDate. Central neuropathic facial pain. Topic last updated January 2024. Also available at <https://www.uptodate.com>
63. UpToDate. Systemic lupus erythematosus in adults: Overview of the management and prognosis. Topic last updated November 2025. Also available at <https://www.uptodate.com>
64. UpToDate. Treatment of Raynaud phenomenon: Initial management. Topic last updated December 2024. Also available at <https://www.uptodate.com>
65. UpToDate. Fibromyalgia: Treatment in adults. Topic last updated September 2025. Also available at <https://www.uptodate.com>
66. Lazaridou A, Paschali M, Vilsmark ES, et al. Biofeedback EMG alternative therapy for chronic low back pain (the BEAT-pain study). *Digit Health.* 2023; 9: 20552076231154386. PMID 36776410
67. Hayes, a symplr company. Health Technology Assessment Biofeedback for Headache and Chronic Musculoskeletal Pain. November 3, 2004. Annual Review December 7, 2008

68. Spada EG, Masiero M, Pizzoli MSF, et al Heart rate variability biofeedback in cancer patients: A scoping review. Behav Sci. 2022, 12, 389 <https://doi.org/10.3390/bs12100389>
69. Ananías J, Vidal C, Ortiz-Muñoz L, et al. Use of electromyographic biofeedback in rehabilitation following anterior cruciate ligament reconstruction: a systematic review and meta-analysis. Physiotherapy. Jun 2024; 123: 19-29. PMID 38244487
70. Yeldan I, Canan GD, Akinci B. Biofeedback Sensor vs. Physiotherapist Feedback During Core Stabilization Training in Patients with Chronic Nonspecific Low Back Pain. Appl Psychophysiol Biofeedback. Mar 2024; 49(1): 103-113. PMID 37878122
71. Calderone A, Mazzurco Masi VM, De Luca R, et al. The impact of biofeedback in enhancing chronic pain rehabilitation: A systematic review of mechanisms and outcomes. Heliyon. Jan 30 2025; 11(2): e41917. PMID 39897804

CODES

To report provider services, use appropriate CPT codes, HCPCS codes, Revenue codes, and/or ICD diagnosis codes.

Codes	Number	Description
CPT		
	90875	Individual psychophysiological therapy incorporating biofeedback training by any modality (face-to-face with the patient), with psychotherapy (e.g., insight oriented, behavior modifying or supportive psychotherapy); 30 minutes
	90876	Individual psychophysiological therapy incorporating biofeedback training by any modality (face-to-face with the patient), with psychotherapy (e.g., insight oriented, behavior modifying or supportive psychotherapy); 45 minutes
	90901	Biofeedback training by any modality
	90912	Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry, when performed initial 15 minutes of one-on-one physician or other qualified health care professional contact with the patient
	90913	Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry, when performed; each additional 15 minutes of one-on-one physician or other qualified health care professional contact with the patient (List separately in addition to code for primary procedure)
HCPCS		
	E0746	Electromyography (EMG), biofeedback device

	E1399	Durable medical equipment, miscellaneous (may be used for home biofeedback devices which may include the following: FreeSpira, Heart Rate Variability, Ileva® Pelvic Health System, or StressEraser)
	E3200	Gait modulation system, rhythmic auditory stimulation, including restricted therapy software, all components and accessories, prescription only (InTandem)
Type of Service	Medicine	
Place of Service	Physician Office	

POLICY HISTORY

Date	Action	Action
December 2025	Annual Review	Policy Revised
December 2024	Annual Review	Policy Revised
August 2024	Annual Review	Policy Renewed
August 2023	Annual Review	Policy Revised – Content moved from retired medical policy “Biofeedback”
June 2022	Interim Review	Policy Revised and biofeedback content incorporated into this medical policy from medical policies: Biofeedback/Neurofeedback as a Treatment of Headaches and Other Disorders and Temporomandibular Joint Dysfunction (TMJ/TMD) Treatment
June 2021	Annual Review	Policy Revised
July 2020	Annual Review	Policy Revised
July 2019	Annual Review	Policy Revised
July 2018	Annual Review	Policy Revised
July 2017	Annual Review	Policy Revised
July 2016	Annual Review	Policy Revised
August 2015	Annual Review	Policy Revised
September 2014	Annual Review	Policy Revised

Date	Action	Action
April 2014	Interim Review	Policy Revised
October 2013	Annual Review	Policy Revised
December 2012	Annual Review	Policy Renewed
December 2011	Annual Review	Policy Renewed
December 2010	Annual Review	Policy Renewed

New information or technology that would be relevant for Wellmark to consider when this policy is next reviewed may be submitted to:

Wellmark Blue Cross and Blue Shield
 Medical Policy Analyst
 PO Box 9232
 Des Moines, IA 50306-9232

*CPT® is a registered trademark of the American Medical Association.