

02.01.17 Bio-Engineered Skin and Soft Tissue Substitutes

Original Effective Date: May 1999

Review Date: May 2026

Revised: July 2026

DISCLAIMER/INSTRUCTIONS FOR USE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations, or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

This Medical Policy document describes the status of medical technology at the time the document was developed. Since that time, new technology may have emerged, or new medical literature may have been published. This Medical Policy will be reviewed regularly and updated as scientific and medical literature becomes available; therefore, policies are subject to change without notice.

Related Policies:

- [02.01.60 Amniotic Membrane and Amniotic Fluid Grafts](#)

Summary

Description

Bioengineered skin and soft tissue substitutes may be derived from human tissue (autologous or allogeneic), nonhuman tissue (xenographic), synthetic materials, or a composite of these materials. Bioengineered skin and soft tissue substitutes are being evaluated for a variety of conditions, including breast reconstruction (prophylactic or oncologic indications) and healing lower-extremity ulcers and severe burns. Acellular dermal matrix (ADM) products are also being evaluated for soft tissue repair.

Summary of Evidence

Breast Reconstruction

For individuals who are undergoing breast reconstruction who receive allogeneic acellular dermal matrix (ADM) products, the evidence includes randomized controlled trials (RCTs) and systematic reviews. Relevant outcomes are symptoms, morbid events, functional outcomes, quality of life (QOL), and treatment-related morbidity. A systematic review found no difference in overall complication rates with ADM allograft compared with standard procedures for breast reconstruction. Reconstructions with ADM have been reported to have higher seroma, infection, and necrosis rates than reconstructions without ADM. However, capsular contracture and malposition of implants may be reduced. Thus, in cases where there is limited tissue coverage, the available evidence may inform patient decision making about reconstruction options. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Tendon Repair

For individuals who are undergoing tendon repair who receive GraftJacket, the evidence includes an RCT. Relevant outcomes are symptoms, morbid events, functional outcomes, QOL, and treatment-related morbidity. The RCT identified found improved outcomes with the GraftJacket ADM allograft for rotator cuff repair. Although these results were positive, additional studies with a larger number of individuals is needed to evaluate the consistency of the effect. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Surgical Repair of Hernias or Parastomal Reinforcement

For Individuals undergoing surgical repair of hernias or parastomal reinforcement who receive Alloderm, Strattice or OviTex, the evidence includes RCTs. Relevant outcomes are symptoms, morbid events, functional outcomes, QOL, and treatment-related morbidity. Several comparative studies including RCTs have shown no difference in outcomes between tissue-engineered skin substitutes and either standard synthetic mesh or no reinforcement. The evidence is insufficient to determine that technology results in an improvement in the net health outcome. However, even though there is a paucity in the peer-reviewed scientific evidence in showing consistent evidence regarding the effectiveness of the use of tissue - engineered skin substitutes for individuals undergoing surgical repair of hernias or parastomal reinforcement, Alloderm, Strattice or OviTex will be considered medically necessary when the criteria below is met, see [Policy](#).

Diabetic Lower-Extremity Ulcers

For individuals who have diabetic lower-extremity ulcers who receive AlloPatch, Apligraf, Dermagraft, Integra, mVASC, or TheraSkin the evidence includes RCTs. Relevant outcomes are symptoms, change in disease status, morbid events, and QOL. RCTs reporting complete wound healing outcomes with at least 12 weeks follow-up have demonstrated the efficacy of AlloPatch (reticular ADM), Apligraf and Dermagraft (living cell therapy), Integra (biosynthetic), mVASC, TheraSkin, and Omega3 Wound (Kerecis) over the standard of care (SOC). The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have diabetic lower-extremity ulcers who receive bioengineered skin substitutes other than AlloPatch, Apligraf, Dermagraft, Integra, mVASC, TheraSkin, or Omega3 Wound (Kerecis), the evidence includes RCTs. Relevant outcomes are symptoms, change in disease status, morbid events,

and QOL. Results from a multicenter RCT showed some benefit of DermACELL that was primarily for the subgroup of individuals who only required a single application of the ADM. Studies are needed to further define the population who might benefit from this treatment. Additional study with a larger number of subjects is needed to evaluate the effect of GraftJacket, DermACELL, Cytal, PriMatrix, and Oasis Wound Matrix, compared with current SOC or other advanced wound therapies. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Lower-Extremity Ulcers due to Venous Insufficiency

For individuals who have lower-extremity ulcers due to venous insufficiency who receive Apligraf or Oasis Wound Matrix, the evidence includes RCTs. Relevant outcomes are symptoms, change in disease status, morbid events, and QOL. RCTs have demonstrated the efficacy of Apligraf living cell therapy and xenogeneic Oasis Wound Matrix over the SOC. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have lower-extremity ulcers due to venous insufficiency who receive bioengineered skin substitutes other than Apligraf or Oasis Wound Matrix, the evidence includes RCTs. Relevant outcomes are disease-specific survival, symptoms, change in disease status, morbid events, and QOL. In a moderately large RCT, Dermagraft was not shown to be more effective than controls for the primary or secondary endpoints in the entire population and was only slightly more effective than controls (an 8%-15% increase in healing) in subgroups of patients with ulcer durations of 12 months or less or size of 10 cm or less. Additional studies with a larger number of subjects is needed to evaluate the effect of the xenogeneic PriMatrix skin substitute versus the current SOC. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Dystrophic Epidermolysis Bullosa

For individuals who have dystrophic epidermolysis bullosa who receive OrCel, the evidence includes case series. Relevant outcomes are symptoms, change in disease status, morbid events, and QOL. OrCel was approved under a humanitarian drug exemption for use in individuals with dystrophic epidermolysis bullosa undergoing hand reconstruction surgery, to close and heal wounds created by the surgery, including those at donor sites. Outcomes have been reported in a small series (e.g., 5 patients). The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Deep Dermal Burns

For individuals who have deep dermal burns who receive bioengineered skin substitutes (i.e., Epicel, Integra, Biobrane, EZ Derm, AlloSkin, TheraSkin, Trancyte), the evidence includes RCTs. Relevant outcomes are symptoms, change in disease status, morbid events, functional outcomes, QOL, and treatment-related morbidity. Overall, few skin substitutes have been approved, and the evidence is limited for each product. Epicel (living cell therapy) has received U.S. Food and Drug administration approval under a humanitarian device exemption for the treatment of deep dermal or full-thickness burns comprising a total body surface area of 30% or more. Comparative studies have demonstrated improved outcomes for biosynthetic skin substitute Integra Dermal Regeneration Template for the treatment of burns. Based on clinical practice standards, the below-mentioned studies, and the overall clinical experience with Biobrane, EZ Derm, AlloSkin, TheraSkin and Trancyte an acceptable level of safety and efficacy has been established for the use of this product for the treatment of burns. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with deep dermal burns who are treated with the ReCell autologous cell harvesting device, the evidence includes 2 RCTs. Relevant outcomes are symptoms, change in disease status, morbid events, functional outcomes, QOL, and treatment-related morbidity. One RCT evaluated ReCell for use in combination with meshed autografting and the other compared ReCell head-to-head with skin grafting. Although the ReCell device was comparable to standard care on outcomes such as complete wound closure, confidence in the strength of the overall body of evidence is limited by individual study limitations and heterogeneity of populations, interventions, and outcome measures across studies. Additional RCT evidence in the intended use population is needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Additional Information

Not applicable

OBJECTIVE

The objective of this review is to determine whether the use of artificial skin and soft tissue substitutes for reinforcement for surgical procedures and healing of chronic wounds and burns improves the net health outcome.

PRIOR APPROVAL

Not applicable.

POLICY

Breast Reconstruction Surgery

The following tissue engineered skin substitute products may be considered **medically necessary** when used for breast reconstruction surgery (see [Policy Guidelines](#)):

- AlloDerm® (Q4116)
- Cortiva® (AlloMax™) (Q4433)
- DermACELL™ (Q4122)
- DermaMatrix™ (Q4433)
- FlexHD®, FlexHD® Pliable™ (Q4128)
- Strattice (Q4130)

The tissue engineered skin substitute products listed above are considered **investigational** when the above criteria is not met and for all other indications because the evidence is insufficient to determine that the technology results in an improvement in net health outcomes.

Burns

The following tissue engineered skin substitute products may be considered **medically necessary** when used for the treatment of second and third-degree burns:

- Biobrane and Biobrane Glove (A2043, A2044)

- Ez Derm (Q4136)
- Epicel® (Q4431, Q4432, Q4433) (for the treatment of deep dermal or full-thickness burns comprising a total body surface area $\geq 30\%$ when provided in accordance with the HDE specifications of the FDA, see medical policy [10.01.14 Humanitarian Use Devices](#))
- Fresh frozen unprocessed allograft skin products (for example AlloSkin (Q4115), TheraSkin Q4121); or
- Integra® Bilayer Matrix Wound Dressing (Q4104)
- Integra® Omnigraft™ Dermal Regeneration Template (Q4105)
- Transcyte (Q4182)

Treatment of fresh, clean, split-thickness donor site wounds in burn victims using the following tissue-engineered skin substitute product may be considered **medically necessary**:

- OrCel (Q4431)

The tissue engineered skin substitute products listed above are considered **investigational** when the above criteria is not met and for all other indications because the evidence is insufficient to determine that the technology results in an improvement in net health outcomes.

Surgical Repair of Hernias or Parastomal Reinforcement

The following tissue engineered skin substitute products may be considered **medically necessary** for the surgical repair of complex abdominal wall wounds to include repair of hernias or parastomal reinforcement:

- AlloDerm® (Q4116)
- OviTex™ (A4100 or Q4432)
- Strattice (Q4130)

The tissue engineered skin substitute products listed above are considered **investigational** when the above criteria is not met and for all other indications because the evidence is insufficient to determine that the technology results in an improvement in net health outcomes.

Diabetic Lower Extremity Ulcer(s)

The following tissue engineered skin substitute products may be considered **medically necessary** for the treatment of chronic, non-infected full thickness diabetic lower extremity ulcer(s) that have not healed with standard conservative therapy (such as surgical debridement, complete off loading, and standard dressing changes) attempted for at least 2 to 4 weeks (past start of therapy) with the following supporting documentation: the ulcer(s) has failed to decrease in size and depth, or there is no change in baseline size or depth with no sign of improvement, or no indication that improvement is likely:

- AlloPatch® (Q4128)
- Apligraf® (Q4101)
- Integra® Omnigraft™ Dermal Regeneration Matrix (also known as Omnigraft) Q4105 and Integra® Flowable™ Wound Matrix (Q4114)
- Kerecis™ Omega3 (Q4158)
- mVASC® (A4100 or Q4433)
- TheraSkin® (Q4182)

Note: For utilization of amniotic membrane and amniotic fluid grafts for diabetic lower extremity ulcer(s) see medical policy [02.01.60 Amniotic Membrane and Amniotic Fluid Grafts](#).

The tissue engineered skin substitute products listed above are considered **investigational** when the above criteria is not met and for all other indications because the evidence is insufficient to determine that the technology results in an improvement in net health outcomes.

Venous Stasis Ulcers

The following tissue engineered skin substitute products may be considered **medically necessary** for the treatment of chronic, non-infected partial or full-thickness lower extremity skin ulcers due to venous insufficiency which have not adequately responded following a 1-month period of conventional ulcer therapy:

- Apligraf® (Q4101)
- Oasis™ Wound Matrix (Q4102)

Note: For utilization of amniotic membrane and amniotic fluid grafts for venous stasis ulcer(s) see medical policy [02.01.60 Amniotic Membrane and Amniotic Fluid Grafts](#).

The tissue engineered skin substitute products listed above are considered **investigational** when the above criteria are not met and for all other indications because the evidence is insufficient to determine that the technology results in an improvement in net health outcomes.

Dystrophic Epidermolysis Bullosa

The following tissue engineered skin substitute product may be considered **medically necessary** for the treatment of dystrophic epidermolysis bullosa:

- OrCel (Q4431) (for the treatment of mitten-hand deformity when standard wound therapy has failed and when provided in accordance with the humanitarian device exemption [HDE] specifications of the U.S. Food and Drug Administration [FDA], see medical policy [10.01.14 Humanitarian Use Devices](#))

The tissue engineered skin substitute product listed above is considered **investigational** when the above criteria are not met and for all other indications because the evidence is insufficient to determine that the technology results in an improvement in net health outcomes.

Investigational

All other tissue engineered skin substitute products including but not limited to the following are considered **investigational** for all indications because the evidence is insufficient to determine that the technology results in an improvement in net health outcomes:

- ACell® UBM Hydrated/Lyophilized Wound Dressing
- AC5 Advanced wound System (Ac5)
- Allomend®
- AlloSkin™ RT
- Apis®
- Aongen™ Collagen Matrix

- Architect® ECM, PX, FX
- Artacent® Wound
- ArthroFlex™ (Flex Graft)
- AxoGuard® Nerve Protector (AxoGen)
- Bio-ConneKt® Wound Matrix
- Cohealyx™ collagen dermal matrix
- CollaCare®
- CollaCare® Dental
- Collagen Wound Dressing (Oasis Research)
- CollaGUARD®
- CollaMend™
- CollaWound™
- Coll-e-derm
- Collexa®
- Collieva®
- Conexa™
- Coreleader Colla-Pad
- CorMatrix®
- Cymetra™ (Micronized AlloDerm)™
- Cytal™ (previously MatriStem®)
- DeNovoSkin™
- Dermadapt™ Wound Dressing
- Derma-gide
- DermaPure™
- DermaSpan™
- DressSkin
- Durepair Regeneration Matrix®
- Endoform Dermal Template™
- *ENDURAG*en™
- Excellagen®
- ExpressGraft™
- Flexible Collagen Nerve Cuff
- FlowerDerm™
- Foundation Dermal Regeneration Scaffold (DRS) + DUO
- Foundation Dermal Regeneration Scaffold (DRS) + Solo
- G4Derm™ Plus
- GammaGraft
- Geistlich Derma-Gide™
- Graftjacket®
- GraftJacket® Xpress, injectable
- Helicoll™
- hMatrix®
- Hyalomatrix®
- Hyalomatrix® PA
- Innovaburn
- InnovaMatrix® FD
- InnovaMatrix® PD
- InnovaMatrix® XL

- Integra™ Bilayer Wound Matrix
- Integra® Matrix Wound Dressing (previously Avagen)
- InteguPly®
- Keramatrix®
- Keroxx™
- Marigen™ Pacto
- MatriDerm®
- MatriStem
- Matrix HD™
- Mediskin®
- MemoDerm™
- Microlyte Matrix®
- Microlyte Paingaurd®
- MicroTract wound matrix
- Microderm® biologic wound matrix
- Miro3d
- Miro3d fibers, per milligram
- Mirodry wound matrix
- Mochida Nerve Cuff
- MyOwn skin
- Myriad matrix
- Myriad morcells, 4 milligrams
- Neomatrix
- Nerve Align Nerve Cuff
- Nerve Tape
- Neurawrap
- NeuroMend
- NeuroShield
- Novashield or Novagen Wound Matrix
- NovoSorb™ Biodegradable Temporizing Matrix (BMT)
- Oasis® Burn Matrix
- Oasis® Ultra
- Ologen™ Collagen Matrix
- Omega3 Wound (originally Merigen wound dressing)
- Omeza® Collagen Matrix
- Permacol™
- PermeaDerm® B
- PermaDerm® C
- PermaDerm® Glove
- Phoenix™ Wound Matrix
- PriMatrix™
- PriMatrix™ Dermal Repair Scaffold
- Progenamatrix
- Puracol® and Puracol® Plus Collagen Wound Dressings
- PuraPly™ Wound Matrix (previously FortaDerm™)
- PuraPly™ AM (Antimicrobial Wound Matrix)
- Puros® Dermis
- RegenePro™

- Reinforce Flexible Collagen Nerve Cuff
- Repliform®
- ReCell®
- Repriza™
- Resolve matrix
- Resstrata®/Restrata® minimatrix
- SkinTE™
- StrataGraft®
- Suprathel®
- SurgiMend®
- Symphony™
- Talymed®
- TenoGlide™
- TenSIX™ Acellular Dermal Matrix
- TissueMend
- TheraForm™ Standard/Sheet
- TruSkin™
- Tutomesh™ Fenestrated Bovine Pericardium
- Veritas® Collagen Matrix
- Versawrap nerve protector
- Xcellistem®
- XCM Biologic® Tissue Matrix
- XenMatrix™ AB

POLICY GUIDELINES

Category III codes are a set of temporary (T) codes for emerging technologies, services, and procedures that allow for data collection by the American Medical Association's (AMA). If a Category III code is available, providers must use that code instead of an unlisted or deleted Category I code. The services or procedures represented by Category III codes may not have FDA approval, may not be performed by many health care professionals across the country, and the service or procedure may not have proven clinical efficacy. Certain T codes may be addressed as a covered service in a Wellmark BlueCross BlueShield Medical Coverage Policy. But, unless there is explicit Policy criteria that specifically extends coverage to a particular Category III code, the code would generally be considered experimental, investigational, or unproven.

There is no standard definition of "skin substitute". Products in this review cover products that do not require U.S. Food and Drug Administration (FDA) approval or clearance as well as a number of products cleared through the 510(k) pathway with a variety of FDA product codes. The FDA product codes that include these products are not limited to skin substitute products and may include other indications not related to wounds. **Note:** *The list of products named in this review is not a complete list of all commercially available products.*

Breast Reconstruction

For breast reconstruction surgery using one of the various acellular dermal matrix products used in breast reconstruction (i.e., including but not limited to the above-referenced medically necessary products) may be considered medically necessary when one of the following are met:

- When there is insufficient tissue expander or implant coverage by the pectoralis major muscle and additional coverage is required.
- When there is a viable but compromised or thin postmastectomy skin flaps that are at risk for dehiscence or necrosis.
- The inframammary fold and lateral mammary folds have been undermined during mastectomy and reestablishment of these landmarks is needed.

Diabetic Lower-Extremity Ulcer(s) or Venous Insufficiency

Required Documentation

Prior to the application of the skin substitute for diabetic or venous insufficiency ulcers, it is expected that the medical record documentation will contain evidence of the following:

- Documentation that conservative measures have failed, or the medical record supports that the wound is so clinically severe that it requires immediate, aggressive therapy.
- Documentation that the ulcer has failed to decrease in size and depth, or that there has been no change in baseline size or depth with no sign of improvement, or no indication that improvement is likely.
- Documentation of venous insufficiency either by history of deep venous thrombosis in the affected leg, or documentation of valvular reflux by duplex ultrasound, venography, or air/photo plethysmography
- The individual's medical record must reflect the frequency of skin substitute application, and this should be consistent with each individual's history and response to the skin substitute application and product safety criteria. Repeated application to the same wound within the same treatment period without signs of improvement may be denied.
- Documentation that the skin substitute product was appropriately applied and immobilized in accordance with the manufacturer's label instructions.

Grading System for Diabetic Ulcers

The following grading system for diabetic ulcers is frequently utilized:

Wagner Grading System

- A. Grade 1: Superficial Diabetic Ulcer
- B. Grade 2: Ulcer extension
 - 1. Involves ligament, tendon, joint capsule or fascia
 - 2. No abscess or Osteomyelitis
- C. Grade 3: Deep ulcer with abscess or Osteomyelitis (infection inferred)
- D. Grade 4: Gangrene to portion of forefoot (infection inferred)
- E. Grade 5: Extensive gangrene of foot (infection inferred)

Conventional Non-Surgical Therapy for Chronic Venous Insufficiency Ulcers and Chronic Diabetic Foot Ulcers include:

- Control of edema, venous hypertension, or lymphedema
- Control of any nidus of infection or colonization with bacterial or fungal elements
- Elimination of underlying cellulitis, osteomyelitis, foreign body, or malignant process
- Appropriate debridement of necrotic tissue or foreign body (exposed bone or tendon)
- For diabetic foot ulcers, appropriate non-weight bearing or off-loading pressure

- For venous stasis ulcers, compression therapy provided with documented diligent use of multilayer dressings, compression stockings of greater than 20mmHg pressure, or pneumatic compression
- Provision of wound environment to promote healing (protection from trauma and contaminants, elimination of inciting or aggravating processes)
- Management of concomitant and inciting medical issues (e.g., diabetes, tobacco use).

Failed Response - defined as an ulcer or skin deficit that has failed to respond to documented appropriate wound-care measures, has increased in size or depth, or has not changed in baseline size or depth and has no indication that improvement is likely (i.e., granulation, epithelialization, or progress towards closing)

Re-Application - refers to an additional application of skin substitute to the same ulcer within the same treatment period.

Retreatment - refers to a new treatment period where the same ulcer is being treated again because the initial treatment has failed.

Unsuccessful treatment - defined as increase in size or depth of an ulcer or no change in baseline size or depth and no sign of improvement or indication that improvement is likely (such as granulation, epithelialization, or progress towards closing) for a period of 2-4 weeks past start of therapy.

Coding

See the [Codes](#) table for details.

BACKGROUND

Skin and Soft Tissue Substitutes

Bioengineered skin and soft tissue substitutes may be either acellular or cellular. Acellular products (e.g., dermis with cellular material removed) contain a matrix or scaffold composed of materials such as collagen, hyaluronic acid, and fibronectin. Acellular dermal matrix (ADM) products can differ in a number of ways, including by species source (human, bovine, porcine), tissue source (e.g., dermis, pericardium, intestinal mucosa), additives (e.g., antibiotics, surfactants), hydration (wet, freeze-dried), and required preparation (multiple rinses, rehydration).

Cellular products contain living cells such as fibroblasts and keratinocytes within a matrix. The cells contained within the matrix may be autologous, allogeneic, or derived from other species (e.g., bovine, porcine). Skin substitutes may also be composed of dermal cells, epidermal cells, or a combination of dermal and epidermal cells, and may provide growth factors to stimulate healing. Bioengineered skin substitutes can be used as either temporary or permanent wound coverings.

Applications

There are a large number of potential applications for artificial skin and soft tissue products. One large category is nonhealing wounds, which potentially encompasses diabetic neuropathic ulcers, vascular insufficiency ulcers, and pressure ulcers. A substantial minority of such wounds do not heal adequately with standard wound care, leading to prolonged morbidity and increased risk of mortality. For example, nonhealing lower-extremity wounds represent an ongoing risk for infection, sepsis, limb amputation, and death. Bioengineered skin and soft tissue substitutes have the potential to improve rates of healing and reduce secondary complications.

Other situations in which bioengineered skin products might substitute for living skin grafts include certain postsurgical states (e.g., breast reconstruction) in which skin coverage is inadequate for the procedure performed, or for surgical wounds in patients with compromised ability to heal. Second- and third-degree burns are another indication in which artificial skin products may substitute for auto- or allografts. Certain primary dermatologic conditions that involve large areas of skin breakdown (e.g., bullous diseases) may also be conditions in which artificial skin products can be considered as substitutes for skin grafts. ADM products are also being evaluated in the repair of other soft tissues including rotator cuff repair, following oral and facial surgery, hernias, and other conditions.

Regulatory Status

The U.S. Food and Drug Administration (FDA) does not refer to any single product or class of products as “skin substitutes”. Products in this review cover products that do not require FDA approval or clearance as well as a number of products cleared through the 510(k) pathway with a variety of FDA product codes. A large number of artificial skin and soft-tissue products are commercially available or in development. Commercial availability is not a reflection of a product's regulatory status. The following section summarizes a subset of commercially available skin and soft-tissue substitutes. This is not a complete list of all commercially available products. Information on additional products is available in a 2020 Technical Brief on skin substitutes for treating chronic wounds that was commissioned by the Agency for Healthcare Research and Quality.

Acellular Dermal Matrix Products

Allograft ADM products derived from donated cadaveric human skin tissue are supplied by tissue banks compliant with standards of the American Association of Tissue Banks and U.S. Food and Drug Administration (FDA) guidelines. The processing removes the cellular components (i.e., epidermis, all viable dermal cells) that can lead to rejection and infection. ADM products from human skin tissue are regarded as minimally processed and not significantly changed in structure from the natural material; FDA classifies ADM products as banked human tissue and, therefore, not requiring FDA approval for homologous use.

In 2017, FDA published clarification of what is considered minimal manipulation and homologous use for human cells, tissues, and cellular and tissue-based products (HCT/Ps).

HCT/Ps are defined as human cells or tissues that are intended for implantation, transplantation, infusion, or transfer into a human recipient. If an HCT/P does not meet the criteria below and does not qualify for any of the stated exceptions, the HCT/P will be regulated as a drug, device, and/or biological product and applicable regulations and premarket review will be required.

An HCT/P is regulated solely under section 361 of the PHS Act and 21 CFR Part 1271 if it meets all the following criteria:

1. "The HCT/P is minimally manipulated;
2. The HCT/P is intended for homologous use only, as reflected by the labeling, advertising, or other indications of the manufacturer's objective intent;
3. The manufacture of the HCT/P does not involve the combination of the cells or tissues with another article, except for water, crystalloids, or a sterilizing, preserving, or storage agent, provided that the addition of water, crystalloids, or the sterilizing, preserving, or storage agent does not raise new clinical safety concerns with respect to the HCT/P; and
4. Either:

- i. The HCT/P does not have a systemic effect and is not dependent upon the metabolic activity of living cells for its primary function; or
 - ii. The HCT/P has a systemic effect or is dependent upon the metabolic activity of living cells for its primary function, and: a) Is for autologous use; b) Is for allogeneic use in a first-degree or second-degree blood relative; or c) Is for reproductive use."
- AlloDerm® (LifeCell Corp.) is an ADM (allograft) tissue-replacement product created from native human skin and processed so that the basement membrane and cellular matrix remain intact. Originally, AlloDerm® required refrigeration and rehydration before use. It is currently available in a ready-to-use product stored at room temperature. An injectable micronized form of AlloDerm® (Cymetra) is available.
 - Cortiva® (previously marketed as AlloMax™ Surgical Graft and before that NeoForm™) is an acellular non-cross-linked human dermis allograft.
 - AlloPatch® (Musculoskeletal Transplant Foundation) is an acellular human dermis allograft derived from the reticular layer of the dermis and marketed for wound care. This product is also marketed as FlexHD® for postmastectomy breast reconstruction.
 - FlexHD® and the newer formulation FlexHD® Pliable™ (Musculoskeletal Transplant Foundation) are acellular hydrated reticular dermis allograft derived from donated human skin.
 - DermACELL™ (LifeNet Health) is an allogeneic ADM processed with proprietary technologies MATRACELL® and PRESERVON®.
 - DermaMatrix™ (Synthes) is a freeze-dried ADM derived from donated human skin tissue. DermaMatrix Acellular Dermis is processed by the Musculoskeletal Transplant Foundation.
 - DermaPure™ (Tissue Regenix Wound Care) is a single layer decellularized human dermal allograft for the treatment of acute and chronic wounds.
 - GraftJacket® Regenerative Tissue Matrix (also called GraftJacket Skin Substitute; KCI) is an acellular regenerative tissue matrix that has been processed from human skin supplied from U.S. tissue banks. The allograft is minimally processed to remove the epidermal and dermal cells while preserving dermal structure. GraftJacket Xpress® is an injectable product.
 - mVASC® (MicroVascular Tissues, Inc.) is a microvascular tissue structural allograft made of small blood vessels and extracellular matrix, inherent non-viable cells, and associated biological signaling factors harvested from subcutaneous tissue of cadaveric human donors.

Although frequently used by surgeons for breast reconstruction, FDA does not consider this homologous use and has not cleared or approved any surgical mesh device (synthetic, animal collagen-derived, or human collagen-derived) for use in breast surgery. The indication of surgical mesh for general use in "Plastic and reconstructive surgery" was cleared by the FDA before surgical mesh was described for breast reconstruction in 2005. FDA states that the specific use of surgical mesh in breast procedures represents a new intended use and that a substantial equivalence evaluation via 510(k) review is not appropriate and a pre-market approval evaluation is required.

In March 2019, the FDA held an Advisory Committee meeting on breast implants, at which time the panel noted that while there is data about ADM for breast reconstruction, the FDA has not yet determined the safety and effectiveness of ADM use for breast reconstruction. The panel recommended that patients are informed and also recommended studies to assess the benefit and risk of ADM use in breast reconstruction.

In March 2021, FDA issued a Safety Communication to inform patients, caregivers, and health care providers that certain ADM products used in implant-based breast reconstruction may have a higher chance for complications or problems. An FDA analysis of patient-level data from real-world use of ADMs

for implant-based breast reconstruction suggested that 2 ADMs—FlexHD and Allomax—may have a higher risk profile than others.

In October 2021, an FDA advisory panel on general and plastic surgery voted against recommending FDA approval of the SurgiMend mesh for the specific indication of breast reconstruction. The advisory panel concluded that the benefits of using the device did not outweigh the risks.

FDA product codes: FTM, OXF.

Xenogeneic Products

Cytal™ (previously called MatriStem®) Wound Matrix, Multilayer Wound Matrix, Pelvic Floor Matrix, MicroMatrix, and Burn Matrix (all manufactured by ACell) are composed of porcine-derived urinary bladder matrix.

Helicoll (Encol) is an acellular collagen matrix derived from bovine dermis. In 2004, it was cleared for marketing by the FDA through the 510(k) process for topical wound management that includes partial and full-thickness wounds, pressure ulcers, venous ulcers, chronic vascular ulcers, diabetic ulcers, trauma wounds (e.g., abrasions, lacerations, second-degree burns, skin tears), and surgical wounds including donor sites/grafts.

Keramatrix® (Keraplast Research) is an open-cell foam comprised of freeze-dried keratin that is derived from acellular animal protein. In 2009, it was cleared for marketing by the FDA through the 510(k) process under the name of Keratec. The wound dressings are indicated in the management of the following types of dry, light, and moderately exuding partial and full-thickness wounds: pressure (stage I-IV) and venous stasis ulcers, ulcers caused by mixed vascular etiologies, diabetic ulcers, donor sites, and grafts.

Kerecis™ Omega3 Wound (Kerecis) is an ADM derived from fish skin. It has a high content of omega 3 fatty acids and is intended for use in burn wounds, chronic wounds, and other applications.

Permacol™ (Covidien) is xenogeneic and composed of cross-linked porcine dermal collagen. Cross-linking improves tensile strength and long-term durability but decreases pliability.

PriMatrix™ (TEI Biosciences; a subsidiary of Integra Life Sciences) is a xenogeneic ADM processed from fetal bovine dermis. It was cleared for marketing by the FDA through the 510(k) process for partial- and full-thickness wounds; diabetic, pressure, and venous stasis ulcers; surgical wounds; and tunneling, draining, and traumatic wounds. FDA product code: KGN.

SurgiMend® PRS (TEI Biosciences, a subsidiary of Integra Life Sciences) is a xenogeneic ADM processed from fetal and neonatal bovine dermis.

Strattice™ Reconstructive Tissue Matrix (LifeCell Corp.) is a xenogeneic non-cross-linked porcine-derived ADM. There are pliable and firm versions, which are stored at room temperature and come fully hydrated.

Oasis™ Wound Matrix (Cook Biotech) is a collagen scaffold (extracellular matrix) derived from porcine small intestinal submucosa. In 2000, it was cleared for marketing by the FDA through the 510(k) process for the management of partial- and full-thickness wounds, including pressure ulcers, venous ulcers, diabetic ulcers, chronic vascular ulcers, tunneled undermined wounds, surgical wounds, trauma wounds, and draining wounds.

OviTex® (TELA Bio Inc.) is a next generation soft tissue repair platform that utilizes layers of ovine (sheep) rumen interwoven with just enough polymer suture for added strength (2014). OviTex PRS long term resorbable received FDA approval (2023) for one time use to reinforce soft tissue where weakness exists in individuals requiring soft tissue repair or reinforcement in plastic and reconstructive surgery. In 2024, the FDA approval was expanded to include OviTex PRS (STR) (or OviTex PRS Short Term Resorbable) and OviTex PRS PP (or OviTex PRS Permanent).

FDA Product code: KGN, FTL, FTM.

Living Cell Therapy

Apligraf® (Organogenesis) is a bilayered living cell therapy composed of an epidermal layer of living human keratinocytes and a dermal layer of living human fibroblasts. Apligraf® is supplied as needed, in 1 size, with a shelf-life of 10 days. In 1998, it was approved by the FDA for use in conjunction with compression therapy for the treatment of noninfected, partial- and full-thickness skin ulcers due to venous insufficiency and in 2001 for full-thickness neuropathic diabetic lower-extremity ulcers nonresponsive to standard wound therapy.

Dermagraft® (Organogenesis) is composed of cryopreserved human-derived fibroblasts and collagen derived from newborn human foreskin and cultured on a bioabsorbable polyglactin mesh scaffold. Dermagraft has been approved by the FDA for repair of diabetic foot ulcers.

TheraSkin® (Soluble Systems) is a cryopreserved split-thickness human skin allograft composed of living fibroblasts and keratinocytes and an extracellular matrix in epidermal and dermal layers. TheraSkin® is derived from human skin allograft supplied by tissue banks compliant with the American Association of Tissue Banks and FDA guidelines. It is considered a minimally processed human cell, tissue, and cellular- and tissue-based product by the FDA.

Epicel® (Genzyme Biosurgery) is an epithelial autograft composed of a patient's own keratinocytes cultured ex vivo and is FDA-approved under a humanitarian device exemption for the treatment of deep dermal or full-thickness burns comprising a total body surface area of 30% or more. It may be used in conjunction with split-thickness autografts or alone in patients for whom split-thickness autografts may not be an option due to the severity and extent of their burns.

OrCel™ (Forticell Bioscience; formerly Composite Cultured Skin) is an absorbable allogeneic bilayered cellular matrix, made of bovine collagen, in which human dermal cells have been cultured. It was approved by FDA premarket approval for healing donor site wounds in burn victims and under a humanitarian device exemption for use in patients with recessive dystrophic epidermolysis bullosa undergoing hand reconstruction surgery to close and heal wounds created by the surgery, including those at donor sites.

FDA product codes: FTM, PFC, OCE, ODS.

Autologous Cell Harvesting Device

Recell® (Avita Medical) was initially approved by the FDA in September 2018 under the PMA process (PMA BP170122). It is an autologous cell harvesting device indicated for the treatment of acute partial-thickness thermal burn wound when used by an appropriately licensed healthcare professional at the patient's point of care to prepare autologous RES Regenerative Epidermal Suspension. The initial indication was for use in patients 18 years of age and older in combination with meshed autografting.

Subsequently, indications were expanded to include direct application to acute partial-thickness thermal burn wounds in patients 18 years of age and older or application in combination with meshed autografting for acute full-thickness thermal burn wounds in pediatric as well as adult patients.

FDA product code: QCZ.

Biosynthetic Products

Biobrane®/Biobrane-L (Smith & Nephew) is a biosynthetic wound dressing constructed of a silicon film with a nylon fabric partially embedded into the film. The fabric creates a complex 3-dimensional structure of trifilament thread, which chemically binds collagen. Blood/sera clot in the nylon matrix, adhering the dressing to the wound until epithelialization occurs.

Integra® Dermal Regeneration Template (also marketed as Omnigraft Dermal Regeneration Matrix; Integra LifeSciences) is a bovine, collagen/glycosaminoglycan dermal replacement covered by a silicone temporary epidermal substitute. It was approved by the FDA for use in the post-excisional treatment of life-threatening full-thickness or deep partial-thickness thermal injury where sufficient autograft is not available at the time of excision or not desirable because of the physiologic condition of the patient, and for certain diabetic foot ulcers. Integra® Matrix Wound Dressing and Integra® Meshed Bilayer Wound Matrix are substantially equivalent skin substitutes and were cleared for marketing by the FDA through the 510(k) process for other indications. Integra® Bilayer Matrix Wound Dressing (Integra LifeSciences) is designed to be used in conjunction with negative pressure wound therapy. The meshed bilayer provides a flexible wound covering and allows drainage of wound exudate.

TransCyte™ (Advanced Tissue Sciences) consists of human dermal fibroblasts grown on nylon mesh, combined with a synthetic epidermal layer, and was approved by the FDA in 1997. TransCyte is intended as a temporary covering over burns until autografting is possible. It can also be used as a temporary covering for some burn wounds that heal without autografting.

FDA product codes: FRO, MDD, MGR.

Synthetic Products

Suprathel® (PolyMedics Innovations) is a synthetic copolymer membrane fabricated from a tripolymer of polylactide, trimethylene carbonate, and s-caprolactone. It is used to provide temporary coverage of superficial dermal burns and wounds. Suprathel® is covered with gauze and a dressing that is left in place until the wound has healed.

Nerve Wraps

Nerve wraps can be used for peripheral nerve repair. They are often made from biocompatible materials like collagen, designed to encase injured peripheral nerves. It provides a barrier between the nerve and surrounding tissue, minimizing scarring and promoting a conducive environment for nerve healing. Their application is ideal for cases where the nerve is intact, but needs protection from scarring or compression. Nerve conduits, addressed in evidence review 7.01.177, would be more suitable in cases of severed nerves.

AxoGuard® nerve connector (Axogen, Inc) is an implant derived from small intestine submucosa designed to protect injured and compressed nerves. Other FDA 510K approved nerve wraps include: Flexibile Collagen Nerve Cuff (Collagen Matrix, Inc), Mochida Nerve Cuff (Mochida Pharmaceutical Co.),

NervAlign Nerve Cuff (Renerve, Ltd), Nerve tape (BioCircuit Technologies, Inc), Neurawrap (Integra LifeSciences, Corp), NeuroMend (Stryker Orthopedics), NeuroShield (Monarch bioimplants, GmbH), Reinforce flexible Collagen Nerve Cuff (Collagen Matrix, Inc), and Versawrap nerve protector (Alafair Biosciences, Inc).

FDA product code: JXI.

RATIONALE

This evidence review was created in May 1999 and has been updated regularly with searches of the PubMed database. The most recent literature update was performed through May 2026.

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life (QOL), and ability to function, including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

There is no standard definition of “skin substitute”. Products reviewed in the following sections include products that do not require U.S. Food and Drug Administration (FDA) approval or clearance as well as a number of products cleared through the 510(k) pathway with a variety of FDA product codes. The FDA product codes that include these products are not limited to skin substitute products and may include other indications not related to wound healing or wound care.

Breast Reconstruction

Clinical Context and Therapy Purpose

A variety of breast reconstruction techniques are used postmastectomy, including implant-based (immediate or delayed following use of a tissue expander) and those using autologous tissue flaps. Some of these techniques have been used with acellular dermal matrix (ADM) to provide additional support or tissue coverage. The purpose of bioengineered soft tissue substitutes in individuals who are undergoing breast reconstruction (prophylactic or oncologic indications) is to provide a treatment option that is an alternative to or an improvement on breast reconstruction without use of a biological or biosynthetic matrix.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals who are undergoing breast reconstruction, typically following mastectomy (prophylactic or oncologic indications).

Interventions

The therapy being considered is bioengineered soft tissue substitutes as a biological matrix that is used to facilitate one-stage tissue expander reconstruction. As noted in the regulatory status section, the U.S. Food and Drug Administration (FDA) has not cleared or approved any surgical mesh device (synthetic, animal collagen-derived, or human collagen-derived) for use in breast surgery. In October 2021, an FDA advisory panel on general and plastic surgery voted against recommending FDA approval of the SurgiMend mesh for the specific indication of breast reconstruction. The advisory panel concluded that the benefits of using the device did not outweigh the risks.

Comparators

The following therapies are currently being used to make decisions about soft tissue substitutes or biological matrices: 2-stage tissue expander reconstruction without a biological matrix.

Outcomes

The general outcomes of interest are symptoms, morbid events, functional outcomes, QOL, and treatment-related morbidity. Specific outcomes are the time to permanent implant, pain during and after the procedure, and adverse events including seroma, infection, and necrosis rates, rates of capsular contracture, and malposition of implants. Short-term outcomes would be measured within 3 months with longer-term outcomes apparent by 2 years.

Study Selection Criteria

- To assess efficacy outcomes, we sought comparative controlled prospective trials, with preference for RCTs*.
- In the absence of such trials, we sought comparative observational studies, with preference for prospective studies.
- To assess longer-term outcomes and adverse effects, we sought single-arm studies that capture longer periods of follow-up and/or larger populations.
- Within each category of study design, we prefer larger sample size studies and longer duration studies.
- We excluded studies with duplicative or overlapping populations.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials.

Review of Evidence

The literature on ADM for breast reconstruction consists primarily of retrospective, uncontrolled series and systematic reviews of these studies.

A 2013 study used data from the American College of Surgeon's National Surgical Quality Improvement Program to compare ADM-assisted tissue expander breast reconstruction (n=1717) to submuscular tissue expander breast reconstruction (n=7442) after mastectomy. Complication rates did not differ

significantly between the ADM-assisted (5.5%) and the submuscular tissue expander groups (5.3%; $p=.68$). Rates of reconstruction-related complications, major complications, and 30-day reoperation did not differ significantly between cohorts.

Systematic Reviews

Ng et al (2024) conducted a systematic review and meta-analysis comparing postoperative complications and patient-reported outcomes between patients who received ADM and those who did not. Prospective cohort studies and RCTs were included (9 studies; $N=3161$). There were no significant differences in postoperative outcomes between the ADM and non-ADM groups for key complications such as seroma ($p=.51$), hematomas ($p=.20$), infections ($p=.21$), wound dehiscence ($p=.09$), reoperations ($p=.70$), implant loss ($p=.27$), or skin necrosis ($p=.21$).

A meta-analysis by Lee and Mun (2016) included 23 studies (total $N=6199$ cases) on implant-based breast reconstruction that were published between February 2011 and December 2014. The analysis included an RCT and 3 prospective comparative cohort studies; the remainder was retrospective comparative cohort studies. Use of ADM did not affect the total complication rate (see Table 1). ADM significantly increased the risk of major infection, seroma, and flap necrosis, but reduced risks of capsular contracture and implant malposition. Use of ADM allowed for significantly greater intraoperative expansion (mean difference, 79.63; 95% confidence interval [CI], 41.99 to 117.26; $p<.001$) and percentage of intraoperative filling (mean difference, 13.30; 95% CI, 9.95 to 16.65; $p<.001$), and reduced the frequency of injections to complete expansion (mean difference, -1.56; 95% CI, -2.77 to -0.35; $p=.01$).

Table 1. Meta-Analysis of Breast Reconstruction Outcomes With and Without ADM

Outcome Measure	Relative Risk	95% Confidence Interval	p
Infection	1.42	1.02 to 1.99	.04
Seroma	1.41	1.12 to 1.78	.004
Mastectomy flap necrosis	1.44	1.11 to 1.87	.006
Unplanned return to the operating room	1.09	0.63 to 1.90	NS
Implant loss	1.00	0.68 to 1.48	NS
Total complications	1.08	0.87 to 1.34	NS
Capsular contracture	0.26	0.15 to 0.47	<.001
Implant malposition	0.21	0.07 to 0.59	.003

Adapted from Lee and Mun (2016).⁶
ADM: acellular dermal matrix; NS: not significant.

AlloDerm

Randomized Controlled Trials

McCarthy et al (2012) reported on a multicenter, blinded RCT of AlloDerm in 2-stage expander/implant reconstruction. Seventy patients were randomized to AlloDerm ADM-assisted tissue expander/implant

reconstruction or to submuscular tissue expander/implant placement. The trial was adequately powered to detect clinically significant differences in immediate postoperative pain but underpowered to detect the secondary endpoint of pain during tissue expansion. There were no significant differences between the groups in the primary outcomes of immediate postoperative pain (54.6 AlloDerm vs. 42.8 controls on a 100-point visual analog scale) or pain during the expansion phase (17.0 AlloDerm vs. 4.6 controls) or in the secondary outcome of rate of tissue expansion (91 days AlloDerm vs. 108 days controls) and patient-reported physical well-being. There was no significant difference in adverse events, although the total number of adverse events was small.

Comparison Between Products

AlloDerm Versus AlloMax

Hinchcliff et al (2017) conducted an RCT that compared AlloDerm with AlloMax (n=15 each) for implant-based breast reconstruction. Complications were assessed 7, 14, and 30 days postoperatively and biopsies of the ADMs were taken during implant exchange. Vessel density in the AlloMax biopsies was higher than in the AlloDerm biopsies. Complications were reported in 26.1% of AlloMax cases and 8.0% of AlloDerm cases; these complication rates did not differ statistically with the 30 patients in this trial.

AlloDerm Versus DermaMatrix

Mendenhall et al (2017) conducted an RCT that compared AlloDerm with DermaMatrix in 111 patients (173 breasts). There were no significant differences in overall rates of complications (AlloDerm, 15.4%; DermaMatrix, 18.3%; $p=.8$) or implant loss (AlloDerm, 2.2%; DermaMatrix, 3.7%; $p=.5$) between the 2 ADMs at 3 months postoperative. There were no statistically difference in the overall complication rates (6% vs 13%; $p=.3$), severity of complications, or patient satisfaction between the AlloDerm and DermMatrix groups at 2 years after definitive reconstruction.

AlloDerm Versus DermACELL

Davison et al (2024) conducted a prospective randomized trial comparing AlloDerm with DermACELL in 55 patients undergoing bilateral nipple and/or skin-sparing mastectomies. Patients served as their own controls and were blinded to the random assignment of the two products to the left or right breast. The findings revealed no significant differences in drain removal time or average drain output between the two groups. However, a notable difference was observed in seroma rates, with 30.91% of AlloDerm breasts experiencing seromas compared to 14.55% in DermACELL breasts ($p<.05$). Additionally, incorporation rates were significantly higher for DermACELL at 99.8% compared to AlloDerm's 93.4% ($p<.05$). Both AlloDerm and DermACELL demonstrated a high success rate of 94.55% for reconstruction outcomes. Nonetheless, AlloDerm was associated with a higher incidence of seromas and a trend towards lower incorporation rates.

AlloDerm Versus Cortiva

Keane et al (2024) conducted an RCT comparing Cortiva with AlloDerm in patients who underwent either direct-to-implant (DTI) or tissue expander (TE) reconstruction (N=302). The primary outcome measured was reconstructive failure, defined as premature explantation of TEs or DTI reconstructions before 3 months postoperatively. 151 patients received AlloDerm (280 breasts) and 151 received Cortiva (277 breasts). The results showed no significant difference in reconstructive failure rates between the two ADMs, with AlloDerm at 9.3% and Cortiva at 8.3% ($p=.68$). Additionally, there were no notable differences in other complications or patient-reported outcomes between the groups. Seroma formation was more

prevalent in the AlloDerm group (12%) compared to Cortiva (7.6%) and was statistically significant (odds ratio: 1.93; 95% CI: 1.01 to 3.67; p=.047).

Strattice

Dikmans et al (2017) reported on early safety outcomes from an open-label multicenter RCT that compared porcine ADM-assisted 1-stage expansion with 2-stage implant-based breast reconstruction (see Table 2). One-stage breast reconstruction with porcine ADM was associated with a higher risk of surgical complications, reoperation, and with removal of implant, ADM, or both (see Table 3). The trial was stopped early due to safety concerns, but it cannot be determined from this study design whether the increase in complications was due to the use of the xenogeneic ADM or to the comparison between 1-stage and 2-stage reconstruction.

Table 2. Summary of Key RCT Characteristics

Author	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Dikmans et al (2017)	EU	8	2013-2015	Women intending to undergo skin-sparing mastectomy and immediate IBBR	59 patients (91 breasts) undergoing 1-stage IBBR with ADM	62 women (92 breasts) undergoing 2-stage IBBR

ADM: acellular dermal matrix; EU: European Union; IBBR: implant-based breast reconstruction; RCT: randomized controlled trial.

Table 3. Summary of Key RCT Outcomes

Study	Surgical Complications	Severe Adverse Events	Reoperation	Removal of Implant, ADM, or Both
Dikmans et al (2017)				
1-stage with ADM, n (%)	27 (46)	26 (29)	22 (37)	24 (26)
2-stage with ADM, n (%)	11 (18)	5 (5)	9 (15)	4 (5)
OR (95% CI)	3.81 (2.67 to 5.43)		3.38 (2.10 to 5.45)	8.80 (8.24 to 9.40)
p	<.001		<.001	<.001

ADM: acellular dermal matrix; CI: confidence interval; OR: odds ratio; RCT: randomized controlled trial.

Section Summary: Breast Reconstruction

Results of a systematic review found no difference in overall complication rates between ADM allograft and standard procedures for breast reconstruction. Although reconstructions with ADM have been

reported to have higher seroma, infection, and necrosis rates than reconstructions without ADM, rates of capsular contracture and malposition of implants may be reduced. Thus, in cases where there is limited tissue coverage, the available studies may be considered sufficient to permit informed decision-making about risks and benefits of using allogeneic ADM for breast reconstruction.

Tendon Repair

Clinical Context and Therapy Purpose

The purpose of bioengineered soft tissue substitutes in individuals who are undergoing tendon repair is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals undergoing tendon repair.

Interventions

The therapy being considered is bioengineered soft-tissue substitutes.

Comparators

The following therapies are currently being used to make decisions about tendon repair: tendon repair without bioengineered soft-tissue substitutes.

Outcomes

The general outcomes of interest are symptoms, morbid events, functional outcomes, QOL, and treatment-related morbidity. Short-term outcomes would be measured within 3 months with longer-term outcomes apparent by 2 years.

Study Selection Criteria

- To assess efficacy outcomes, we sought comparative controlled prospective trials, with preference for RCTs*.
- In the absence of such trials, we sought comparative observational studies, with preference for prospective studies.
- To assess longer-term outcomes and adverse effects, we sought single-arm studies that capture longer periods of follow-up and/or larger populations.
- Within each category of study design, prefer larger sample size studies and longer duration studies.
- We excluded studies with duplicative or overlapping populations.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials.

Review of Evidence

GraftJacket

Barber et al (2012) reported an industry-sponsored multicenter RCT of augmentation with GraftJacket human ADM for arthroscopic repair of large (>3 cm) rotator cuff tears involving 2 tendons. Twenty-two patients were randomized to GraftJacket augmentation and 20 patients to no augmentation. At a mean follow-up of 24 months (range, 12-38 months), the American Shoulder and Elbow Surgeons score improved from 48.5 to 98.9 in the GraftJacket group and from 46.0 to 94.8 in the control group ($p=.035$). The Constant score improved from 41 to 91.9 in the GraftJacket group and from 45.8 to 85.3 in the control group ($p=.008$). The University of California, Los Angeles score did not differ significantly between groups. Gadolinium-enhanced magnetic resonance imaging (MRI) scans showed intact cuffs in 85% of repairs in the GraftJacket group and 40% of repairs in the control group. However, no correlation was found between MRI findings and clinical outcomes. Rotator cuff retears occurred in 3 (14%) patients in the GraftJacket group and 9 (45%) patients in the control group.

Rashid et al (2020) reported disruption of the native extracellular matrix with either GraftJacket or Permacol (porcine acellular dermis) as a patch overlay for rotator cuff repair in a small controlled study with 13 patients. The disruption was greater in the Permacol group and there was an immune response in 1 of 3 patients following use of the xenograft.

Section Summary: Tendon Repair

One small RCT was identified that found improved outcomes with GraftJacket ADM allograft for rotator cuff repair. Although results of this trial were promising, additional study with a larger number of patients is needed to corroborate these findings and determine the effects of this technology with greater certainty.

Surgical Repair of Hernias or Parastomal Reinforcement

Clinical Context and Therapy Purpose

The purpose of bioengineered soft tissue substitutes in individuals who are undergoing surgical repair of hernias or require parastomal reinforcement is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals undergoing surgical repair of hernias or requiring parastomal reinforcement.

Interventions

The therapy being considered is bioengineered matrix support.

Comparators

The following therapies are currently being used for surgical repair of hernias or parastomal reinforcement: synthetic mesh.

Outcomes

The general outcomes of interest are symptoms, morbid events, functional outcomes, QOL, and treatment-related morbidity. Specific outcomes are surgical site occurrence of postoperative infection, seroma/hematoma, pain, bulging, dehiscence, fistula, or mechanical failure. Short-term outcomes would be measured within 3 months with longer-term outcomes apparent by 2 years.

Study Selection Criteria

- To assess efficacy outcomes, we sought comparative controlled prospective trials, with preference for RCTs*.
- In the absence of such trials, we sought comparative observational studies, with preference for prospective studies.
- To assess longer-term outcomes and adverse effects, we sought single-arm studies that capture longer periods of follow-up and/or larger populations.
- Within each category of study design, prefer larger sample size studies and longer duration studies.
- We excluded studies with duplicative or overlapping populations.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials.

Review of Evidence

Systematic Reviews

A 2013 systematic review evaluated the clinical effectiveness of acellular collagen-based scaffolds for the repair of incisional hernias. The bioprosthetic materials could be harvested from bovine pericardium, human cadaveric dermis, porcine small intestine mucosa, porcine dermal collagen, or bovine dermal collagen. Products included in the search were Surgisis, Tutomesh, Veritas, AlloDerm, FlexHD, AlloMax, CollaMend, Permacol, Strattice, FortaGen, ACell, DermaMatrix, XenMatrix, and SurgiMend. Sixty publications with 1212 repairs were identified and included in the review, although meta-analysis could not be performed. There were 4 level III studies (2 AlloDerm, 2 Permacol); the remainder were level IV or V. The largest number of publications were on AlloDerm (n=27) and Permacol (n=18). No publications on incisional hernia repair were identified for AlloMax, FortaGen, DermaMatrix, or ACell. The overall incidence of a surgical site occurrence (e.g., postoperative infection, seroma/hematoma, pain, bulging, dehiscence, fistula, mechanical failure) was 82.6% for porcine small intestine mucosa, 50.7% for xenogeneic dermis, 48.3% for human dermis, and 6.3% for xenogeneic pericardium. No comparative data were identified that could establish superiority to permanent synthetic meshes.

AlloDerm as an Overlay

Espinosa-de-los-Monteros et al (2007) retrospectively reviewed 39 abdominal wall reconstructions with AlloDerm performed in 37 patients and compared them with 39 randomly selected cases. They reported a significant decrease in recurrence rates when human cadaveric acellular dermis was added as an overlay to primary closure plus rectus muscle advancement and imbrication in patients with medium-sized hernias. However, no differences were observed when adding human cadaveric acellular dermis as an overlay to patients with large-size hernias treated with underlay mesh.

Comparison Between Products

AlloDerm Versus Surgisis Gold

Gupta et al (2006) compared the efficacy and complications associated with use of AlloDerm and Surgisis bioactive mesh in 74 patients who underwent ventral hernia repair. The first 41 procedures were performed using Surgisis Gold 8-ply mesh formed from porcine small intestine submucosa, and the remaining 33 patients had ventral hernia repair with AlloDerm. Patients were seen 7 to 10 days after discharge from the hospital and at 6 weeks. Any signs of wound infection, diastasis, hernia recurrence, changes in bowel habits, and seroma formation were evaluated. The use of the AlloDerm mesh resulted in 8 (24%) hernia recurrences. Fifteen (45%) of the AlloDerm patients developed a diastasis or bulging at the repair site. Seroma formation was only a problem in 2 patients.

AlloDerm Versus FlexHD

A 2013 study compared AlloDerm with FlexHD for complicated hernia surgery. From 2005 to 2007, AlloDerm was used to repair large (>200 cm²) symptomatic complicated ventral hernias that resulted from trauma or emergency surgery (n=55). From 2008 to 2010, FlexHD was used to repair large, complicated ventral hernias in patients meeting the same criteria (n=40). The 2 groups were comparable at baseline. At 1 year follow-up, all AlloDerm patients were diagnosed with hernia recurrence (abdominal laxity, functional recurrence, true recurrence) requiring a second repair. Eleven (31%) patients in the FlexHD group required a second repair. This comparative study is limited by the use of nonconcurrent comparisons, which is prone to selection bias and does not control for temporal trends in outcomes.

FlexHD Versus Strattice

Roth et al (2017) reported on a prospective study assessing clinical and QOL outcomes following complex hernia repair with a human (FlexHD) or porcine (Strattice) ADM. The study was funded by the Musculoskeletal Transplant Foundation, which prepares and supplies FlexHD. Patients were enrolled if they had a hernia at least 6 cm in the transverse dimension, active or prior infection of the abdominal wall, and/or enterocutaneous fistula requiring mesh removal. Eighteen (51%) of the 35 patients had undergone a previous hernia repair. After abdominal wall repair with the ADM, 20 (57%) patients had a surgical site occurrence, and nearly one-third had hospital readmission. The type of biologic material did not impact hernia outcomes. There was no comparison with synthetic mesh in this study, limiting interpretation.

Strattice Versus Synthetic Mesh

Bellows et al (2014) reported early results of an industry-sponsored multicenter RCT that compared Strattice (non-cross-linked porcine ADM, n=84) with a standard synthetic mesh (n=88) for the repair of inguinal hernias. The trial was designed by the surgeons and was patient- and assessor-blinded to reduce risk of bias. Blinding continued through 2 years of follow-up. The primary outcome was resumption of activities of daily living at 1 year. Secondary outcomes included complications, recurrences, or chronic pain (i.e., pain that did not disappear by 3 months post-surgery). At 3-month follow-up, there were no significant differences in either the occurrence or type of wound events (relative risk, 0.98; 95% CI, 0.52 to 1.86). Pain was reduced from 1 to 3 days postoperative in the group treated with Strattice, but at 3-month follow-up pain scores did not differ significantly between groups.

Dhanani et al (2025) reported 3-year outcomes of a single-center pilot RCT comparing biologic mesh (non-cross-linked porcine acellular dermal matrix) with synthetic mesh (medium-density, macroporous

polypropylene) for complex open ventral hernia repair. Of the 87 patients randomized (44 biologic, 43 synthetic), 61 (70%) completed 3-year follow-up. No significant differences were observed between biologic and synthetic mesh for major complications (50% vs 30%; RR, 2.34; 95% CI, 0.80 to 6.90; $p=.123$), hernia recurrence (39% vs 24%; $p=.214$), mesh infection (14% vs 3%; $p=.144$), or reoperation (18% vs 12%; $p=.531$). Both groups demonstrated clinically meaningful improvements in functional status and pain scores at 3 years. The study was limited by its small sample size and 30% loss to follow-up, and was underpowered to detect significant differences in complication rates.

Seefeldt et al (2025) reported the BIOLAP multicenter, randomized, self-controlled clinical trial comparing biological versus synthetic mesh in laparoendoscopic inguinal hernia repair. A total of 491 patients with primary bilateral inguinal hernias were enrolled across 21 certified German hernia centers; each patient received a non-cross-linked, acellular, collagenous biological mesh on one side and a lightweight synthetic mesh on the other, with randomization determining side assignment. The biological meshes came from 3 different manufacturers, and both patients and follow-up clinicians were blinded to mesh allocation. At 6 months, there was no significant difference in pain between biological and synthetic mesh (mean VAS score at rest, 0.3 for both; $p=.76$). At 2 years, the recurrence rate was significantly higher for biological meshes (11.2%) than for synthetic meshes (2.5%; $p<.001$), and seroma rates were also significantly higher with biological mesh (33.4% vs 21.6%; $p<.001$). On multivariate analysis, biological mesh was the strongest predictor of recurrence (OR, 4.53; 95% CI, 2.42 to 8.49; $p<.001$).

Remulla et al (2026) reported the 5-to-10-year follow-up of a multicenter RCT (NCT02451176) comparing biologic mesh (Strattice Reconstructive Tissue Matrix, a non-cross-linked porcine ADM) with synthetic mesh (Bard Soft Mesh, medium-weight polypropylene) for single-stage retromuscular repair of clean-contaminated and contaminated ventral hernias. Of 253 patients originally randomized, 203 (80.2%) achieved minimum 5-year follow-up (median, 5.4 years). Synthetic mesh was associated with a significant reduction in midline hernia recurrence risk (HR, 0.46; 95% CI, 0.25 to 0.86; $p=.015$). The overall 5-to-10-year midline recurrence rate was 17.8% across both groups: 23.6% in the biologic group versus 11.8% in the synthetic group, corresponding to an absolute risk reduction of 11.8% (95% CI, 2.1 to 21.4) with synthetic mesh. This advantage was primarily observed within the first 2 years; the difference in recurrence rates beyond 2 years was not statistically significant. No new mesh infections or mesh excisions occurred in either group beyond 2 years postoperatively. Three patients (1.2%) required intervention for ongoing wound-related issues.

Strattice Versus No Reinforcement

A retrospective case-control study of 80 subjects undergoing ventral hernia repair with either Strattice ($n=40$) or conventional open repair ($n=40$) was reported by Richmond (2014). Mean follow-up was 33.1 months. The authors reported that the defect size was greater in the Strattice group (mean, 372.5 vs. 283.7 cm², $p=0.01$) as was the percentage Ventral Hernia Working Group Grade III/IV hernias (65.0% vs. 30.0%, $p=0.03$). Despite this, the number of recurrences were lower in the Strattice group (13.2% vs. 37.5%, $p=0.02$), and infection rates were lower as well (0% vs. 23%, respectively, $p=0.002$). Finally, the indications for reoperation, including recurrence or complications requiring reoperation, were also lower in the Strattice group (17.5% vs. 52.5%, $p=0.002$).

Also in 2014, the Parastomal Reinforcement with Strattice (PRISM) Study Group reported a multicenter, double-blinded, randomized trial of Strattice for parastomal reinforcement in patients undergoing surgery for permanent abdominal wall ostomies. Patients were randomized to standard stoma construction with no reinforcement ($n=58$) or stoma construction with Strattice as parastomal reinforcement ($n=55$). At 24-month follow-up ($n=75$), the incidence of parastomal hernias was similar for the 2 groups (13.2% of controls, 12.2% of study group).

Adverse Events

Permacol (porcine acellular dermal matrix) was reported in a case series of 13 patients to result in recurrent intestinal fistulation and intestinal failure when used for abdominal reconstructive surgery.

OviTex

OviTex is a decellularized ovine forestomach extracellular matrix mesh reinforced with five percent polymer fiber. OviTex has both permanent and resorbable variations.

Sivarajah and colleagues (2022) reported a retrospective analysis of 109 subjects, 18 years of age and older, from 2002-2021 who underwent ventral hernia repair to assess the association of mesh type with complications and surgical site occurrence. Ventral hernia repair was performed with OviTex reinforced biologic ovine rumen (n=50) or synthetic polypropylene mesh (n=59). Synthetic products used included: Prolene (n=29), Parietex (n=15), and Physiomesher (n=15). All subjects with OviTex had prior abdominal surgery compared to 86.4% of subjects who received synthetic mesh (p=0.01). Subjects who received OviTex had lower rates of adverse events (16.0 vs 30.5%, p=0.12), and similar hernia recurrence rates (4.0 vs. 6.78%, p=0.68) compared to subjects with synthetic mesh. Synthetic mesh was associated with increased odds for overall complications (3.78, p<0.05) and adverse events (3.87, p<0.05). The authors concluded that OviTex had a superior profile to other mesh products due to comparable hernia recurrences and decreased rate of adverse events.

The same investigators (Sivaraj 2022) also reported a retrospective analysis of 141 subjects who underwent ventral hernia repair with varying biologic mesh types which compared postoperative donor site complications and hernia recurrence rates amongst 1) Strattice (n=51), 2) Permacol (n=17), 3) OviTex (n=36), and 4) Surgimend (n=37). There were less complications in subjects with OviTex (16.7%), compared to Permacol (52.9%), Strattice (47.1%), and Surgimend (43.2%) (p=0.015 for OviTex vs. all other products). Rates of hernia recurrence were also lower in subjects that received OviTex (2.78%) and Strattice (13.7%) compared to Permacol (29.4%) and Surgimend (24.3%) (p=0.022). These results reiterated the authors' previous findings that OviTex decreased both abdominal complications and recurrence rates after ventral hernia repair compared to the other products.

OviTex is supported by sufficient evidence showing that it is at least commensurate with other accepted approaches for hernia repair and may provide a lower risk of adverse events.

Section Summary: Surgical Repair of Hernias or Parastomal Reinforcement

Current evidence does not support a benefit of ADMs in hernia repair or prevention of parastomal hernia. Additional RCTs are needed to compare biologic mesh with synthetic mesh and to determine if there is a patient population that would benefit from these products.

Diabetic Lower-Extremity Ulcers

Clinical Context and Therapy Purpose

The purpose of bioengineered soft tissue substitutes in individuals who have diabetic lower extremity ulcers is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with diabetic lower extremity ulcers.

Interventions

The therapy being considered is bioengineered skin substitutes.

Comparators

The following therapies are currently being used: standard wound care which involves regular debridement and moist wound covering.

Outcomes

The general outcomes of interest are symptoms, change in disease status, morbid events, and QOL.

The primary endpoints of interest for trials of wound closure are as follows, consistent with guidance from the FDA for industry in developing products for treatment of chronic cutaneous ulcer and burn wounds:

- Incidence of complete wound closure.
- Time to complete wound closure (reflecting accelerated wound closure).
- Incidence of complete wound closure following surgical wound closure.
- Pain control.

Time to wound closure can be measured at 12 weeks and 6 months with longer-term outcomes apparent by 1 year. More complex wounds may require more than 6 months to heal.

Study Selection Criteria

- To assess efficacy outcomes, we sought comparative controlled prospective trials, with preference for RCTs.
- In the absence of such trials, we sought comparative observational studies, with preference for prospective studies.
- To assess longer-term outcomes and adverse effects, we sought single-arm studies that capture longer periods of follow-up and/or larger populations.
- Within each category of study design, prefer larger sample size studies and longer duration studies.
- We excluded studies with duplicative or overlapping populations.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials.

Review of Evidence

Systematic Reviews

A 2016 Cochrane review evaluated skin substitutes for the treatment of diabetic foot ulcers. Seventeen trials (N=1655) were included in the meta-analysis. Most trials identified were industry-sponsored, and an asymmetric funnel plot indicated publication bias. Pooled results of published trials found that skin

substitutes increased the likelihood of achieving complete ulcer closure compared with standard of care (SOC) alone (relative risk, 1.55; 95% CI, 1.30 to 1.85). Use of skin substitutes also led to a statistically significant reduction in amputations (relative risk, 0.43; 95% CI, 0.23 to 0.81), although the absolute risk difference was small. Analysis by individual products found a statistically significant benefit on ulcer closure for Apligraf, EpiFix, and Hyalograft-3D. The products that did not show a statistically significant benefit for ulcer closure were Dermagraft, GraftJacket, Kaloderm, and OrCel.

Apligraf, Dermagraft, AlloPatch, Integra Dermal Regeneration Template, or Integra Flowable Wound Matrix, mVASC, or TheraSkin

Apligraf

Veves et al (2001) reported on a randomized prospective trial on the effectiveness of Apligraf (previously called Graftskin), a living skin equivalent, in treating noninfected nonischemic chronic plantar diabetic foot ulcers. The trial involved 24 centers in the United States; 208 patients were randomized to ulcer treatment with Apligraf (112 patients) or saline-moistened gauze (96 patients, control group). Standard state-of-the-art adjunctive therapy, including extensive surgical debridement and adequate foot off-loading, was provided in both groups. Apligraf was applied at the beginning of the study and weekly thereafter for a maximum of 4 weeks (maximum of 5 applications) or earlier if complete healing occurred. At the 12-week follow-up visit, 63 (56%) Apligraf-treated patients achieved complete wound healing compared with 36 (38%) in the control group ($p=.004$). The Kaplan-Meier method median time to complete closure was 65 days for Apligraf, which was significantly lower than the 90 days observed in the control group ($p=.003$). The rates of adverse reactions were similar between groups, except osteomyelitis and lower-limb amputations, both of which were less frequent in the Apligraf group. Trialists concluded that application of Apligraf for a maximum of 4 weeks resulted in higher healing rates than state-of-the-art treatment and was not associated with any significant adverse events. This trial was reviewed in a 2001 TEC Assessment, which concluded that Apligraf, in conjunction with good local wound care, met the TEC criteria for the treatment of diabetic ulcers that fail to respond to conservative management.

Dermagraft

A 2003 pivotal multicenter FDA regulated trial randomized 314 patients with chronic diabetic ulcers to Dermagraft (human-derived fibroblasts cultured on mesh) or control. Over the 12-week study, patients received up to 8 applications of Dermagraft. All patients received pressure-reducing footwear and were encouraged to stay off their study foot as much as possible. At 12 weeks, the median percent wound closure for the Dermagraft group was 91% compared with 78% for the control group. Ulcers treated with Dermagraft closed significantly faster than ulcers treated with conventional therapy. No serious adverse events were attributed to Dermagraft. Ulcer infections developed in 10.4% of the Dermagraft patients compared with 17.9% of the control patients. Together, there was a lower rate of infection, cellulitis, and osteomyelitis in the Dermagraft-treated group (19% vs. 32.5%). A 2015 retrospective analysis of the trial data found a significant reduction in amputation/bone resection rates with Dermagraft (5.5% vs. 12.6%, $p=.031$). Of the 28 cases of amputation/bone resection, 27 were preceded by ulcer-related infection.

AlloPatch

AlloPatch Pliable human reticular acellular dermis was compared with SOC in an industry-sponsored multicenter trial by Zelen et al (2017, 2018). The initial trial with 20 patients per group was extended to determine the percent healing at 6 weeks with 40 patients per group. Healing was evaluated by the site investigator and confirmed by an independent panel. At 6 weeks, 68% (27/40) of wounds treated using AlloPatch had healed compared with 15% (6/40) in the SOC-alone group ($p<.001$). At 12 weeks, 80%

(32/40) of patients in the AlloPatch group had healed compared to 30% (12/40) in the control group. Mean time to heal within 12 weeks was 38 days (95% CI: 29-47 days) for the ADM group and 72 days (95% CI: 66 to 78 days) for the SOC group ($p<.001$).

Integra Omnigraft Dermal Regeneration Template or Integra Flowable Wound Matrix

Integra Dermal Regeneration Template is a biosynthetic skin substitute that is FDA approved for life-threatening thermal injury. The FOUNDER (Foot Ulcer New Dermal Replacement) multicenter study (32 sites) assessed Integra Dermal Regeneration Template (marketed as Omnigraft) for chronic nonhealing diabetic foot ulcers under an FDA regulated investigational device exemption. A total of 307 patients with at least 1 chronic diabetic foot ulcer were randomized to treatment with the Integra Template or a control condition (sodium chloride gel 0.9%). Treatment was given for 16 weeks or until wound closure. There was a modest increase in wound closure with the Integra Template (51% vs. 32%, $p=.001$) and a shorter median time to closure (43 days vs. 78 days, $p=.001$). There was a strong correlation between investigator-assessed and computerized planimetry assessment of wound healing ($r=0.97$). Kaplan-Meier analysis showed the greatest difference between groups in wound closure up to 10 weeks, with diminishing differences after 10 weeks. Trial strengths included adequate power to detect an increase in wound healing of 18%, which was considered to be clinically significant, secondary outcomes of wound closure and time to wound closure by computerized planimetry, and intention-to-treat (ITT) analysis.

Integra Flowable Wound Matrix is composed of a porous matrix of cross-linked bovine tendon collagen and glycosaminoglycan. It is supplied as a granular product that is mixed with saline. Campitiello et al (2017) published an RCT that compared the flowable matrix with wet dressing in 46 patients who had Wagner grade 3 diabetic foot ulcers. The ulcers had developed over 39 weeks. Complete healing at 6 weeks was achieved in significantly more patients in the Integra Flowable Wound Matrix group than in the control group, while the risk of rehospitalization and major amputation was reduced with Integra Flowable Wound Matrix (see Table 4).

Table 4. Probability of Wound Healing with IFWM Versus SOC

Study	Complete Wound Healing	Rehospitalization	Major Amputation
Campitiello et al (2017)			
IFWM, n (%)	20 (86.95)	2 (6.69)	1 (4.34)
SOC, n (%)	12 (52.17)	10 (43.47)	7 (30.43)
RR (95% CI)	1.67 (1.09 to 2.54)	0.10 (0.01 to 0.72)	0.16 (0.02 to 1.17)
p	.010	.001	.028

CI: confidence interval; IFWM: Integra Flowable Wound Matrix; RR: relative risk; SOC: standard of care.

mVASC

Tables 5 and 6 summarize the trial characteristics and results for RCTs of mVASC. Tables 7 and 8 evaluate study limitations.

Gould et al (2023) reported results of the HIFLO (Healing in Diabetic Foot Ulcers with Microvascular Tissue) Trial, a multicenter (6 US sites) RCT comparing weekly application of the processed

microvascular tissue (PMVT) allograft, mVASC in addition to a standardized diabetic foot ulcer protocol versus standard wound care with a collagen alginate dressing control in 100 adults with Wagner Grade 1 and 2 diabetic foot ulcers of ≥ 4 weeks and < 52 weeks duration.³¹ Wound and local peripheral neuropathy assessment were performed weekly. The primary outcome of the study was complete wound closure at 12 weeks. The investigator and a blinded physician made the initial determination of wound closure, followed by adjudication and confirmation by an independent, blinded panel of plastic surgeons. All participants who attended at least 1 treatment visit were included in the analysis. There was missing data for 15 participants at week 12 (3 in mVASC vs. 12 in control) and 14 of these were missing due to adverse events related to the wound. These were included in the primary analysis and counted as wound healing failures. The mean age of participants was 60 years, 90% of participants were White and 10% were Black, and 66% of participants were men. At randomization, the mean size of the wound area was 3.3 cm and the mean duration of the wound was 15 weeks. The proportion of participants with complete wound closure at week 12 was 74% (37/50) for mVASC versus 38% (19/50) for control ($p < .001$). Of the wounds that healed, the mean time to healing was also statistically significantly faster for the mVASC group (54 days; 95% CI, 46 to 61 vs 64 days; 95% CI, 57 to 72; $p = .009$). The 10-point Semmes-Weinstein monofilament (SWM) test of peripheral neuropathy also favored mVASC (118% vs. 11%; $p = .028$). No adverse events or serious adverse events related to the study treatment or the procedure were reported. There were 11 adverse events (3, mVASC vs. 8, control) reported that were related to the wound.

Table 5. Randomized Controlled Trial of mVASC for Diabetic Foot Ulcers-Characteristics

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Gould 2023; HIFLO	US	6	2017-2020	Adults with chronic Wagner Grade 1 or 2 DFU Mean age, 60 y 90% White 10% Black 66% Male Mean wound size 3.3 cm	mVASC + SOC (n=50)	SOC (n=50)

DFU: Diabetic Foot Ulcers; HIFLO: Healing in Diabetic Foot Ulcers with Microvascular Tissue; SOC: Standard of Care

Table 6. Randomized Controlled Trial of mVASC for Diabetic Foot Ulcers - Results

Study	Wounds Healed	Time to Heal	% Area Reduction	Adverse events
Gould 2023; HIFLO	at 12 weeks	by 12 weeks	at 12 weeks	
N analyzed	100	56	100	100
mVASC	74% (37/50)	Mean, 54 d	76%	3
SOC	38% (19/50)	Mean, 64 d	24%	8
p-value	$< .001$	.009	.009	

DFU: Diabetic Foot Ulcers; HIFLO: Healing in Diabetic Foot Ulcers with Microvascular Tissue; SOC: Standard of Care

Table 7. Randomized Controlled Trial of mVASC for Diabetic Foot Ulcers – Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Gould 2023; HIFLO	4. Lack of racial and ethnic diversity				1. follow-up not sufficient to determine ulcer recurrence.

HIFLO: Healing in Diabetic Foot Ulcers with Microvascular Tissue

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 8. Randomized Controlled Trial of mVASC for Diabetic Foot Ulcers – Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Gould 2023; HIFLO			1. Registered retrospectively in European registry			3. Confidence intervals not reported

HIFLO: Healing in Diabetic Foot Ulcers with Microvascular Tissue

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

TheraSkin Versus Standard of Care

Tables 9 and 10 summarize the trial characteristics and results for RCTs of TheraSkin compared to SOC. Tables 11 and 12 evaluate study limitations.

Armstrong et al (2022) reported results of an RCT including 100 adults with non-healing Wagner 1 diabetic foot ulcers comparing TheraSkin (n=50) to SOC (n=50).³² The index ulcer had to have been present for greater than 4 weeks and less than 1 year with a minimum size of 1.0 cm² and a maximum size of 25 cm². Standard of care included glucose monitoring, weekly debridement as appropriate, and an offloading device. The dressing in the SOC group was calcium alginate (Fibracol Plus). The primary outcome was the proportion of full-thickness wounds healed at 12 weeks. Wound healing was assessed initially by the investigator and confirmed by blinded adjudication panel. Wounds were closed when there was 100% re-epithelization and no drainage. The mean age of participants was 60 years; 53% of participants were male, 70% were White, and 15% were Black. The mean wound area at baseline was 4.1 cm². Participants who did not have healing of at least 50% by 6 weeks were allowed to seek alternative rescue wound care (TheraSkin, n=1; SOC, n=11). In addition, 3 participants in the TheraSkin group and 8 in the SOC group had worsening of the wound or an adverse event before week 12. All

enrolled participants were included in analysis and missing data were imputed using last observation carried forward. The percent of participants with complete wound healing at week 12 was 76% (38/50) in the intervention group compared with 36% (18/50) in the SOC group ($p<.01$). The mean percent area reduction at 12 weeks was 77.8% in the TheraSkin group compared with 49.6% in the SOC group ($p<.01$). There were no statistically significant differences between groups in QOL or pain score measures.

Table 9. Randomized Controlled Trial of TheraSkin vs. SOC for Diabetic Foot Ulcers – Characteristics

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Armstrong (2022); NCT04040426	US	5	2019- 2021	Adults with non-healing Wagner 1 DFUs Mean wound area, 4.1 cm ² Mean age, 60 yrs 53% male 70% White 15% Black	TheraSkin (n=50)	SOC with calcium alginate dressing (n=50)

DFU: Diabetic Foot Ulcers;; SOC: standard of care

Table 10. Randomized Controlled Trial of TheraSkin vs. SOC for Diabetic Foot Ulcers – Results

Study	Wounds Healed	Time to Heal	% Area Reduction	Adverse events
Armstrong (2022); NCT04040426	at 12 weeks	by 12 weeks	at 12 weeks	
N analyzed	100	100	100	100
TheraSkin	76% (38/50)	Mean, 47 days (95% CI, 39 to 55)	78% (SD=63)	2
SOC	36% (18/50)	Mean, 65 days (95% CI, 58 to 73)	50% (SD=98)	4
-value	<.01	<.01	<.01	NR

CI: confidence interval; NR: not reported; SD: standard deviation; SOC: standard of care

Table 11. Randomized Controlled Trial of TheraSkin vs. SOC for Diabetic Foot Ulcers – Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Armstrong (2022); NCT04040426	4. Lack of racial and ethnic diversity				1. follow-up not sufficient to determine ulcer recurrence.

SOC: standard of care.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not

supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 12. Randomized Controlled Trial of TheraSkin vs. SOC for Diabetic Foot Ulcers – Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Armstrong (2022); NCT04040426		1. Investigators not blinded		2. Missing data imputed by last observation carried forward; no sensitivity analyses provided		

SOC: standard of care.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

TheraSkin Versus Dermagraft

Sanders et al (2014) reported on an (N=23) industry-funded randomized comparison of TheraSkin (cryopreserved human skin allograft with living fibroblasts and keratinocytes) and Dermagraft for diabetic foot ulcers. Wound size at baseline ranged from 0.5 to 18.02 cm²; the average wound size was about 5 cm² and was similar for the 2 groups (p=.51). Grafts were applied according to manufacturers' instructions over the first 12 weeks of the study until healing, with an average of 4.4 TheraSkin grafts (every 2 weeks) compared with 8.9 Dermagraft applications (every week). At week 12, complete wound healing was observed in 63.6% of ulcers treated with TheraSkin and 33.3% of ulcers treated with Dermagraft (p<.049). At 20 weeks, complete wound healing was observed in 90.9% of the TheraSkin-treated ulcers compared with 66.7% of the Dermagraft group (p=.428).

TheraSkin Versus Apligraf

DiDomenico et al (2011) compared TheraSkin with Apligraf for the treatment of diabetic foot ulcers in a (N=29) RCT. The risk of bias in this study is uncertain because reporting did not include a description of power analysis, statistical analysis, method of randomization, or blinding. The percentage of wounds closed at 12 weeks was 41.3% in the Apligraf group and 66.7% in the TheraSkin group. Results at 20 weeks were not substantially changed from those at 12 weeks, with 47.1% of wounds closed in the Apligraf group and 66.7% closed in the TheraSkin group. The percentage healed in the Apligraf group was lower than expected based on prior studies. The average number of grafts applied was similar for both groups (1.53 for Apligraf, 1.38 for TheraSkin). The low number of dressing changes may have influenced results, with little change in the percentage of wounds closed between 12 and 20 weeks. An adequately powered trial with blinded evaluation of wound healing and a standard treatment regimen would permit greater certainty on the efficacy of this product.

Section Summary: Apligraf, Dermagraft, AlloPatch, Integra, mVASC or TheraSkin for Diabetic Lower-Extremity Ulcers

RCTs reporting complete wound healing outcomes with at least 12 weeks of follow-up have demonstrated the efficacy of Apligraf, Dermagraft, AlloPatch, Integra Dermal Regeneration Template, Integra Flowable Wound Matrix, mVASC, and TheraSkin over SOC for the treatment of diabetic lower-extremity ulcers.

Bioengineered Skin Substitutes other than Apligraf, Dermagraft, AlloPatch, Integra, mVASC, or TheraSkin

GraftJacket Regenerative Tissue Matrix

Brigido et al (2004) reported a (N=40) randomized pilot study comparing GraftJacket with conventional treatment for chronic nonhealing diabetic foot ulcers. Control patients received conventional therapy with debridement, wound gel with gauze dressing, and off-loading. GraftJacket patients received surgical application of the scaffold using skin staples or sutures and moistened compressive dressing. A second graft application was necessary after the initial application for all patients in the GraftJacket group. Preliminary 1-month results showed that, after a single treatment, ulcers treated with GraftJacket healed at a faster rate than conventional treatment. There were significantly greater decreases in wound length (51% vs. 15%), width (50% vs. 23%), area (73% vs. 34%), and depth (89% vs. 25%), respectively. With follow-up to 4 weeks, no data were reported on the proportion with complete closure or the mean time to heal. All grafts were incorporated into the host tissue.

Reyzelman et al (2009) reported an industry-sponsored multicenter randomized study that compared a single application of GraftJacket with SOC in 86 patients with diabetic foot ulcers. Eight patients, 6 in the study group and 2 in the control group, did not complete the trial. At 12 weeks, complete healing was observed in 69.6% of the GraftJacket group and 46.2% of controls. After adjusting for ulcer size at presentation, a statistically significant difference in nonhealing rate was calculated, with odds of healing 2.0 times higher in the study group. Mean healing time was 5.7 weeks for the GraftJacket group versus 6.8 weeks for the control group. The authors did not report whether this difference was statistically significant. Median time to healing was 4.5 weeks for GraftJacket (range, 1-12 weeks) and 7.0 weeks for control (range, 2-12 weeks). Kaplan-Meier method survivorship analysis for time to complete healing at 12 weeks showed a significantly lower nonhealing rate for the study group (30.4%) than for the control group (53.9%). The authors commented that a single application of GraftJacket, as used in this study, was often sufficient for complete healing.

Reyzelman and Bazarov (2015) reported an industry-sponsored meta-analysis of GraftJacket for diabetic foot ulcers that included the 2 studies described above and a third RCT by Brigido (2006) with 28 patients (N=154). The time to heal was estimated for the Brigido (2004) study, based on the average wound reduction per week. The estimated difference in time to heal was considerably larger for Brigido's (2004) study (-4.30 weeks) than for the other 2 studies that measured the difference in time to heal (-1.58 weeks and -1.10 weeks). Analysis of the proportion of wounds that healed included Brigido (2006) and Reyzelman et al (2009). The odds ratio in the smaller study by Brigido (2006) was considerably larger, with a lack of precision in the estimate (odds ratio, 15.0; 95% CI, 2.26 to 99.64), and the combined odds (3.75; 95% CI, 1.72 to 8.19) was not significant when analyzed using a random-effects model. Potential sources of bias, noted by Reyzelman and Bazarov (2015), included publication and reporting biases, study selection biases, incomplete data selection, post hoc manipulation of data, and subjective choice of analytic methods. Overall, results of these studies do not provide convincing evidence that GraftJacket is more effective than SOC for healing diabetic foot ulcers.

DermACELL Versus GraftJacket Regenerative Tissue Matrix or Standard of Care

DermACELL and GraftJacket are both composed of human ADM. Walters et al (2016) reported on a multicenter randomized comparison of DermACELL, GraftJacket, or SOC (2:1:2 ratio) in 168 patients with diabetic foot ulcers. The study was sponsored by LifeNet Health, a nonprofit organ procurement association and processor for DermACELL. At 16 weeks, the proportion of completely healed ulcers was 67.9% for DermACELL, 47.8% for GraftJacket, and 48.1% for SOC. The 20% difference in completely healed ulcers was statistically significant for DermACELL versus SOC ($p=.039$). The mean time to complete wound closure did not differ significantly for DermACELL (8.6 weeks), GraftJacket (8.6 weeks), and SOC (8.7 weeks).

A second report from this study was published in 2017. This analysis compared DermACELL with SOC and did not include the GraftJacket arm. The authors reported that either 1 or 2 applications of DermACELL led to a greater proportion of wounds healed compared with SOC in per-protocol analysis (see Table 5), but there was no significant difference between DermACELL (1 or 2 applications) and SOC when analyzed by ITT. For the group of patients who received only a single application, the percentage of patients who achieved complete wound healing was significantly higher than SOC at 16 and 24 weeks, but not at 12 weeks. Although reported as an ITT analysis, results were analyzed only for the group who received a single application of DermACELL. This would not typically be considered ITT.

Table 13. Probability of Wound Healing in Per Protocol Analysis of DermACELL Versus SOC

	% With Wound Healing at 12 Wk	% With Wound Healing at 16 Wk	% With Wound Healing at 24 Wk	% With Wound Healing at 12 Wk	% With Wound Healing at 16 Wk	% With Wound Healing at 24 Wk
Cazzell et al (2017)						
DermACELL, %	65.0%	82.5%	89.7%	NR	67.9%	83.7%
SOC, %	41.1%	48.1%	67.3%	NR	48.1%	67.3%
HR (95% CI)	1.97 (1.1 to 3.5)	2.40 (1.4 to 4.1)	2.11 (1.3 to 3.5)		1.72 (1.04 to 2.83)	1.55 (0.98 to 2.44)
p	.012	<.001	<.001	NS	.028	.049

CI: confidence interval; HR: hazard ratio; NR: not reported; NS: not significant; SOC: standard of care.

Cyral (MatriStem) Versus Dermagraft

Frykberg et al (2017) reported a prespecified interim analysis of an industry-funded multicenter noninferiority trial of Cyral (a porcine urinary bladder-derived extracellular matrix) versus Dermagraft in 56 patients with diabetic foot ulcers. The mean duration of ulcers before treatment was 263 days (range, 30 to 1095 days). The primary outcome was the percent wound closure with up to 8 weeks of treatment using blinded evaluation of photographs. ITT analysis found complete wound closure in 5 (18.5%) wounds treated with Cyral compared with 2 (6.9%) wounds treated with Dermagraft ($p=\text{not significant [NS]}$). QOL, measured by the Diabetic Foot Ulcer Scale, improved from 181.56 to 151.11 in the Cyral group and from 184.46 to 195.73 in the Dermagraft group ($p=.074$). It should be noted that this scale is a subjective

measure and patients were not blinded to treatment. Power analysis indicated that 92 patients would be required; further recruitment is ongoing for completion of the study.

PriMatrix

Lantis et al (2021) reported on a multicenter RCT comparing PriMatrix plus SOC to PriMatrix alone in 226 patients with diabetic foot ulcers (Tables 14 and 15).

Study subjects underwent a 2-week run-in period of SOC treatment and were excluded if they had a wound reduction of 30% or more. Patients randomized to the SOC group received weekly treatment at the study site identical to the SOC treatment applied during the screening period. In addition, control group patients performed daily dressing changes, which consisted of wound cleaning, application of saline gel and secondary dressings. The primary endpoint was the percentage of subjects with complete wound closure, defined as 100% re-epithelialization without drainage during the 12-week treatment phase.

Significantly more patients in the PriMatrix group experienced complete wound closure at 12 weeks (45.6% vs 27.9%; $p=.008$). It is unclear if this difference (17.7%) is clinically significant; the study was powered to detect a 20% difference between groups. The time to complete healing did not differ between groups for the wounds that healed. Major study limitations include lack of blinding, limited generalizability, and insufficient duration of follow-up to assess wound recurrence (Tables 16 and 17).

Table 14. Randomized Controlled Trial of PriMatrix for Diabetic Foot Ulcers- Characteristics

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Lantis et al (2021) NCT03010319	US	21	2019-2020	Diabetic foot ulcer for a minimum of 2 weeks, adequate vascular perfusion to the affected extremity	PriMatrix plus standard of care n = 103	Standard of care n = 104

Table 15. Randomized Controlled Trial of PriMatrix for Diabetic Foot Ulcers- Results

Study	Wound Healed at 12 weeks	Median Time to Heal, days (range)	AEs
Lantis et al (2021) NCT03010319			
Number analyzed	207	76	226
PriMatrix	47/103 (45.6%)	43 (22 to 93)	Any AE: 44.8%
Standard Care	29/104 (27.9%)	57 (16 to 88)	Any AE: 46.4%
Treatment Effect	HR 2.02 (95% CI 1.3 to 3.2)		

Study	Wound Healed at 12 weeks	Median Time to Heal, days (range)	AEs
p	.008	.362	

AE: adverse events; CI: confidence interval; HR: hazard ratio

Table 16. Randomized Controlled Trial of PriMatrix for Diabetic Foot Ulcers- Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Lantis et al (2021) NCT03010319	4. Race and ethnicity of the study population was not reported and is not included in the demographics table.		3. Standard of care patients received additional dressing changes at home, which could have potentially exposed the wound to unknown factors.		4-week follow-up not sufficient to determine ulcer recurrence.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 17. Randomized Controlled Trial of PriMatrix for Diabetic Foot Ulcers- Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Lantis et al (2021) NCT03010319	3. Allocation concealment not described.	1. Patients and investigator not blinded		1. 24 subjects from the treatment group and 22 from the control group discontinued from each arm prior to meeting the protocol-defined primary endpoint and were counted as treatment failures. 207 of 226 randomized were included in primary analysis (91.6%)		3. Confidence intervals not reported

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important

difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other

Oasis Wound Matrix Versus Regranex Gel

Niezgoda et al (2005) compared healing rates at 12 weeks for full-thickness diabetic foot ulcers treated with OASIS Wound Matrix (a porcine acellular wound care product) to Regranex Gel. This industry-sponsored, multicenter RCT was conducted at 9 outpatient wound care clinics and involved 73 patients with at least 1 diabetic foot ulcer. Patients were randomized to receive either Oasis Wound Matrix (n=37) or Regranex Gel (n=36) and secondary dressing. Wounds were cleaned and debrided, if needed, at a weekly visit. The maximum treatment period for each patient was 12 weeks. After 12 weeks, 18 (49%) Oasis-treated patients had complete wound closure compared with 10 (28%) Regranex-treated patients. Oasis treatment met the noninferiority margin but did not demonstrate that healing in the Oasis group was statistically superior ($p=.055$). Post hoc subgroup analysis showed no significant difference in incidence of healing in patients with type 1 diabetes (33% vs. 25%) but showed a significant improvement in patients with type 2 diabetes (63% vs. 29%). There was also increased healing of plantar ulcers in the Oasis group (52% vs. 14%). These post hoc findings are considered hypothesis-generating. Additional study with a larger number of subjects is needed to compare the effect of Oasis treatment to current SOC.

Autologous Grafting on HYAFF Scaffolds

Uccioli et al (2011) reported a multicenter RCT of cultured expanded fibroblasts and keratinocytes grown on an HYAFF scaffold (benzyl ester of hyaluronic acid) compared with paraffin gauze for difficult diabetic foot ulcers. A total of 180 patients were randomized. At 12 weeks, complete ulcer healing was similar for the 2 groups (24% treated vs. 21% controls). At 20 weeks, complete ulcer healing was achieved in a similar proportion of the treatment group (50%) and the control group (43%, log-rank test = 0.344). Subgroup analysis, adjusted for baseline factors and possibly post-hoc, found a statistically significant benefit of treatment on dorsal ulcers but not plantar ulcers.

Mirragen Advanced Wound Matrix

Armstrong et al (2022) reported a multi-center, single-blinded RCT (N=40; 5 outpatient wound care centers) comparing Mirragen Advanced Wound Matrix (BBGFM), a borate-based bioactive glass fiber matrix, with SOC in subjects with chronic, full-thickness, non-infected, non-ischemic diabetic foot ulcers (University of Texas 1A/Wagner 1). Both groups received SOC including weekly debridement as needed and offloading; the BBGFM group received weekly application of the matrix in addition to SOC. At 12 weeks, complete wound closure was achieved in 70% (14/20) of BBGFM-treated subjects compared with 25% (5/20) of SOC subjects (adjusted $P=.006$); the corresponding number needed to treat was 2.0 (95% CI: 1.4 to 5.8). Mean PAR at 12 weeks was 79% in the BBGFM group compared with 37% in the SOC group (adjusted $P=.027$). The authors noted that investigator blinding was not possible given the absence of a sham product, and that withdrawal of non-responding subjects at 6 weeks resulted in censoring of outcomes.

Armstrong et al (2025) reported a multi-center, single-blind RCT (N=148; 14 sites) comparing BBGFM plus SOC with SOC alone in subjects with chronic, full-thickness, non-infected, non-ischemic diabetic foot ulcers (University of Texas 1A/Wagner Grade 1). The trial was stopped early following a pre-specified interim analysis. In the mITT analysis, 48% (32/67) of BBGFM-treated subjects achieved complete wound closure at 12 weeks compared with 24% (16/66) of SOC subjects ($p=.007$); mean time to healing was 9.1 weeks (95% CI: 8.1 to 10.0) versus 10.4 weeks (95% CI: 9.6 to 11.1) for SOC (adjusted $p=0.042$). In the

per-protocol analysis, 73% (32/44) of BBGFM subjects healed compared with 42% (16/38) of SOC subjects ($p=.007$); mean time to healing was 8.2 weeks (95% CI: 7.0 to 9.4) versus 9.7 weeks (95% CI: 8.6 to 10.7) for SOC (adjusted $p=0.084$, not statistically significant). The authors identified investigator non-blinding and early withdrawal of non-responding subjects as study limitations, noting that it is unknown how many of the 36 withdrawn subjects (24% of the mITT population) might have healed within 12 weeks had they not been withdrawn.

Kerecis OMEGA3 Wound

Systematic Reviews

Ruiz-Muñoz et al (2024) conducted a systematic review and meta-analysis of RCTs comparing fish skin grafts (Kerecis Omega3 Wound) with standard care for diabetic foot ulcer healing. Five RCTs with a total of 411 patients were included. Pooled analysis showed a significantly higher rate of complete ulcer healing in the fish skin graft groups compared with standard care (OR, 3.34; 95% CI, 2.14 to 5.20; $p<.01$), with no heterogeneity between studies ($I^2=0\%$). Sensitivity analysis confirmed the robustness of findings, and publication bias testing did not indicate asymmetry. The authors noted that blinding of participants and personnel was compromised in 3 of the 5 included trials.

Randomized Controlled Trials

Lullove et al (2021, 2022) reported interim results and Lantis et al (2023) of a RCT of Omega3 Wound (Kerecis) plus standard wound care compared to standard care alone in individuals with diabetic lower extremity skin ulcers (Table 18). The primary outcome of the trial was healing at 12 weeks. Complete ulcer healing was based on the site investigator's assessment, as evidenced by complete (100%) re-epithelialization without drainage and need of dressing. An independent panel of wound care experts who were blinded to the patient allocation process and the principal investigator's assessment reviewed all study-related decisions made by the site investigators and confirmed healing status. Secondary outcomes were time to heal and wound area reduction by percentage at 12 weeks. Patients underwent a 2-week run-in period prior to randomization. If the ulcer reduced in area by 20% or more after 14 days of standard care, the patient was excluded as a screening failure. If the wound area was reduced by less than 20%, the patient was randomized and enrolled in the study.

Study results are summarized in Table 19. At 12 weeks, the complete healing rate was significantly higher in the intervention arm (57% vs 31%), but time to healing did not differ between groups for wounds that healed completely. Among the subset of wounds that did not heal completely by 12 weeks ($n = 65$), there was a larger percent wound reduction in the intervention group (86% vs 64%; $p=.03$). Of the 45 participants whose wound healed during the 12 weeks of the trial, 42 were available for follow-up 6 to 12 months following healing.

Dardari et al (2024) reported the Odinn trial (NCT04257370), an international, open-label, multicenter RCT comparing intact fish skin graft (Kerecis Omega3 Wound) with standard wound care in patients with deep diabetic foot ulcers penetrating to bone, joint, or tendon (University of Texas [UT] grade 2 or 3). A total of 255 patients were randomized (129 fish skin graft, 126 standard care) across 15 tertiary care centers in France, Italy, Germany, and Sweden. Treatment was applied weekly for 6 weeks, then biweekly through week 14, with healing assessed at 16 weeks by a blinded adjudication committee. At 16 weeks, 44.0% of wounds healed in the fish skin graft group compared with 26.4% in the standard care group ($p<.001$; OR, 2.55; 95% CI, 1.43 to 4.63), with continued healing observed at 20 weeks (45.7% vs 32.4%) and 24 weeks (55.2% vs 37.8%). Cox regression analysis showed fish skin graft was associated with faster time to healing (HR, 1.59; 95% CI, 1.07 to 2.36). Primary wound infection rates were similar between groups (30.2% vs

29.3%). The study was limited by its open-label design; however, wound closure was confirmed by a blinded adjudication committee. Study results are summarized in Table 19.

Study limitations are detailed in Tables 20 and 21.

Table 18. Randomized Controlled Trial of Omega3 Wound for Diabetic Foot Ulcers- Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Lantis et al (2023) Lullove et al (2021) NCT04133493	US	16	2019-2022	Diabetic foot ulcer for a minimum of 4 weeks, adequate renal function and perfusion to the affected extremity Mean age, 60 y 69% Men 80% White 7% Black Mean wound size, 4.4 cm	Omega3 Wound plus standard of care (n=51)	Standard of care (n=51)
Dardari et al (2024); NCT04257370	France, Italy, Germany, Sweden	15	2020-2022	Diabetic foot ulcer penetrating to bone, joint, or tendon (UT grade 2 or 3) present for ≥1 month, or open amputation wounds; ABPI ≥0.6; 78% Men; UT grade 2: 58.8%, UT grade 3: 41.2%; 50-52% amputation wounds	Intact fish skin graft (Kerecis Omega3 Wound) plus standard of care (n=129)	Standard of care (n=126)

ABPI: ankle brachial pressure index; UT: University of Texas Wound Classification System Grade; y: years.

Table 19. Randomized Controlled Trial of Omega3 Wound for Diabetic Foot Ulcers- Results

Study	Wound Healed at 12 weeks	Time to Heal	Percent Wound Reduction at 12 Weeks for Wounds that did not heal)	Adverse events
Lantis et al (2023) Lullove et al (2021) NCT04133493				
N analyzed	102		65	
Omega3 Wound	57 % (29/ 51)	Mean 7 weeks in both groups	86%	3
Standard Care	31 % (16/ 51)		64%	5

Study	Wound Healed at 12 weeks		Time to Heal	Percent Wound Reduction at 12 Weeks for Wounds that did not heal)	Adverse events
p-value	.02			.03	
Dardari et al (2024) NCT04257370	Wound Healed at 16 weeks	Wound Healed at 24 weeks	Time to Heal		Adverse events
Total N=255					
Omega3 Wound	44.0% (57/129)	55.2%	Mean 17.31 weeks (95% CI, 15.5 to 18.67)		79/129 (61%) experienced ≥ 1 AE; wound infections in 39 (30.2%)
Standard Care	26.4% (33/126)	37.8%	Mean 19.37 weeks (95% CI, 18.09 to 20.66)		67/126 (53%) experienced ≥ 1 AE; wound infections in 37 (29.3%)
p-value	<.001	NR			
HR			1.59 (95% CI, 1.07 to 2.36)		

AE: adverse event; CI: confidence interval; NR: not reported.

Table 20. Randomized Controlled Trial of Omega3 Wound for Diabetic Foot Ulcers- Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Lantis et al 2023 Lullove et al (2021) NCT04133493	4. Lack of racial and ethnic diversity		3. Standard of care patients received additional dressing changes at home, which could have potentially exposed the wound to unknown factors.		
Dardari et al (2024); NCT04257370			3. Standard of care dressings were chosen by the investigator, introducing variability in the control arm		1. Primary endpoint at 16 weeks; follow-up through 24 weeks only

ITT: intension-to-treat; RCT: randomized controlled trial.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of

interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 21. Randomized Controlled Trial of Omega3 Wound for Diabetic Foot Ulcers- Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Lantis et al (2023) Lullove et al (2021) NCT04133493				1, 2. 25% of participants did not complete week 12. Although they were included in the primary ITT analysis, the method of imputation was unclear.		3. Confidence intervals not reported
Dardari et al (2024); NCT04257370		1. Open-label design; patients and wound care staff were unblinded due to differences in appearance of fish skin graft and standard dressings. Wound closure was confirmed by a blinded adjudication committee.				

ITT: intention-to-treat; RCT: randomized controlled trial.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

Section Summary: Bioengineered Skin Substitutes other than Apligraf, Dermagraft, AlloPatch, Integra, mVASC or TheraSkin for Diabetic Lower-Extremity Ulcers

Results from a multicenter RCT showed some benefit of DermACELL that was primarily for the subgroup of individuals who only required a single application of the ADM. Studies are needed to further define the population who might benefit from this treatment. Additional study with a larger number of subjects is needed to evaluate the effect of GraftJacket, DermACELL, Cytal, PriMatrix, and Oasis Wound Matrix, compared with current SOC or other advanced wound therapies.

Lower-Extremity Ulcers due to Venous Insufficiency

Clinical Context and Therapy Purpose

The purpose of bio-engineered soft tissue substitutes in individuals who have lower extremity ulcers due to venous insufficiency is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals who have lower extremity ulcers due to venous insufficiency.

Interventions

The therapy being considered is bioengineered skin substitutes.

Comparators

The following therapies are currently being used: SOC which includes debridement of necrotic tissue and compression.

A Cochrane review by O' Meara et al (2012) that evaluated compression for venous leg ulcers included 48 RCTs with 59 different comparisons. Most RCTs were small. Measures of healing were the time to complete healing, the proportion of ulcers healed within the trial period (typically 12 weeks), the change in ulcer size, and the rate of change in ulcer size. Evidence from 8 trials indicated that venous ulcers healed more rapidly with compression than without. Findings suggested that multicomponent systems (bandages or stockings) were more effective than single-component compression. Also, multicomponent systems containing an elastic bandage appeared more effective than those composed mainly of inelastic constituents. Although these meta-analyses did not include time to healing, studies included in the review reported the mean time to ulcer healing was approximately 2 months, while the median time to healing in other reports was 3 to 5 months.

Outcomes

The general outcomes of interest are symptoms, change in disease status, morbid events, and QOL.

The primary endpoints of interest for trials of wound closure are as follows, consistent with guidance from the FDA for industry in developing products for treatment of chronic cutaneous ulcer and burn wounds:

- Incidence of complete wound closure.
- Time to complete wound closure (reflecting accelerated wound closure).
- Incidence of complete wound closure following surgical wound closure.
- Pain control.

Time to wound closure can be measured at 6 months with longer-term outcomes apparent by 1 year. Complex wounds may require more than 6 months to heal.

Study Selection Criteria

- To assess efficacy outcomes, we sought comparative controlled prospective trials, with preference for RCTs*.
- In the absence of such trials, we sought comparative observational studies, with preference for prospective studies.
- To assess longer-term outcomes and adverse effects, we sought single-arm studies that capture longer periods of follow-up and/or larger populations.
- Within each category of study design, we prefer larger sample size studies and longer duration studies.
- We excluded studies with duplicative or overlapping populations.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials.

Review of Evidence

Apligraf

Falanga et al (1998) reported on a multicenter randomized trial of Apligraf living cell therapy. A total of 293 patients with venous insufficiency and clinical signs of venous ulceration were randomized to compression therapy alone or to compression therapy and treatment with Apligraf. Apligraf was applied up to a maximum of 5 (mean, 3.3) times per patient during the initial 3 weeks. The primary endpoints were the percentage of patients with complete healing by 6 months after initiation of treatment and the time required for complete healing. At 6-month follow-up, the percentage of patients healed was higher with Apligraf (63% vs. 49%), and the median time to complete wound closure was shorter (61 days vs. 181 days). Treatment with Apligraf was superior to compression therapy in healing larger (>1000 mm²) and deeper ulcers and ulcers of more than 6 months in duration. There were no symptoms or signs of rejection, and the occurrence of adverse events was similar in both groups. This study was reviewed in a 2001 TEC Assessment, which concluded that Apligraf (Graftskin), in conjunction with good local wound care, met TEC criteria for the treatment of venous ulcers that fail to respond to conservative management.

Oasis Wound Matrix

Mostow et al (2005) reported on an industry-sponsored multicenter (12 sites) randomized trial that compared weekly treatment using Oasis Wound Matrix (xenogeneic collagen scaffold from porcine small intestinal mucosa) with SOC in 120 patients who had chronic ulcers due to venous insufficiency that had not adequately responded to conventional therapy. Healing was assessed weekly for up to 12 weeks, with follow-up performed after 6 months to assess recurrence. After 12 weeks of treatment, there was a significant improvement in the percentage of wounds healed in the Oasis group (55% vs. 34%). After adjusting for baseline ulcer size, patients in the Oasis group were 3 times more likely to heal than those in the group receiving SOC. Patients in the SOC group whose wounds did not heal by week 12 were allowed to cross over to Oasis treatment. None of the healed patients treated with Oasis wound matrix who was seen for the 6-month follow-up experienced ulcer recurrence.

A research group in Europe has described 2 comparative studies of the Oasis matrix for mixed arteriovenous ulcers. In a quasi-randomized study, Romanelli et al (2007) compared the efficacy of 2

extracellular matrix-based products, Oasis and Hyaloskin (extracellular matrix with hyaluronic acid). Fifty-four patients with mixed arteriovenous leg ulcers were assigned to the 2 arms based on order of entry into the study; 50 patients completed the study. Patients were followed twice weekly, and dressings changed more than once a week, only when necessary. After 16 weeks of treatment, complete wound closure was achieved in 82.6% of Oasis-treated ulcers compared with 46.2% of Hyaloskin-treated ulcers. Oasis treatment significantly increased the time to dressing change (mean, 6.4 days vs. 2.4 days), reduced pain on a 10-point scale (3.7 vs. 6.2), and improved patient comfort (2.5 vs. 6.7).

Romanelli et al(2010) compared Oasis with a moist wound dressing (SOC) in 23 patients with mixed arteriovenous ulcers and 27 patients with venous ulcers. The trial was described as randomized, but the method of randomization was not described. After the 8-week study period, patients were followed monthly for 6 months to assess wound closure. Complete wound closure was achieved in 80% of the Oasis-treated ulcers at 8 weeks compared with 65% of the SOC group. On average, Oasis-treated ulcers achieved complete healing in 5.4 weeks compared with 8.3 weeks for the SOC group. Treatment with Oasis also increased the time to dressing change (5.2 days vs. 2.1 days) and the percentage of granulation tissue formed (65% vs. 38%).

Section Summary: Apligraf or Oasis Wound Matrix for Lower-Extremity Ulcers due to Venous Insufficiency

RCTs have demonstrated the efficacy of Apligraf or Oasis Wound Matrix over SOC for lower-extremity ulcers due to venous insufficiency.

Bioengineered Skin Substitutes other than Apligraf or Oasis Wound Matrix for Lower-Extremity Ulcers due to Venous Insufficiency

Dermagraft

Dermagraft living cell therapy has been approved by the FDA for repair of diabetic foot ulcers. Use of Dermagraft for venous ulcers is an off-label indication. Harding et al (2013) reported an open-label multicenter RCT that compared Dermagraft plus compression therapy (n=186) with compression therapy alone (n=180). The trial had numerous inclusion and exclusion criteria that restricted the population to patients who had nonhealing ulcers with compression therapy but had the capacity to heal. ITT analysis revealed no significant difference between the 2 groups in the primary outcome measure, the proportion of patients with completely healed ulcers by 12 weeks (34% Dermagraft vs. 31% control). Prespecified subgroup analysis revealed a significant improvement in the percentage of wounds healed for ulcers of 12 months or less in duration (52% vs. 37%) and for ulcers of 10 cm or less in diameter (47% vs. 39%). There were no significant differences in the secondary outcomes of time to healing, complete healing by week 24, and percent reduction in ulcer area.

DermACELL

Cazzell (2019) published an RCT on DermACELL ADM for venous leg ulcers in 18 patients (see Table 22). This was part of a larger study of the acellular dermal matrix for chronic wounds of the lower extremity in 202 patients; the component on diabetic lower extremity ulcers was previously reported by Cazzell et al (2017) and is described above. When including patients who required more than 1 application of the ADM, the percent of wounds closed at 24 weeks was 29.4% with DermACELL and 33.3% with SOC, suggesting no benefit DermACELL for the treatment of venous ulcers in this small sub study.

Table 22. Summary of Key RCT Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Cazzell (2019) NCT01970163	US	7	2013- 2016	Venous leg ulcer present for at least 60 days (n=18)	1 or 2 applications of DermACELL plus SOC (n=18)	SOC (debridement and compression, n=10)

RCT: randomized controlled trial; SOC: standard of care

Section Summary: Bioengineered Skin Substitutes other than Apligraf or Oasis Wound Matrix for Lower-Extremity Ulcers due to Venous Insufficiency

In a moderately large RCT, Dermagraft was not shown to be more effective than controls in the primary or secondary endpoints for the entire population and was slightly more effective than controls (an 8%-15% increase in healing) only in subgroups of patients with ulcer duration of 12 months or less or wound diameter of 10 cm or less. An initial study with 18 patients found that and DermACELL (ADM) was not more effective than SOC.

Deep Dermal Burns

Clinical Context and Therapy Purpose

The purpose of bio-engineered soft tissue substitutes in individuals who have deep dermal burns is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with deep dermal burns.

Interventions

The therapy being considered is bioengineered skin substitutes.

Comparators

The following therapies are currently being used: standard therapy for burns.

Outcomes

The general outcomes of interest are disease-specific survival, symptoms, change in disease status, morbid events, functional outcomes, QOL, and treatment-related morbidity.

The primary endpoints of interest for trials of wound closure are as follows, consistent with guidance from the FDA for industry in developing products for treatment of chronic cutaneous ulcer and burn wounds:

- Incidence of complete wound closure.
- Time to complete wound closure (reflecting accelerated wound closure).
- Incidence of complete wound closure following surgical wound closure.
- Pain control.

Time to wound closure can be measured at 6 months with longer-term outcomes apparent by 1 year.

Study Selection Criteria

- To assess efficacy outcomes, we sought comparative controlled prospective trials, with preference for RCTs*.
- In the absence of such trials, we sought comparative observational studies, with preference for prospective studies.
- To assess longer-term outcomes and adverse effects, we sought single-arm studies that capture longer periods of follow-up and/or larger populations.
- Within each category of study design, we prefer larger sample size studies and longer duration studies.
- We excluded studies with duplicative or overlapping populations.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials.

Review of Evidence

Biobrane

Data regarding the use of synthetic Silicone/Nylon Membrane wound dressing (e.g., Biobrane) has been described in four separate randomized controlled trials (RCTs) in peer-reviewed published medical journals (Barret, 2000; Feldman, 1991; Gerding, 1990; Lal, 2000). All of these studies found that in comparison to their various control groups the use of Biobrane significantly improved pain scores and healing times.

Based on clinical practice standards, the above-mentioned evidence, and the overall clinical experience with Biobrane, an acceptable level of safety and efficacy has been established for the use of this product in the treatment of burns.

Epicel

One case series from 2000 has described the treatment of 30 severely burned patients with Epicel. The cultured epithelial autografts were applied to a mean of 37% of total body surface area (TBSA). Epicel achieved permanent coverage of a mean of 26% of TBSA, an area similar to that covered by conventional autografts (mean, 25%). Survival was 90% in these severely burned patients.

EZ Derm

The available evidence addressing the use of the EZ Derm brand porcine derived decellularized fetal skin is limited. Data from two moderately sized retrospective case series studies have been published. The first involved 157 subjects with partial-thickness burns treated with EZ Derm (Troy, 2014). The authors reported 19 complications, including premature separation of graft (n=9, 16%), infection (n=4, 3%), and need for excision (n=6, 4.5%). The other study involved 164 subjects, also with burns (Burkey, 2016). The authors reported a significant decrease in average narcotic dose following treatment ($p < 0.001$) and fewer

dressings changes needed ($p < 0.001$). Only 4 (2.4%) subjects developed infections, although only one of these infections was at the site of the study graft.

Additionally, two trials from over a decade ago addressing the use of EZ Derm for the treatment of burns have been published (Healy, 1989; Vanstraelen, 1992). The Vanstraelen study concluded that hypertrophic scarring occurred in 25% of xenograft-dressed sites, but none was seen in the comparison group. Several allergic reactions were reported to the porcine xenograft. The conclusions of the Healy study found, in comparison to burned participants treated with Jelonet®, individuals treated with EZ Derm did not vary significantly in terms of bacterial colonization rate, need for surgical treatment, time for spontaneous healing, analgesic requirements or frequency of dressing changes.

Based on clinical practice standards, the above-mentioned studies, and the overall clinical experience with DermaMatrix, an acceptable level of safety and efficacy has been established for the use of this product for the treatment of burns.

Integra Dermal Regeneration Template

A 2013 study compared Integra with split-thickness skin graft and with viscose cellulose sponge (Cellonex), using 3, 10'5 cm test sites on each of 10 burn patients. The surrounding burn area was covered with meshed autograft. Biopsies were taken from each site on days 3, 7, 14, and 21, and at months 3 and 12. The tissue samples were stained and examined for markers of inflammation and proliferation. The Vancouver Scar Scale was used to assess scars. At 12-month follow-up, the 3 methods resulted in similar clinical appearance, along with similar histologic and immunohistochemical findings.

Branski et al (2007) reported on a randomized trial that compared Integra with a standard autograft-allograft technique in 20 children with an average burn size of 73% TBSA (71% full-thickness burns). Once vascularized (about 14-21 days), the Silastic epidermis was stripped and replaced with thin (0.05-0.13 mm) epidermal autograft. There were no significant differences between the Integra group and controls in burn size (70% vs. 74% TBSA), mortality (40% vs. 30%), and hospital length of stay (41 vs. 39 days), all respectively. Long-term follow-up revealed a significant increase in bone mineral content and density (24 months) and improved scarring in terms of height, thickness, vascularity, and pigmentation (at 12 months and 18-24 months) in the Integra group. No differences were observed between groups in the time to first reconstructive procedure, cumulative reconstructive procedures required during 2 years, and cumulative operating room time required for these procedures. The authors concluded that Integra can be used for immediate wound coverage in children with severe burns without the associated risks of cadaver skin.

Heimbach et al (2003) reported on a multicenter (13 U.S. burn care facilities) post approval study involving 222 burn injury patients (36.5% TBSA; range, 1%-95%) who were treated with Integra Dermal Regeneration Template. Within 2 to 3 weeks, the dermal layer regenerated, and a thin epidermal autograft was placed over the wound. The incidence of infection was 16.3%. Mean take rate (absence of graft failure) of Integra was 76.2%; the median take rate was 98%. The mean take rate of epidermal autograft placed over Integra was 87.7%; the median take rate was 95%.

Hicks et al (2019) conducted a systematic review of Integra dermal regeneration template for the treatment of acute full thickness burns and burn reconstruction. A total of 72 studies with 1084 patients (4 RCTs, 4 comparative studies, 5 cohort studies, 2 case control studies, 24 case series, and 33 case reports) were included in the review. The majority of patients (74%) were treated with Integra for acute burns, and the remainder (26%) for burn reconstruction. The take of the skin substitute was 86% (range 0–100%) for acute burn injuries and 95% (range 0–100%) for reconstruction. The take of the split-thickness skin graft over the template was 90% for acute burn injuries and 93% for reconstruction. There

was high variability in reporting of outcomes, but studies generally supported satisfactory cosmetic results in patients who have insufficient autograft and improvement in range of motion in patients who were treated with Integra for burn reconstruction. There was an overall complication rate of 13%; primarily due to infection, graft loss, hematoma formation, and contracture.

An infection rate of 18% was noted in a systematic review of complication rates in 10 studies that used Integra dermal regeneration template for burns.

Omega3 Wound

Luze et al (2022) conducted a systematic review of the use of acellular fish skin grafts in burn wound management. The reviewers identified 5 studies of Omega3 Wound but no RCTs. The identified studies were preclinical (animal), case series, retrospective observational, and 1 small (N = 21) cohort study. The review authors concluded that while the approach is promising, large-cohort studies are needed.

ReCell Autologous Cell Harvesting Device

Two RCTs have evaluated Recell for deep dermal burns (Table 23).

In both studies, 2 similar areas with a burn injury in the same individual were randomized to the control or treatment intervention (i.e., all participants received both treatments). The studies differed in their populations, interventions, and outcome measures. In the earlier study, participants all had deep partial thickness burns, while in the 2019 study the population included individuals with mixed-depth, full thickness burns. Holmes 2018 was a head-to-head comparison of ReCell alone versus skin grafting alone, and Holmes et al (2019) compared ReCell in combination with skin grafting. In the earlier study, the primary effectiveness endpoints were the incidence of wound closure at 4 weeks and the incidence of complete donor site healing at 1 week. In the 2019 trial, the co-primary effectiveness endpoints were non-inferiority of the incidence of RECELL-treated site closure by week 8 when compared to the control, and the superiority of the 37% relative reduction in donor skin for the ReCell treatment when compared with the control.

Study results are detailed in Table 24 and limitations in Tables 25 and 26. Although the ReCell device was comparable to standard care on outcomes such as complete wound closure; confidence in the strength of the overall body of evidence is limited by individual study limitations and heterogeneity of populations, interventions, and outcome measures across studies. Additional RCT evidence in the intended use population is needed.

Table 23. Randomized Controlled Trials of ReCell for Thermal Burns- Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Holmes et al (2018) NCT01138917	US	9	2010-2015	Individuals ages 18 to 65 years, with acute, deep partial-thickness thermal burns from 1% to 20% TBSA that required autografting for definitive closure.	ReCell device N = 101	Meshed STSG Treatment N = 101

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Holmes et al (2019) NCT02380612	US	6	2015-2017	Individuals ages 5 years or older, with acute thermal burn involving 5% to 50% of TBSA that underwent autografting for definitive closure	ReCell device treatment applied over STSG N = 30	Meshed STSG Treatment Alone N = 30

STSG:: Split-thickness skin grafts; TBSA: total body surface area.

Table 24. Randomized Controlled Trials of ReCell for Thermal Burns- Results

Study	Wound Closure (95% re-epithelialization) at 4 weeks	Wound Closure (95% re-epithelialization) at 8 weeks	Complete donor site healing at 1 week (100% re-epithelialization)	Relative Reduction in Donor Skin	Pain (VAS)	Participant Satisfaction and Scar Assessment	Adverse Events (Incidence)
Holmes et al (2018) NCT01138917							
ReCell	81/83 (97.6%)		21.8%		NSD at 16 weeks (data in figure)	NSD in subject satisfaction with appearance or in scarring at 16, 24, and 52 weeks (data in figures)	Treatment site: 35.6% Donor site: 4.0%
STSG	83/83 (100%)		10.0%				Treatment site: 21.8% Donor site: 6.9%
Between-group difference	-2.4% (95% CI: -8.4% to 2.3%)		p = .04				Treatment site: p = .0013 Donor site: 6.9% p = .25
Holmes et al (2019) NCT02380612							
ReCell plus STSG	50%	24/26 (92%)		368 (SD 150) cm ²	NSD between groups in treatment	NSD in subject satisfaction with appearance	NSD between groups in pre-specified
STSG alone	48%	22/26 (85%)		264 (SD 119) cm ²			

Study	Wound Closure (95% re-epithelialization) at 4 weeks	Wound Closure (95% re-epithelialization) at 8 weeks	Complete donor site healing at 1 week (100% re-epithelialization)	Relative Reduction in Donor Skin	Pain (VAS)	Participant Satisfaction and Scar Assessment	Adverse Events (Incidence)
Between-group difference		-7.7% Upper limit of the 97.5% CI 6.4% (i.e. within the pre-defined non-inferiority margin (10%))		32%; p <.001	nt area pain from week 1 to week 12 or week 52	or in scar assessment at any time point	safety events 17 individuals (57%) experienced AEs at control and ReCell sites; 27% had mild AEs, 37% moderate AEs. 1 death, attributed to underlying condition

AE: adverse events; CI: confidence interval; NSD: no significant difference; SD: standard deviation; STSG: Split-thickness skin grafts; VAS: visual analog scale.

Table 25. Randomized Controlled Trials of ReCell for Thermal Burns- Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Holmes et al (2018) NCT01138917					
Holmes et al (2019) NCT02380612	2. Participants had mixed depth full-thickness burns			5. Unclear if 32% reduction in donor site skin is clinically meaningful	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 26. Randomized Controlled Trials of ReCell for Thermal Burns- Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Holmes et al (2018) NCT01138917				83/101 participants evaluated in modified per protocol analysis	noninferiority margin based on 90 subjects	
Holmes et al (2019) NCT02380612				26/30 participants evaluated in per protocol analysis		3. confidence intervals not reported

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

Fresh Frozen Unprocessed Allograft Skin for Burns (AlloSkin and TheraSkin)

The use of fresh, unfrozen, unprocessed skin allograft has been used as a treatment of serious burn injuries since the First World War and it has become an accepted standard therapy. There are several brands of fresh, frozen, unprocessed allograft, including AlloSkin and TheraSkin. These products are treated as human tissue for transplantation under the FDA's HCT/P process.

A small number of studies have been published in the peer-reviewed literature addressing the use of TheraSkin. Landsman and colleagues (2010) conducted a single-center, retrospective, uncontrolled case series study of 188 subjects with VSUs and diabetic foot ulcers. The authors used historical controls for comparison, many of which were derived from previously published RCTs. The follow-up time was 20 weeks. The authors reported that at the 12-week follow-up, 60.4% of diabetic ulcers and 60.7% of venous ulcers were closed. At 20 weeks, those numbers increased to 74.1% and 74.6%, respectively. Neither of these differences was statistically significant. The authors conclude that TheraSkin is "highly effective" for the treatment of both VSUs and diabetic foot ulcers. No superiority was found, and in the absence of the desired outcome, the authors proffer that TheraSkin is equivalent to the control treatment. However, such conclusions reflect an inappropriate interpretation of the data from this effectiveness trial, which was not initially designed to test for equivalency, but superiority. To answer the question of equivalency, the authors would have had to have used an equivalency or non-inferiority study design, which would have utilized a different initial hypothesis and different set of assumptions.

DiDomenico and others (2011) conducted a non-controlled comparative trial of TheraSkin compared to Apligraf involving 28 subjects with diabetic foot ulcers, 16 of whom received TheraSkin and 12 received Apligraf. At 12 weeks, 66.7% of the TheraSkin subjects and 41.3% of the Apligraf subjects had closed wounds. These numbers changed only slightly at 20 weeks, to 66.7% and 47.14% respectively. The authors concluded that TheraSkin was more efficacious in healing diabetic foot ulcers. However, it should be noted that randomization problems in this study resulted in uneven blocks of subject enrollment in the

two cohorts, and the small sample size was not sufficiently powered to conclude whether TheraSkin was more effective than Apligraf.

Sanders (2014) reported the results of an underpowered RCT involving 23 subjects with DFUs randomized to receive treatment with either Dermagraft (n=12) or TheraSkin (n=11). At 12 weeks follow-up, 7 (63.6%) subjects in the TheraSkin group and 4 (33.3%) in the Dermagraft group were healed ($p=0.0498$). At the end of the 20-week evaluation period, 90.91% of TheraSkin subjects vs. 66.67% of the Dermagraft subjects were healed ($p=0.4282$). Time to healing in the TheraSkin group was significantly shorter (8.9 weeks) than in the Dermagraft group (12.5 weeks) (log-rank test, $p=0.0323$). The authors noted that the results of this study are similar to previous outcomes reported using these treatment modalities (see above studies) and suggest that, after 12 weeks of care, DFUs managed with TheraSkin are approximately twice as likely to heal as DFUs managed with Dermagraft, with approximately half the number of grafts required. However, they are careful to comment that, "Research confirming these results with a larger sample size and in individuals with different types of wounds is warranted."

Towler (2018) reported the results of a prospective RCT comparing Apligraf (n=12) to TheraSkin (n=15) for the treatment of VSUs. The authors reported no statistical differences between groups with regard to time to complete healing at 12 or 20 weeks ($p=0.294$ and $p=0.569$, respectively). Additionally, no differences were noted between groups with regard to the number of grafts needed ($p=0.119$). No adverse events were reported for either group. The authors concluded that both products are safe and effective to treat VSUs. However, the study was limited by small sample size, lack of blinding and other methodological issues.

In 2018, Choi reported on the use of cadaver allograft (n=698) vs. conventional treatment (n=584) in subjects with burns involving > 30% body surface area. In both unmatched and propensity-matched subject groups, 90-day in-hospital mortality was significantly better in the allograft group vs. controls (31.7 vs. 39.7% for unmatched comparisons and 37.8 vs. 47.3% for propensity matched comparisons). Logistic regression analyses showed a significant association between cadaver skin allograft and lower 90-day in-hospital mortality in the propensity-matched groups (OR, 0.42).

Armstrong (2021) conducted a randomized, prospective, evaluator-blinded study which compared the response of 100 subjects with non-healing DFU's, 50 of which were treated with TheraSkin and 50 treated with SOC. A total of 23 subjects withdrew from the trial, 4 in the TheraSkin group and 19 in the control group. In the TheraSkin group 1 was removed for not achieving > 50% area reduction by 6 weeks, 1 for wound worsening, and 2 due to adverse events, 1 potentially related to the study treatment and the other not related. In the control group, 11 subjects were removed for not achieving > 50% area reduction by 6 weeks, 1 due to reopened wound, 3 due to serious adverse events, 2 which were potentially treatment related, and 4 due to adverse events, 1 of which was possibly related to study treatment. In the ITT analysis the results at 12 weeks showed that 76% (38/50) of the TheraSkin-treated DFUs healed compared to 36% (18/50) of controls treated with SOC (adjusted $p=0.00056$). The mean percent area reduction at 12 weeks was 77.8% in the TheraSkin group vs. 49.6% in the SOC group (adjusted $p=0.0019$). The average time for closure within the 12-week period was 46.9 days for the TheraSkin group vs. 65.3 days for controls ($p=0.0019$). The authors concluded that wounds treated with TheraSkin in addition to SOC improved wound healing compared to SOC alone.

The use of fresh, unfrozen, unprocessed skin allograft products has been a part of standard medical practice for the treatment of burns for almost a century. However, concerns regarding the risk of disease transmission and shelf life continue to be an issue, and other products have been proposed as an alternative. One of the most commonly used alternative products is fresh frozen skin allograft. Unfortunately, the current level of evidence addressing the safety and efficacy of fresh frozen skin allograft is weak. No solid conclusions can be made regarding the superiority, equivalency, or inferiority of these types of products in relation to other treatment options. However, despite this lack of evidence, a decades-long anecdotal track record for these products, easy access and availability, and a higher degree

of certainty that the product is free from communicable pathogens has led to their acceptance as the standard of care in the burn treatment community. The use of fresh frozen skin allograft beyond the treatment of burns has not been established and is not widely accepted by the practicing community.

Trancyte

TransCyte was granted premarket approval (PMA) by the FDA in 1997 for "for use as a temporary wound covering for surgically excised full-thickness and deep partial-thickness thermal burn wounds in patients who require such a covering prior to autograft placement." TranCyte was not indicated for chronic wounds. TransCyte consists of human dermal fibroblasts grown on nylon mesh, combined with a synthetic epidermal layer. TransCyte can be used as a temporary covering over full thickness and some partial-thickness burns until autografting is possible. It can also be used as a temporary covering for some burn wounds that heal without autografting.

Section Summary: Deep Dermal Burns

Epicel is FDA-approved under a humanitarian device exemption (HDE) for the treatment of deep dermal or full-thickness burns comprising a TBSA of 30% or more, with patient survival of 90%. Integra Dermal Regeneration Template has been compared with autograft in a within-subject study and with autograft-allograft in a small RCT with 10 patients per group. Outcomes are at least as good as with autograft or allograft, with a reduction in scarring and without risks associated with cadaver skin. This product has also been studied in a large series with over 222 burn patients, showing a take rate of 76% and with a take rate of epidermal autograft placed over Integra of 87.7%.

The ReCell device has been evaluated in 2 RCTs. One RCT evaluated ReCell as an adjunct to meshed autologous skin grafting and the other compared ReCell head-to-head with skin grafting. Although the ReCell device was comparable to standard care on outcomes such as complete wound closure, confidence in the strength of the overall body of evidence is limited by individual study limitations and heterogeneity of populations, interventions, and outcome measures across studies. Additional RCT evidence in the intended use population is needed.

The use of for Biobrane, EZ Derm, Integra, Alloskin, Theraskin and Trancyte have been found to be efficacious in the treatment of full-thickness or deep partial-thickness burns based on current evidence and standard of care practices.

Other Indications

Dystrophic Epidermolysis Bullosa

OrCel was approved under an HDE (see Medical Policy [10.01.14 Humanitarian Use Devices](#)) for use in patients with dystrophic epidermolysis bullosa undergoing hand reconstruction surgery, to close and heal wounds created by the surgery, including those at donor sites. HDE status has been withdrawn for Dermagraft for this indication.

Fivenson et al (2003) reported the off-label use of Apligraf in 5 patients with recessive dystrophic epidermolysis bullosa who underwent syndactyly release.

Section Summary: Dystrophic Epidermolysis Bullosa

Dystrophic epidermolysis bullosa is a rare disorder. Because this is a rare disorder, it is unlikely that RCTs will be conducted to evaluate whether OrCel improves health outcomes for this condition.

Punch Biopsy Wounds

Baldursson et al (2015) reported a double-blinded RCT with 81 patients (162 punch biopsy wounds) that compared Kerecis Omega3 Wound (derived from fish skin) with Oasis SIS ECM (porcine small intestinal submucosa extracellular matrix). The primary outcome (the percentage of wounds healed at 28 days) was similar for the fish skin ADM (95%) and the porcine SIS ECM (96.3%). The rate of healing was faster with Kerecis Omega3 ($p=.041$). At 21 days, 72.5% of the fish skin ADM group had healed compared with 56% of the porcine SIS ECM group. Interpretation of this study is limited because it did not include an accepted control condition for this indication.

Split-Thickness Donor Sites

There is limited evidence to support the efficacy of OrCel compared with SOC for the treatment of split-thickness donor sites in burn patients. Still et al (2003) examined the safety and efficacy of bilayered OrCel to facilitate wound closure of split-thickness donor sites in 82 severely burned patients. Each patient had 2 designated donor sites that were randomized to a single treatment of OrCel or standard dressing (Biobrane-L). The healing time for OrCel sites was significantly shorter than for sites treated with a standard dressing, enabling earlier recropping. OrCel sites also exhibited a nonsignificant trend for reduced scarring. Additional studies are needed to evaluate the effect of this product on health outcomes.

Pressure Ulcers

Brown-Etris et al (2019) reported an RCT of 130 patients with stage 3 or stage 4 pressure ulcers who were treated with Oasis Wound Matrix (extracellular collagen matrix derived from porcine small intestinal submucosa) plus SOC or SOC alone. At 12 weeks, the proportion of wounds healed in the collagen matrix group was 40% compared to 29% in the SOC group. This was not statistically significant ($p=.111$). There was a statistical difference in the proportion of patients who achieved 90% wound healing (55% vs. 38% $p=.037$), but complete wound healing is the preferred and most reliable measure. It is possible that longer follow-up may have identified a significant improvement in the percent of wounds healed. The study did include 6-month follow-up, but there was high loss to follow-up and an insufficient number of patients at this time point for statistical comparison.

In the propensity matched study by Gurtner et al (2020) described above, Theraskin improved the healing rate of pressure ulcers by 20% (66.7% vs 46.8%).

Peripheral Nerve Injuries

The Cochrane Collaboration published a meta-analysis of bioengineered nerve conduits and wraps for repairs of peripheral nerves of the upper extremity. The authors included only RCTs or quasi-RCT experimental studies and found 5 which included the desired interventions and had follow-up periods of at least 12 months. A total of 213 participants were included in the studies, which compared nerve reconstruction with artificial wraps or conduits to standard repair either with direct end-to-end epineural repair or with autologous nerve grafting. Sensory recovery assessed with the British Medical Research Council (BMRC) grading scale was higher in the wrap or conduit group than in standard repair with very low certainty of evidence on Grading of Recommendations, Assessment, Development, and Evaluations

(GRADE) at 12 months (mean difference [MD], .03; range, -0.43 to 0.49) and 24 months follow-up (MD, .01; 95% CI, -0.06 to 0.08). Rosen model instrument (RMI) comparisons between conduit or wrap versus standard repair revealed no between-group differences through 24 months (MD, -0.17; 95% CI, -0.38 to 0.05; $p=.13$) and was determined to have low certainty of evidence; findings at 5 years follow-up in a single study found a greater improvement in the conduit or wrap group, but the estimate also had low certainty of evidence (MD, 0.23; 95% CI, 0.07 to 0.38). The rate of adverse event occurrence may be greater in patients treated with nerve wraps or conduits than with standard techniques, but the evidence had a GRADE rating reflected a very low certainty of evidence (risk ratio [RR], 7.15; 95% CI, 1.74 to 29.42). The authors also sought BMRC muscle strength scores, which were not reported in the included studies. The authors concluded that based on the currently available high-quality evidence, the use of currently available nerve repair devices is not supported over the standard of care due to heterogeneity in included participants, the pattern of injury, timing of repair, timing of outcome assessment, and choice of outcome measurement scales. A limitation of this systematic review is that they did not explicitly separate studies by the use of nerve conduits versus wraps for further analysis.

Miscellaneous

In addition to indications previously reviewed, off-label uses of bioengineered skin substitutes have included inflammatory ulcers (e.g., pyoderma gangrenosum, vasculitis), scleroderma digital ulcers, post keloid removal wounds, genetic conditions, and variety of other conditions. Products that have been FDA-approved or -cleared for one indication (e.g., lower-extremity ulcers) have also been used off-label in place of other FDA-approved or -cleared products (e.g., for burns). No RCTs were identified for these miscellaneous indications.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

National Institute for Health and Care Excellence

In 2023, NICE updated its guidance on the prevention and management of diabetic foot problems. The Institute recommended that clinicians “consider dermal or skin substitutes as an adjunct to standard care when treating diabetic foot ulcers, only when healing has not progressed and on the advice of the multidisciplinary foot care service.”

In 2019, NICE published guidance on the ReCell system for treating skin loss, scarring, and depigmentation after burn injury. The guidance recommended that additional research was needed to address the uncertainties regarding the potential benefits of ReCell.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review can be located at clinicaltrials.gov.

REFERENCES

1. Snyder DL, Sullivan N, Margolis DJ, Schoelles K. Skin substitutes for treating chronic wounds. Technology Assessment Program Project ID No. WNDT0818. (Prepared by the ECRI Institute-Penn Medicine Evidence-based Practice Center under Contract No. HHS A 290-2015-00005-I) Rockville, MD: Agency for Healthcare Research and Quality. February 2020. Available at: https://effectivehealthcare.ahrq.gov/sites/default/files/pdf/skin-substitute_0.pdf . Accessed January 5, 2023.
2. U.S. Food and Drug Administration. Regulatory Considerations for Human Cells, Tissues, and Cellular and Tissue-Based Products: Minimal Manipulation and Homologous Use. December 2017. <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/regulatory-considerations-human-cells-tissues-and-cellular-and-tissue-based-products-minimal> . Accessed December 22, 2022.
3. U.S. Food and Drug Administration. Executive Summary Breast Implant Special Topics. March 2019. <https://www.fda.gov/media/122956/download> . Accessed December 23, 2022.
4. U.S. Food & Drug Administration. Acellular Dermal Matrix (ADM) Products Used in Implant-Based Breast Reconstruction Differ in Complication Rates: FDA Safety Communication. March 2021. <https://www.fda.gov/medical-devices/safety-communications/acellular-dermal-matrix-adm-products-used-implant-based-breast-reconstruction-differ-complication> . Accessed December 27, 2022.
5. Davila AA, Seth AK, Wang E, et al. Human Acellular Dermis versus Submuscular Tissue Expander Breast Reconstruction: A Multivariate Analysis of Short-Term Complications. Arch Plast Surg. Jan 2013; 40(1): 19-27. PMID 23362476
6. Lee KT, Mun GH. Updated Evidence of Acellular Dermal Matrix Use for Implant-Based Breast Reconstruction: A Meta-analysis. Ann Surg Oncol. Feb 2016; 23(2): 600-10. PMID 26438439
7. McCarthy CM, Lee CN, Halvorson EG, et al. The use of acellular dermal matrices in two-stage expander/implant reconstruction: a multicenter, blinded, randomized controlled trial. Plast Reconstr Surg. Nov 2012; 130(5 Suppl 2): 57S-66S. PMID 23096987
8. Hinchcliff KM, Orbay H, Busse BK, et al. Comparison of two cadaveric acellular dermal matrices for immediate breast reconstruction: A prospective randomized trial. J Plast Reconstr Aesthet Surg. May 2017; 70(5): 568-576. PMID 28341592
9. Mendenhall SD, Anderson LA, Ying J, et al. The BREASTrial Stage II: ADM Breast Reconstruction Outcomes from Definitive Reconstruction to 3 Months Postoperative. Plast Reconstr Surg Glob Open. Jan 2017; 5(1): e1209. PMID 28203509
10. Liu DZ, Mathes DW, Neligan PC, et al. Comparison of outcomes using AlloDerm versus FlexHD for implant-based breast reconstruction. Ann Plast Surg. May 2014; 72(5): 503-7. PMID 23636114
11. Chang EI, Liu J. Prospective unbiased experience with three acellular dermal matrices in breast reconstruction. J Surg Oncol. Sep 2017; 116(3): 365-370. PMID 28444764

12. Pittman TA, Fan KL, Knapp A, et al. Comparison of Different Acellular Dermal Matrices in Breast Reconstruction: The 50/50 Study. *Plast Reconstr Surg*. Mar 2017; 139(3): 521-528. PMID 28234811
13. Dikmans RE, Negenborn VL, Bouman MB, et al. Two-stage implant-based breast reconstruction compared with immediate one-stage implant-based breast reconstruction augmented with an acellular dermal matrix: an open-label, phase 4, multicentre, randomised, controlled trial. *Lancet Oncol*. Feb 2017; 18(2): 251-258. PMID 28012977
14. Barber FA, Burns JP, Deutsch A, et al. A prospective, randomized evaluation of acellular human dermal matrix augmentation for arthroscopic rotator cuff repair. *Arthroscopy*. Jan 2012; 28(1): 8-15. PMID 21978432
15. Rashid MS, Smith RDJ, Nagra N, et al. Rotator cuff repair with biological graft augmentation causes adverse tissue outcomes. *Acta Orthop*. Dec 2020; 91(6): 782-788. PMID 32691656
16. Bellows CF, Smith A, Malsbury J, et al. Repair of incisional hernias with biological prosthesis: a systematic review of current evidence. *Am J Surg*. Jan 2013; 205(1): 85-101. PMID 22867726
17. Espinosa-de-los-Monteros A, de la Torre JI, Marrero I, et al. Utilization of human cadaveric acellular dermis for abdominal hernia reconstruction. *Ann Plast Surg*. Mar 2007; 58(3): 264-7. PMID 17471129
18. Gupta A, Zahriya K, Mullens PL, et al. Ventral herniorrhaphy: experience with two different biosynthetic mesh materials, Surgisis and Alloderm. *Hernia*. Oct 2006; 10(5): 419-25. PMID 16924395
19. Bochicchio GV, De Castro GP, Bochicchio KM, et al. Comparison study of acellular dermal matrices in complicated hernia surgery. *J Am Coll Surg*. Oct 2013; 217(4): 606-13. PMID 23973102
20. Roth JS, Zachem A, Plymale MA, et al. Complex Ventral Hernia Repair with Acellular Dermal Matrices: Clinical and Quality of Life Outcomes. *Am Surg*. Feb 01 2017; 83(2): 141-147. PMID 28228200
21. Bellows CF, Shadduck P, Helton WS, et al. Early report of a randomized comparative clinical trial of Strattice™ reconstructive tissue matrix to lightweight synthetic mesh in the repair of inguinal hernias. *Hernia*. Apr 2014; 18(2): 221-30. PMID 23543334
22. Fleshman JW, Beck DE, Hyman N, et al. A prospective, multicenter, randomized, controlled study of non-cross-linked porcine acellular dermal matrix fascial sublay for parastomal reinforcement in patients undergoing surgery for permanent abdominal wall ostomies. *Dis Colon Rectum*. May 2014; 57(5): 623-31. PMID 24819103
23. Kalaiselvan R, Carlson GL, Hayes S, et al. Recurrent intestinal fistulation after porcine acellular dermal matrix reinforcement in enteric fistula takedown and simultaneous abdominal wall reconstruction. *Hernia*. Jun 2020; 24(3): 537-543. PMID 31811593
24. Santema TB, Poyck PP, Ubbink DT. Skin grafting and tissue replacement for treating foot ulcers in people with diabetes. *Cochrane Database Syst Rev*. Feb 11 2016; 2(2): CD011255. PMID 26866804
25. Martinson M, Martinson N. A comparative analysis of skin substitutes used in the management of diabetic foot ulcers. *J Wound Care*. Oct 01 2016; 25(Sup10): S8-S17. PMID 27681811

26. Guo X, Mu D, Gao F. Efficacy and safety of acellular dermal matrix in diabetic foot ulcer treatment: A systematic review and meta-analysis. *Int J Surg.* Apr 2017; 40: 1-7. PMID 28232031
27. Veves A, Falanga V, Armstrong DG, et al. Graftskin, a human skin equivalent, is effective in the management of noninfected neuropathic diabetic foot ulcers: a prospective randomized multicenter clinical trial. *Diabetes Care.* Feb 2001; 24(2): 290-5. PMID 11213881
28. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Graftskin for the treatment of skin ulcers. *TEC Assessments.* 2001;Volume 16:Tab 12.
29. Steinberg JS, Edmonds M, Hurley DP, et al. Confirmatory data from EU study supports Apligraf for the treatment of neuropathic diabetic foot ulcers. *J Am Podiatr Med Assoc.* 2010; 100(1): 73-7. PMID 20093548
30. Kirsner RS, Warriner R, Michela M, et al. Advanced biological therapies for diabetic foot ulcers. *Arch Dermatol.* Aug 2010; 146(8): 857-62. PMID 20713816
31. Marston WA, Hanft J, Norwood P, et al. The efficacy and safety of Dermagraft in improving the healing of chronic diabetic foot ulcers: results of a prospective randomized trial. *Diabetes Care.* Jun 2003; 26(6): 1701-5. PMID 12766097
32. Frykberg RG, Marston WA, Cardinal M. The incidence of lower-extremity amputation and bone resection in diabetic foot ulcer patients treated with a human fibroblast-derived dermal substitute. *Adv Skin Wound Care.* Jan 2015; 28(1): 17-20. PMID 25407083
33. Zelen CM, Orgill DP, Serena T, et al. A prospective, randomized, controlled, multicenter clinical trial examining healing rates, safety and cost to closure of an acellular reticular allogenic human dermis versus standard of care in the treatment of chronic diabetic foot ulcers. *Int Wound J.* Apr 2017; 14(2): 307-315. PMID 27073000
34. Zelen CM, Orgill DP, Serena TE, et al. An aseptically processed, acellular, reticular, allogenic human dermis improves healing in diabetic foot ulcers: A prospective, randomized, controlled, multicentre follow-up trial. *Int Wound J.* Oct 2018; 15(5): 731-739. PMID 29682897
35. Driver VR, Lavery LA, Reyzelman AM, et al. A clinical trial of Integra Template for diabetic foot ulcer treatment. *Wound Repair Regen.* 2015; 23(6): 891-900. PMID 26297933
36. Campitiello F, Mancone M, Della Corte A, et al. To evaluate the efficacy of an acellular Flowable matrix in comparison with a wet dressing for the treatment of patients with diabetic foot ulcers: a randomized clinical trial. *Updates Surg.* Dec 2017; 69(4): 523-529. PMID 28497218
37. Brigido SA, Boc SF, Lopez RC. Effective management of major lower extremity wounds using an acellular regenerative tissue matrix: a pilot study. *Orthopedics.* Jan 2004; 27(1 Suppl): s145-9. PMID 14763548
38. Reyzelman A, Crews RT, Moore JC, et al. Clinical effectiveness of an acellular dermal regenerative tissue matrix compared to standard wound management in healing diabetic foot ulcers: a prospective, randomised, multicentre study. *Int Wound J.* Jun 2009; 6(3): 196-208. PMID 19368581
39. Reyzelman AM, Bazarov I. Human acellular dermal wound matrix for treatment of DFU: literature review and analysis. *J Wound Care.* Mar 2015; 24(3): 128; 129-34. PMID 25764957

40. Brigido SA. The use of an acellular dermal regenerative tissue matrix in the treatment of lower extremity wounds: a prospective 16-week pilot study. *Int Wound J.* Sep 2006; 3(3): 181-7. PMID 16984575
41. Walters J, Cazzell S, Pham H, et al. Healing Rates in a Multicenter Assessment of a Sterile, Room Temperature, Acellular Dermal Matrix Versus Conventional Care Wound Management and an Active Comparator in the Treatment of Full-Thickness Diabetic Foot Ulcers. *Eplasty.* 2016; 16: e10. PMID 26933467
42. Cazzell S, Vayser D, Pham H, et al. A randomized clinical trial of a human acellular dermal matrix demonstrated superior healing rates for chronic diabetic foot ulcers over conventional care and an active acellular dermal matrix comparator. *Wound Repair Regen.* May 2017; 25(3): 483-497. PMID 28544150
43. Gurtner GC, Garcia AD, Bakewell K, et al. A retrospective matched-cohort study of 3994 lower extremity wounds of multiple etiologies across 644 institutions comparing a bioactive human skin allograft, TheraSkin, plus standard of care, to standard of care alone. *Int Wound J.* Feb 2020; 17(1): 55-64. PMID 31729833
44. Sanders L, Landsman AS, Landsman A, et al. A prospective, multicenter, randomized, controlled clinical trial comparing a bioengineered skin substitute to a human skin allograft. *Ostomy Wound Manage.* Sep 2014; 60(9): 26-38. PMID 25211605
45. DiDomenico L, Landsman AR, Emch KJ, et al. A prospective comparison of diabetic foot ulcers treated with either a cryopreserved skin allograft or a bioengineered skin substitute. *Wounds.* Jul 2011; 23(7): 184-9. PMID 25879172
46. Frykberg RG, Cazzell SM, Arroyo-Rivera J, et al. Evaluation of tissue engineering products for the management of neuropathic diabetic foot ulcers: an interim analysis. *J Wound Care.* Jul 2016; 25 Suppl 7: S18-25. PMID 27410467
47. Lantis JC, Snyder R, Reyzelman AM, et al. Fetal bovine acellular dermal matrix for the closure of diabetic foot ulcers: a prospective randomised controlled trial. *J Wound Care.* Jul 01 2021; 30(Sup7): S18-S27. PMID 34256588
48. Kavros SJ, Dutra T, Gonzalez-Cruz R, et al. The use of PriMatrix, a fetal bovine acellular dermal matrix, in healing chronic diabetic foot ulcers: a prospective multicenter study. *Adv Skin Wound Care.* Aug 2014; 27(8): 356-62. PMID 25033310
49. Karr JC. Retrospective comparison of diabetic foot ulcer and venous stasis ulcer healing outcome between a dermal repair scaffold (PriMatrix) and a bilayered living cell therapy (Apligraf). *Adv Skin Wound Care.* Mar 2011; 24(3): 119-25. PMID 21326023
50. Niezgoda JA, Van Gils CC, Frykberg RG, et al. Randomized clinical trial comparing OASIS Wound Matrix to Regranex Gel for diabetic ulcers. *Adv Skin Wound Care.* Jun 2005; 18(5 Pt 1): 258-66. PMID 15942317
51. Uccioli L, Giurato L, Ruotolo V, et al. Two-step autologous grafting using HYAFF scaffolds in treating difficult diabetic foot ulcers: results of a multicenter, randomized controlled clinical trial with long-term follow-up. *Int J Low Extrem Wounds.* Jun 2011; 10(2): 80-5. PMID 21693443
52. Lullove EJ, Liden B, Winters C, et al. A Multicenter, Blinded, Randomized Controlled Clinical Trial Evaluating the Effect of Omega-3-Rich Fish Skin in the Treatment of Chronic, Nonresponsive Diabetic Foot Ulcers. *Wounds.* Jul 2021; 33(7): 169-177. PMID 33872197
53. Lullove EJ, Liden B, McEneaney P, et al. Evaluating the effect of omega-3-rich fish skin in the treatment of chronic, nonresponsive diabetic foot ulcers: penultimate analysis of a

- multicenter, prospective, randomized controlled trial. *Wounds*. Apr 2022; 34(4): E34-E36. PMID 35797557
54. O'Meara S, Cullum N, Nelson EA, et al. Compression for venous leg ulcers. *Cochrane Database Syst Rev*. Nov 14 2012; 11(11): CD000265. PMID 23152202
 55. Falanga V, Margolis D, Alvarez O, et al. Rapid healing of venous ulcers and lack of clinical rejection with an allogeneic cultured human skin equivalent. *Human Skin Equivalent Investigators Group. Arch Dermatol*. Mar 1998; 134(3): 293-300. PMID 9521027
 56. Mostow EN, Haraway GD, Dalsing M, et al. Effectiveness of an extracellular matrix graft (OASIS Wound Matrix) in the treatment of chronic leg ulcers: a randomized clinical trial. *J Vasc Surg*. May 2005; 41(5): 837-43. PMID 15886669
 57. Romanelli M, Dini V, Bertone M, et al. OASIS wound matrix versus Hyaloskin in the treatment of difficult-to-heal wounds of mixed arterial/venous aetiology. *Int Wound J*. Mar 2007; 4(1): 3-7. PMID 17425543
 58. Romanelli M, Dini V, Bertone MS. Randomized comparison of OASIS wound matrix versus moist wound dressing in the treatment of difficult-to-heal wounds of mixed arterial/venous etiology. *Adv Skin Wound Care*. Jan 2010; 23(1): 34-8. PMID 20101114
 59. Harding K, Sumner M, Cardinal M. A prospective, multicentre, randomised controlled study of human fibroblast-derived dermal substitute (Dermagraft) in patients with venous leg ulcers. *Int Wound J*. Apr 2013; 10(2): 132-7. PMID 23506344
 60. Cazzell S. A Randomized Controlled Trial Comparing a Human Acellular Dermal Matrix Versus Conventional Care for the Treatment of Venous Leg Ulcers. *Wounds*. Mar 2019; 31(3): 68-74. PMID 30720443
 61. Carsin H, Ainaud P, Le Bever H, et al. Cultured epithelial autografts in extensive burn coverage of severely traumatized patients: a five year single-center experience with 30 patients. *Burns*. Jun 2000; 26(4): 379-87. PMID 10751706
 62. Lagus H, Sarlomo-Rikala M, Böhling T, et al. Prospective study on burns treated with Integra®, a cellulose sponge and split thickness skin graft: comparative clinical and histological study--randomized controlled trial. *Burns*. Dec 2013; 39(8): 1577-87. PMID 23880091
 63. Branski LK, Herndon DN, Pereira C, et al. Longitudinal assessment of Integra in primary burn management: a randomized pediatric clinical trial. *Crit Care Med*. Nov 2007; 35(11): 2615-23. PMID 17828040
 64. Heimbach DM, Warden GD, Luterman A, et al. Multicenter postapproval clinical trial of Integra dermal regeneration template for burn treatment. *J Burn Care Rehabil*. 2003; 24(1): 42-8. PMID 12543990
 65. Hicks KE, Huynh MN, Jeschke M, et al. Dermal regenerative matrix use in burn patients: A systematic review. *J Plast Reconstr Aesthet Surg*. Nov 2019; 72(11): 1741-1751. PMID 31492583
 66. Gonzalez SR, Wolter KG, Yuen JC. Infectious Complications Associated with the Use of Integra: A Systematic Review of the Literature. *Plast Reconstr Surg Glob Open*. Jul 2020; 8(7): e2869. PMID 32802634
 67. Luze H, Nischwitz SP, Smolle C, et al. The Use of Acellular Fish Skin Grafts in Burn Wound Management-A Systematic Review. *Medicina (Kaunas)*. Jul 09 2022; 58(7). PMID 35888631

68. Holmes JH, Molnar JA, Shupp JW, et al. Demonstration of the safety and effectiveness of the RECELL ® System combined with split-thickness meshed autografts for the reduction of donor skin to treat mixed-depth burn injuries. *Burns*. Jun 2019; 45(4): 772-782. PMID 30578048
69. Holmes JH, Molnar JA, Carter JE, et al. A Comparative Study of the ReCell® Device and Autologous Split-Thickness Meshed Skin Graft in the Treatment of Acute Burn Injuries. *J Burn Care Res*. Aug 17 2018; 39(5): 694-702. PMID 29800234
70. Fivenson DP, Scherschun L, Cohen LV. Apligraf in the treatment of severe mitten deformity associated with recessive dystrophic epidermolysis bullosa. *Plast Reconstr Surg*. Aug 2003; 112(2): 584-8. PMID 12900618
71. Baldursson BT, Kjartansson H, Konráðsdóttir F, et al. Healing rate and autoimmune safety of full-thickness wounds treated with fish skin acellular dermal matrix versus porcine small-intestine submucosa: a noninferiority study. *Int J Low Extrem Wounds*. Mar 2015; 14(1): 37-43. PMID 25759413
72. Still J, Glat P, Silverstein P, et al. The use of a collagen sponge/living cell composite material to treat donor sites in burn patients. *Burns*. Dec 2003; 29(8): 837-41. PMID 14636761
73. Brown-Etris M, Milne CT, Hodde JP. An extracellular matrix graft (Oasis ® wound matrix) for treating full-thickness pressure ulcers: A randomized clinical trial. *J Tissue Viability*. Feb 2019; 28(1): 21-26. PMID 30509850
74. Lazic T, Falanga V. Bioengineered skin constructs and their use in wound healing. *Plast Reconstr Surg*. Jan 2011; 127 Suppl 1: 75S-90S. PMID 21200276
75. Saffle JR. Closure of the excised burn wound: temporary skin substitutes. *Clin Plast Surg*. Oct 2009; 36(4): 627-41. PMID 19793557
76. National Institute for Health and Care Excellence (NICE). Diabetic Foot Problems: Prevention and Management [NG19]. 2019; <https://www.nice.org.uk/guidance/ng19/evidence>.
77. Peirce SC, Carolan-Rees G. ReCell ® Spray-On Skin System for Treating Skin Loss, Scarring and Depigmentation after Burn Injury: A NICE Medical Technology Guidance. *Appl Health Econ Health Policy*. Apr 2019; 17(2): 131-141. PMID 30635844
78. Centers for Medicare & Medicaid Services (CMS). National Coverage Determination (NCD) for Porcine Skin and Gradient Pressure Dressings (270.5). n.d.; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=139&ncdver=1&bc>
79. Centers for Medicare & Medicaid Services (CMS). Fact Sheet: CMS finalizes Medicare Hospital Outpatient Prospective Payment System and Ambulatory Surgical Center Payment System changes for 2019 <https://www.cms.gov/newsroom/fact-sheets/cms-finalizes-medicare-hospital-outpatient-prospective-payment-system-and-ambulatory-surgical-center> .
80. Centers for Medicare & Medicaid Services. 2022. Fact Sheet. Calendar Year (CY) 2023 Medicare Physician Fee Schedule Final Rule. <https://www.cms.gov/newsroom/fact-sheets/calendar-year-cy-2023-medicare-physician-fee-schedule-final-rule> .
81. Hayes Inc. Medical Code Brief A2020 March 14, 2023
82. Hayes Inc. Medical Code Brief A2019 March 14, 2023

83. Sobti N, Liao EC. Surgeon-controlled study and meta-analysis comparing FlexHD and AlloDerm in immediate breast reconstruction outcomes. *Plast Reconstr Surg*. 2016;138(5):959-967
84. Garvey PB, Giordano SA, Baumann DP, et al. Long-term outcomes after abdominal wall reconstruction with acellular dermal matrix. *J Am Coll Surg*. 2017;224(3):341-350
85. Booth JH, Garvey PB, Baumann DP, et al. Primary fascial closure with mesh reinforcement is superior to bridged mesh repair for abdominal wall reconstruction. *J Am Coll Surg*. 2013;217(6):999-1009
86. Giordano S, Garvey PB, Baumann DP, et al. Primary fascial closure with biologic mesh reinforcement results in lesser complication and recurrence rates than bridged biologic mesh repair for abdominal wall reconstruction: A propensity score analysis. *Surgery*. 2017a;161(2):499-508
87. Giordano S, Schaverien M, Garvey PB, et al. Advanced age does not affect abdominal wall reconstruction outcomes using acellular dermal matrix: A comparative study using propensity score analysis. *Am J Surg*. 2017b;213(6):1046-1052
88. Mericli AF, Garvey PB, Giordano S, et al. Abdominal wall reconstruction with concomitant ostomy-associated hernia repair: Outcomes and propensity score analysis. *J Am Coll Surg*. 2017;224(3):351-361
89. Romain B, Story F, Meyer N, et al. Comparative study between biologic porcine dermal meshes: Risk factors of postoperative morbidity and recurrence. *J Wound Care*. 2016;25(6):320-325
90. Sbitany H, Kwon E, Chern H, et al. Outcomes analysis of biologic mesh use for abdominal wall reconstruction in clean-contaminated and contaminated ventral hernia repair. *Ann Plast Surg*. 2015;75(2):201-204
91. Fagotti L, Soares E, Bolia IK, et al. Early outcomes after arthroscopic hip capsular reconstruction using iliotibial band allograft versus dermal allograft. *Arthroscopy*. 2019;35(3):778-786
92. Li HY, Xiao SC, Zhu SH, et al. Successful treatment of a patient with an extraordinarily large deep burn. *Med Sci Monit*. 2011;17(4):CS47-CS51
93. Moravvej H, Hormozi AK, Hosseini SN, et al. Comparison of the application of allogeneic fibroblast and autologous mesh grafting with the conventional method in the treatment of third-degree burns. *J Burn Care Res*. 2016;37(1):e90-e95
94. Sivaraj D, Fischer KS, Kim TS, et al. Outcomes of biosynthetic and synthetic mesh in ventral hernia repair. *Plast Reconstr Surg Glob Open*. 2022; 10(12): e4707
95. Sivaraj D, Henn D, Fischer KS, et al. Reinforced biologic mesh reduces postoperative complications compared to biologic mesh after ventral hernia repair. *Plast Reconstr Surg Glob Open*. 2022; 10(2): e4083
96. Bain MA, Koullias GJ, Morse K, et al. Type I collagen matrix plus polyhexamethylene biguanide antimicrobial for the treatment of cutaneous wounds. *J Comp Eff Res*. 2020; 9(10):691-703
97. Koullias GJ, Bain MA, Thibodeaux K, Sabolinski M. A prospective noninterventional study of type I collagen matrix plus polyhexamethylene biguanide antimicrobial for the treatment of venous leg ulcers: a secondary analysis. *Wound Manag Prev*. 2022; 68(6):11-17

98. Lintzeris D, Vernon K, Percise H, et al. Effect of a new purified collagen matrix with polyhexamethylene biguanide on recalcitrant wounds of various etiologies: a case series. *Wounds*. 2018; 30(3):72-78
99. Menack MJ, Thibodeaux KT, Trabanco C, Sabolinski ML. Effectiveness of type I collagen matrix plus polyhexamethylene biguanide antimicrobial for the treatment of pressure injuries. *Wounds*. 2022; 34(6):159-164
100. Tierney BP. Comparison of 30-day clinical outcomes with SimpliDerm and AlloDerm RTU in immediate breast reconstruction. *Plast Reconstr Surg Glob Open*. 2021; 9(6): e3648
101. Tierney B P, De La Garza M, Jennings G R, et al. Clinical outcomes of acellular dermal matrix (SimpliDerm and AlloDerm Ready-to-Use) in immediate breast reconstruction. *Cureus*. 2022; 14(2): e22371
102. Gibson ALF, Holmes JH 4th, Shupp JW, et al. A phase 3, open-label, controlled, randomized, multicenter trial evaluating the efficacy and safety of StrataGraft® construct in patients with deep partial-thickness thermal burns. *Burns*. 2021; 47(5):1024-1037
103. Holmes JH 4th, Schurr MJ, King BT, et al. An open-label, prospective, randomized, controlled, multicenter, phase 1b study of StrataGraft skin tissue versus autografting in patients with deep partial-thickness thermal burns. *Burns*. 2019; pii: S0305-4179(19)30432-2
104. Kalstrup J, Balslev Willert C, Brinch-Møller Weitemeyer M, et al. Immediate direct-to-implant breast reconstruction with acellular dermal matrix: Evaluation of complications and safety. *Breast*. 2021; 60:192-198
105. Lohmander F, Lagergren J, Roy PG, et al. Implant based breast reconstruction with acellular dermal matrix: safety data from an open-label, multicenter, randomized, controlled trial in the setting of breast cancer treatment. *Ann Surg*. 2019; 269(5):836-841
106. Paprottka FJ, Krezdorn N, Sorg H, et al. Evaluation of complication rates after breast surgery using acellular dermal matrix: median follow-up of three years. *Plast Surg Int*. 2017; 2017:1283735
107. Wilson RL, Kirwan CC, O'Donoghue JM, et al. BROWSE: A multicentre comparison of nine year outcomes in acellular dermal matrix based and complete submuscular implant-based immediate breast reconstruction-aesthetics, capsular contracture and patient reported outcomes. *Eur J Surg Oncol*. 2021 Nov 6:S0748-7983(21)00776-9
108. Asaad M, Selber JC, Adelman DM, et al. Allograft vs xenograft bioprosthetic mesh in tissue expander breast reconstruction: a blinded prospective randomized controlled trial. *Aesthet Surg J*. 2021. 41(12):NP1931-NP1939
109. Barmettler A, Heo M. A Prospective, randomized comparison of lower eyelid retraction repair with autologous auricular cartilage, bovine acellular dermal matrix (SurgiMend), and porcine acellular dermal matrix (Enduragen) spacer grafts. *Ophthalmic Plast Reconstr Surg*. 2018; 34(3):266-273
110. Mazari FAK, Wattoo GM, Kazzazi NH, et al. The comparison of Stratitice and SurgiMend in acellular dermal matrix-assisted, implant-based immediate breast reconstruction. *Plast Reconstr Surg*. 2018; 141(2):283-293
111. Schefflan M, Grinberg-Rashi H, Hod K. Bovine acellular dermal matrix in immediate breast reconstruction: a retrospective, observational study with SurgiMend. *Plast Reconstr Surg*. 2018; 141(1):1e-10e

112. Alexandridis V, Teleman P, Rudnicki M. Efficacy and safety of pelvic organ prolapse surgery with porcine small intestinal submucosa graft implantation. *Eur J Obstet Gynecol Reprod Biol.* 2021; 267:18-22
113. Jayne DG, Scholefield J, Tolan D, et al. Anal fistula plug versus surgeon's preference for surgery for trans-sphincteric anal fistula: the FIAT RCT. *Health Technol Assess.* 2019; 23(21):1-76
114. Jayne DG, Scholefield J, Tolan D, et al.; FIAT Trial Collaborative Group. A multicenter randomized controlled trial comparing safety, efficacy, and cost-effectiveness of the Surgisis Anal Fistula Plug versus surgeon's preference for transsphincteric fistula-in-ano: The FIAT Trial. *Ann Surg.* 2021; 273(3):433-441
115. Korwar V, Adjepong S, Pattar J, Sigurdsson A. Biological mesh repair of paraesophageal hernia: an analysis of our outcomes. *J Laparoendosc Adv Surg Tech A.* 2019; (11):1446-1450
116. Lin H, Jin Z, Zhu Y, et al. Anal fistula plug vs rectal advancement flap for the treatment of complex cryptoglandular anal fistulas: a systematic review and meta-analysis of studies with long-term follow-up. *Colorectal Dis.* 2019; 21(5):502-515
117. Thomas PW, Blackwell JEM, Herrod PJJ, et al. Long-term outcomes of biological mesh repair following extra levator abdominoperineal excision of the rectum: an observational study of 100 patients. *Tech Coloproctol.* 2019; 23(8):761-767
118. Armstrong D, Galiano R, Orgill D, et al. Multi-centre prospective randomised controlled clinical trial to evaluate a bioactive split thickness skin allograft vs standard of care in the treatment of diabetic foot ulcers. *Int Wound J.* 2022; 19(4):932-944
119. Barbul A, Gurtner GC, Gordon H, et al. Matched-cohort study comparing bioactive human split-thickness skin allograft plus standard of care to standard of care alone in the treatment of diabetic ulcers: A retrospective analysis across 470 institutions [published correction appears in *Wound Repair Regen.* 2020;28(3):431]. *Wound Repair Regen.* 2020; 28(1):81-89
120. Gurtner GC, Garcia AD, Bakewell K, Alarcon JB. A retrospective matched-cohort study of 3994 lower extremity wounds of multiple etiologies across 644 institutions comparing a bioactive human skin allograft, TheraSkin, plus standard of care, to standard of care alone. *Int Wound J.* 2020; 17(1):55-64
121. Towler MA, Rush EW, Richardson MK, Williams CL. randomized, prospective, blinded-enrollment, head-to-head towler venous leg ulcer healing trial comparing living, bioengineered skin graft substitute (Apligraf) with living, cryopreserved, human skin allograft (TheraSkin). *Clin Podiatr Med Surg.* 2018; 35(3):357-365
122. Eichler C, Efremova J, Brunnert K, et al. A head to head comparison between SurgiMend® - fetal bovine acellular dermal matrix and Tutomesh® - a bovine pericardium collagen membrane in breast reconstruction in 45 cases. *In Vivo.* 2017; 31(4):677-682
123. Paprottka FJ, Krezdorn N, Sorg H, et al. Evaluation of complication rates after breast surgery using acellular dermal matrix: median follow-up of three years. *Plast Surg Int.* 2017; 2017:1283735
124. Quah GS, French JR, Cocco A, et al. Veritas in immediate implant-based breast reconstruction is associated with higher complications compared with tiLOOP. *Plast Reconstr Surg Glob Open.* 2019; 7(12):e2533

125. Li H, Lim P, Stanley E, et al. Experience with NovoSorb® Biodegradable Temporising Matrix in reconstruction of complex wounds. *ANZ J Surg.* 2021; 91(9):1744-1750
126. Lo CH, Brown JN, Dantzer EJG, et al. Wound healing and dermal regeneration in severe burn patients treated with NovoSorb® Biodegradable Temporising Matrix: A prospective clinical study. *Burns.* 2022 May;48(3):529-538. 2022; 48(3):529-538
127. Li H, Lim P, Stanley E, et al. Experience with NovoSorb® Biodegradable Temporising Matrix in reconstruction of complex wounds. *ANZ J Surg.* 2021; 91(9):1744-1750
128. Gould LJ, Orgill DP, Armstrong DG, et al. Improved healing of chronic diabetic foot wounds in a prospective randomised controlled multi-centre clinical trial with a microvascular tissue allograft. *Int Wound J.* 2022; 19(4):811-825
129. Kim TH, Park JH, Jeong HG, Wee SY. The utility of novel fish-skin derived acellular dermal matrix (Kerecis) as a wound dressing material. *Wound Manag Res.* 2021; 17(1):39-47
130. Kirsner RS, Margolis DJ, Baldursson BT, et al. Fish skin grafts compared to human amnion/chorion membrane allografts: a double-blind, prospective, randomized clinical trial of acute wound healing. *Wound Repair Regen.* 2020; 28(1):75-80
131. Lullove EJ, Liden B, Winters C, et al. A multicenter, blinded, randomized controlled clinical trial evaluating the effect of omega-3-rich fish skin in the treatment of chronic, nonresponsive diabetic foot ulcers. *Wounds.* 2021;33(7):169-177
132. Lullove EJ, Liden B, McEneaney P, et al. Evaluating the effect of omega-3-rich fish skin in the treatment of chronic, nonresponsive diabetic foot ulcers: penultimate analysis of a multicenter, prospective, randomized controlled trial. *Wounds.* 2022; 34(4): E34-E36
133. Lullove EJ, Liden B, McEneaney P, et al. Fish-skin grafts for the management of chronic diabetic foot ulcers: final efficacy and cost analysis of a prospective, multicenter, randomized, controlled clinical trial, and cost effectiveness versus a standardized care plan. *Wounds.* 2023; In press
134. Michael S, Winters C, Khan M. Acellular fish skin graft use for diabetic lower extremity wound healing: a retrospective study of 58 ulcerations and a literature review. *Wounds.* 2019; 31(10):262-268
135. Hicks CW, Zhang GQ, Canner JK, et al. Outcomes and predictors of wound healing among patients with complex diabetic foot wounds treated with a dermal regeneration template (Integra). *Plast Reconstr Surg.* 2020; 146(4):893-902
136. Othman S, Lukowiak T, Shakir S, et al. Bilayer wound matrix-based cutaneous scalp reconstruction: A multidisciplinary case control analysis of factors associated with reconstructive success and failure. *J Plast Reconstr Aesthet Surg.* 2021; 74(11):3008-3014
137. Wang CQ, Tran T, Montera B, K, et al. Symptomatic, radiological, and quality of life outcome of paraesophageal hernia repair with urinary bladder extracellular surgical matrix: comparison with primary repair. *Surg Laparosc Endosc Percutan Tech.* 2019; 29(3):182-186
138. Adams WP Jr, Baxter R, Glicksman C, et al. The use of poly-4-hydroxybutyrate (P4HB) scaffold in the ptotic breast: a multicenter clinical study. *Aesthet Surg J.* 2018; 38(5):502-518
139. Maxwell DW, Hart AM, Keifer OP Jr, et al. A Comparison of acellular dermal matrices in abdominal wall reconstruction. *Ann Plast Surg.* 2019; 82(4):435-440

140. Broyles JM, Liao EC, Kim J, et al. Acellular dermal matrix-associated complications in implant-based breast reconstruction: a multicenter, prospective, randomized controlled clinical trial comparing two human tissues. *Plast Reconstr Surg*. 2021; 148(3):493-500
141. Barmettler A, Heo M. A prospective, randomized comparison of lower eyelid retraction repair with autologous auricular cartilage, bovine acellular dermal matrix (SurgiMend), and porcine acellular dermal matrix (Enduragen) spacer grafts. *Ophthalmic Plast Reconstr Surg*. 2018; 34(3):266-273
142. Cazzell S, Moyer PM, Samsell B, et al. A prospective, multicenter, single-arm clinical trial for treatment of complex diabetic foot ulcers with deep exposure using acellular dermal matrix. *Adv Skin Wound Care*. 2019; 32(9):409-415
143. Cazzell S, Vayser D, Pham H, et al. A randomized clinical trial of a human acellular dermal matrix demonstrated superior healing rates for chronic diabetic foot ulcers over conventional care and an active acellular dermal matrix comparator. *Wound Repair Regen*. 2017; 25(3):483-497
144. Chang EI, Liu J. Prospective unbiased experience with three acellular dermal matrices in breast reconstruction. *J Surg Oncol*. 2017; 116(3):365-370
145. Ortiz JA. Clinical outcomes in breast reconstruction patients using a sterile acellular dermal matrix allograft. *Aesthetic Plast Surg*. 2017; 41(3):542-550
146. Pittman TA, Fan KL, Knapp A, et al. Comparison of different acellular dermal matrices in breast reconstruction: the 50/50 study. *Plast Reconstr Surg*. 2017; 139(3):521-528
147. Powers JM, Reuter Muñoz KD, Parkerson J, et al. From salvage to prevention: a single-surgeon experience with acellular dermal matrix and infection in prepectoral breast reconstruction. *Plast Reconstr Surg*. 2021; 148(6):1201-1208
148. Lindsey JT Jr, Boyd CJ, Davis CB, et al. Alloderm and Cortiva have similar perioperative wound complications in abdominal wall reconstruction. *J Surg Res*. 2020; 255:255-260
149. Moyer HR, Hart AM, Yeager J, Losken A. A histological comparison of two human acellular dermal matrix products in prosthetic-based breast reconstruction. *Plast Reconstr Surg Glob Open*. 2017; 5(12):e1576
150. Parikh RP, Tenenbaum MM, Yan Y, Myckatyn TM. Cortiva versus AlloDerm Ready-To-Use in prepectoral and submuscular breast reconstruction: prospective randomized clinical trial study design and early findings. *Plast Reconstr Surg Glob Open*. 2018; 6(11):e2013
151. Urquia LN, Hart AM, Liu DZ, Losken A. Surgical outcomes in prepectoral breast reconstruction. *Plast Reconstr Surg Glob Open*. 2020; 8(4):e2744
152. Ashfaq A, Brown T, Reemtsen B. Repair of complete atrioventricular septal defects with decellularized extracellular matrix: initial and midterm outcomes. *World J Pediatr Congenit Heart Surg*. 2017; 8(3):310-314
153. Hu K, Siddiqi U, Lee B, et al. Pediatric aortic valve repair: any development in the material for cusp extension valvuloplasty? *J Card Surg*. 2021; 36(11):4054-4060
154. Kelley TM Jr, Kashem M, Wang H, et al. Anterior leaflet augmentation with CorMatrix porcine extracellular matrix in twenty-five patients: unexpected patch failures and histologic analysis. *Ann Thorac Surg*. 2017; 103(1):114-120
155. Maillot C, Harly E, Demezou H, Le Huec JC. Surgical repair of large-to-massive rotator cuff tears seems to be a better option than patch augmentation or débridement and biceps

- tenotomy: a prospective comparative study. *J Shoulder Elbow Surg.* 2018; 27(9):1545-1552
156. Rogers AD, Blackport E, Cartotto R. The use of Biobrane® for wound coverage in Stevens-Johnson Syndrome and toxic epidermal necrolysis. *Burns.* 2017; 43(7):1464-1472
 157. Parikh RP, Brown G, Sharma K, et al. Immediate implant-based breast reconstruction with acellular dermal matrix: a comparison of sterile and aseptic AlloDerm in 2039 consecutive cases. *Plast Reconstr Surg.* 2018; 142(6):1401-1409
 158. Park SJ, Kim Y, Jang SY. The application of an acellular dermal allograft (AlloDerm) for patients with insufficient conjunctiva during evisceration and implantation surgery. *Eye (Lond).* 2018; 32(1):136-141
 159. Kim KY, Woo YJ, Jang SY, et al. Correction of lower eyelid retraction using acellular human dermis during orbital decompression. *Ophthalmic Plast Reconstr Surg.* 2017 ;33(3):168-172
 160. Woo A, Harless C, Jacobson SR. Revisiting an old place: single-surgeon experience on post-mastectomy subcutaneous implant-based breast reconstruction. *Breast J.* 2017; 23(5):545-553
 161. Choi YH, Cho YS, Lee JH, et al. Cadaver skin allograft may improve mortality rate for burns involving over 30% of total body surface area: a propensity score analysis of data from four burn centers. *Cell Tissue Bank.* 2018; 19(4):645-651
 162. UpToDate. Overview of the management of epidermolysis bullosa. Topic last updated November 2024. Also available at <https://www.uptodate.com>
 163. UpToDate. Implant-based breast reconstruction and augmentation. Topic last updated February 2025. Also available at <https://www.uptodate.com>
 164. UpToDate. Overview of breast reconstruction. Topic last updated May 2023. Also available at <https://www.uptodate.com>
 165. UpToDate. Management of the open abdomen in adults. Topic last updated October 2024. Also available at <https://www.uptodate.com>
 166. UpToDate. Skin Substitutes. Topic last updated. July 2025. Also available at <https://www.uptodate.com>
 167. UpToDate. Evaluation and management chronic venous insufficiency including venous leg ulcer. Topic last updated April 2026. Also available at <https://www.uptodate.com>
 168. UpToDate. Local care of diabetes-related foot ulcers/wounds. Topic last updated February 2026. Also available at <https://www.uptodate.com>
 169. Hayes, a symplr company. Health Technology Assessment Acellular Skin Substitute for Chronic Foot Ulcers in Adults with Diabetes Mellitus May 10, 2023
 170. Hayes, a symplr company. Evidence Analysis Research Brief Skin Substitutes of Acute Wounds January 31, 2024
 171. Hayes, a symplr company. Health Technology Assessment Skin Substitutes for Venous Leg Ulcers in Adults. July 23, 2020/Annual Review July 11, 2023
 172. Hayes, a symplr company. Evidence Analysis Research Brief. Bioengineered Skin Substitutes for Treatment of Dystrophic Epidermolysis Bullosa. December 17, 2025

173. Davison SP, Harbour S, Fassihi E. Comparison of Different Acellular Dermal Matrix in Breast Reconstruction: A Skin-to-Skin Study. *Aesthet Surg J.* Jul 15 2024; 44(8): 829-837. PMID 38377366
174. Keane AM, Chiang SN, Tao Y, et al. Cortiva versus AlloDerm in Prepectoral and Partial Submuscular Implant-Based Breast Reconstruction: A Randomized Clinical Trial. *Plast Reconstr Surg.* Oct 01 2024; 154(4S): 13S-26S. PMID 38085977
175. Ng TP, Loo BYK, Yong N, et al. Review: Implant-Based Breast Reconstruction After Mastectomy for Breast Cancer: A Meta-analysis of Randomized Controlled Trials and Prospective Studies Comparing Use of Acellular Dermal Matrix (ADM) Versus Without ADM. *Ann Surg Oncol.* May 2024; 31(5): 3366-3376. PMID 38285304
176. Thomson SE, Ng NY, Riehle MO, et al. Bioengineered nerve conduits and wraps for peripheral nerve repair of the upper limb. *Cochrane Database Syst Rev.* Dec 07 2022; 12(12): CD012574. PMID 36477774
177. Khamise A, Lapid H, Mishra A, et al. Acute healing outcomes in paediatric partial thickness burns using Epiprotect and Biobrane®: A retrospective comparative study. *J Plast Reconstr Aesthet Surg* 2024 Aug; 95:55-61. PMID 38875874
178. Lee A, McKean A, Wormald J, et. al. A clinician survey of skin substitute use in burns care in the UK and Ireland: The CASSIUS study. *J Plast Reconstr Aesthet Surg.* 2025 Mar;102:238-246. PMID 39938465
179. Delgado-Miguel C, Moran AG, Gomez LF, et. al. Comparison of the effectiveness of three different skin substitutes for the treatment of pediatric burns. *Eur J Pediatr* 2024 Dec 13;184(1):80. PMID 39671108
180. Samuels K, Millet E and Wong L. Efficacy of acellular dermal matrix type in the treatment of capsular contracture in breast augmentation: A systematic review and meta-analysis. *Aesthet Surg J.* 2023 Dec 14;44(1):26-35. PMID 37595294
181. Williams ML and Holewinski JE. Use of human acellular dermal matrix in patients with complex wounds and comorbidities. *J Wound Care.* 2015 Jun;24(6):261-2, 264-7. PMID 26075374
182. Armstrong DG, Orgill DP, Galiano RD, et al. A multi-centre, single-blinded randomised controlled clinical trial evaluating the effect of resorbable glass fibre matrix in the treatment of diabetic foot ulcers. *Int Wound J.* May 2022; 19(4): 791-801. PMID 34418302
183. Armstrong DG, Orgill DP, Galiano RD, et al. A Borate-Based Bioactive Glass Advances Wound Healing in Non-Healing Wagner Grade 1 Diabetic Foot Ulcers: A Randomised Controlled Clinical Trial. *Int Wound J.* Oct 2025; 22(10): e70763. PMID 41014175
184. Dardari D, Piaggese A, Potier L, et al. Intact Fish Skin Graft to Treat Deep Diabetic Foot Ulcers. *NEJM Evid.* Dec 2024; 3(12): EVIDoa2400171. PMID 39365895
185. Dhanani NH, Lyons NB, Lee KH, et al. Synthetic Versus Biologic Mesh for Complex Open Ventral Hernia Repair: 3-Year Follow-Up of a Pilot Randomized Controlled Trial. *HCA Healthc J Med.* 2025; 6(1): 43-50. PMID 40071184
186. Ruiz-Muñoz M, Martinez-Barrios FJ, Cervera-Garvi P, et al. Fish skin grafts versus standard of care on wound healing of chronic diabetic foot ulcers: A systematic review and meta-analysis. *Prim Care Diabetes.* Jun 2024; 18(3): 291-298. PMID 38644082

187. Seefeldt CS, Knievel J, Deeb M, et al. Biological vs Synthetic Mesh in Laparoendoscopic Inguinal Hernia Repair: The BIOLAP Randomized Clinical Trial. *JAMA Surg.* Dec 01 2025; 160(12): 1309-1316. PMID 41060645
188. U.S. Food and Drug Administration. General and Plastic Surgery Devices Advisory Committee Meeting. October 20, 2021. <https://www.fda.gov/media/153413/download>. Accessed May 2026
189. Remulla D, Carvalho A, Birrell AM, et al. Mesh-related Outcomes of Biologic Versus Synthetic Mesh for Single-stage Repair of Contaminated Ventral Hernias: A 5 to 10-year Analysis of a Randomized Controlled Trial. *Ann Surg.* Mar 01, 2026; 283(3): 390-397. PMID 40792628
190. Bruggeman A, Van Schouwen KF. Results of bioresorbable hydrogel sheet utilized as an adhesion barrier in spine surgery. *JSM Neurosurgery and Spine* 9(1):1104 April 2022
191. Hones KM, Nichols DS, Barker H et. al. Outcomes following use of VersaWrap nerve protector in treatment of patients with recurrent compression neuropathies. *Front. Surg.* 10:1123375. doi: 10.3389/fsurg.2022.1123375
192. McDermott E, Bowers Z, Nuelle J. The application of hyaluronic acid/Alginate flexor pollicis longus tendon repair to prevent adhesion formation: A second look. *Cureus.* 2023 doi:10.7759/cureus.33147
193. Adu Y, Harder J, Cox C, et. al. Evaluating the effect of VersaWrap tendon protector on functional outcomes in operative tendon repairs. *Front. Surg.* 11:1447515. Doi: 10.3389/fsurg.2024.1447515
194. Bridges T, McCahon J, Parhk S. Surgical arthroscopy with intra-articular hyaluronic acid/Alginate adjunct in the treatment of ankle osteoarthritis. *Tech Foot and Ankle* 2024 Volume 00 Number 00. DOI: 10.1097/BTF.0000000000000422
195. U.S. Food and Drug Administration. Alafair Biosciences Inc. FDA Approval VersaWrap October 29, 2021. K212163
196. Shi L and Ronfard V. Biochemical and biomechanical characterization of porcine small intestinal submucosa (SIS): a mini review. *Int J Burns Trauma.* 2013 Nov 1;3(4):173-9. PMID 24273692
197. Salgado R, Bravo L, Garcia M, et. al. Histomorphometric Analysis of early epithelialization and dermal changes in mid-partial-thickness burn wounds in humans treated with porcine small intestinal submucosa and silver-containing hydrofiber. *Journal of Burn Care & Research*, Volume 35, Issue 5, September-October 2014, Pages e330-e337, <https://doi.org/10.1097/BCR.0000000000000015>
198. Guesmi F, Saidi I, Abbassi R et al Therapeutic potential of second degree's skin burns by topical dressing of teucrium ramosissimum that promotes re-epithelialization. *Dermatol Ther.*2022 May;35(5):e15428. PMID 35261131
199. Betar N, Maher D, Wheatley L et al Clinical outcomes and resource utilization in patients with major burns treated with NovoSorb® BTM. *Burns.* 2023 Nov;49(7):1663-1669
200. Jacoby S, Bachoura A, Chen N, et. al. One-stage integra coverage for fingertip injuries. *Hand (NY).* 2013 Sep;8(3):291-5. Doi:10.1007/s11552-013-9513-x. PMID 24426936
201. Hamdan AM, Al-Chalabi MMM, Wan Sulaiman WA. Integra: An alternative option for reconstruction of extensive finger defects with exposed bones. *Cureus.* 2021 Jul 7; 13(7): e16223. doi: 10.7759/cureus.16223. PMID 34367823

202. Bracey J. Use of Integra in traumatic, nonburn injuries of the hand. Journal of Hand Surgery Volume 39, Issue 9, Supplement, E56, September 2014
203. Kahn S, Carter J, Wilde S, et. al. Autologous skin cell suspension for full-thickness skin defect reconstruction: Current evidence and health economic expectations. Adv Ther. 2024;41(3):891-900. doi: 10.1007/s12325-02302777-7.Epub 2024 Jan 22
204. Henry S, Mapula S, Grevious M, et. al. Maximizing wound coverage in full-thickness skin defects: A randomized-controlled trial of autologous skin cell suspension and widely meshed autograft versus standard autografting. J Trauma Acute Care Surg. 2024 Jan 1; 96(1):85-93. Doi: 10.1097/TA.0000000000001420. Epub 2023 Sep 1 PMID 38098145

CODES

To report provider services, use appropriate CPT codes, HCPCS codes, Revenue codes, and/or ICD diagnosis codes.

Codes	Number	Description
CPT		
		Harvesting and Application of SkinTE® Product
	1044T	Harvest of full-thickness for autologous heterogenous skin-construct graft, including direct closure of donor site; first 5 sq cm or less
	1045T	Harvest of full-thickness skin for autologous heterogenous skin-construct graft, including direct closure of donor site; each additional 5 sq cm, or part thereof (List separately in addition to code for primary procedure)
	1046T	Autologous heterogenous skin-construct graft application, trunk, arms, legs; first 50 sq cm or less, or 0.5% of body area of infants and children
	1047T	Autologous heterogenous skin-construct graft application, trunk, arms, legs; each additional 50 sq cm, or each additional 0.5% of body area of infants and children, or part thereof (List separately in addition to code for primary procedure)
	1048T	Autologous heterogenous skin-construct graft application, face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits; each additional 50 sq cm, or each additional 0.5% of body area of infants and children, or part thereof (List separately in addition to code for primary procedure)
	1049T	Autologous heterogenous skin-construct graft application, face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits; ach additional 50 sq cm, or each additional 0.5% of body area of infants and children, or part thereof (List separately in addition to code for primary procedure)

		Autologous Cell Harvesting Device (Recell®)
	15011	Harvest of skin for skin cell suspension autograft; first 25 sq cm or less
	15012	Harvest of skin for skin cell suspension autograft; each additional 25 sq cm or part thereof (List separately in addition to code for primary procedure)
	15013	Preparation of skin cell suspension autograft, requiring enzymatic processing, manual mechanical disaggregation of skin cells, and filtration; first 25 sq cm or less of harvested skin
	15014	Preparation of skin cell suspension autograft, requiring enzymatic processing, manual mechanical disaggregation of skin cells, and filtration; each additional 25 sq cm or less of harvested skin or part thereof (List separately in addition to code for primary procedure)
	15015	Application of skin cell suspension autograft to wound and donor sites, including application of primary dressing, trunk, arm, legs; first 480 sq cm or less
	15016	Application of skin cell suspension autograft to wound and donor sites, including application of primary dressing, trunk, arm, legs; each additional 480 sq cm or part thereof (List separately in addition to code for primary procedure)
	15017	Application of skin cell suspension autograft to wound and donor sites, including application of primary dressing, face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits; first 480 sq cm or less
	15018	Application of skin cell suspension autograft to wound and donor sites, including application of primary dressing, face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits; each additional 480 sq cm or part thereof (List separately in addition to code for primary procedure)
		Application Codes
	15150	Tissue cultured skin autograft, trunk, arms, legs; first 25 sq cm or less
	15151	Tissue cultured skin autograft, trunk, arms, legs; additional 1 sq cm to 75 sq cm
	15152	Tissue cultured skin autograft, trunk, arms, legs; each additional 100 sq cm, or each additional 1% of body area of infants and children, or part thereof
	15155	Tissue cultured skin autograft, face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits; first 25 sq cm or less

	15156	Tissue cultured skin autograft, face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits; additional 1 sq cm to 75 sq cm
	15157	Tissue cultured skin autograft, face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits; each additional 100 sq cm, or each additional 1% of body area of infants and children, or part thereof
	15271	Application of skin substitute graft to trunk, arms, legs, total wound surface area up to 100 sq cm; first 25 sq cm or less wound surface area
	15272	Application of skin substitute graft to trunk, arms, legs, total wound surface area up to 100 sq cm; each additional 25 sq cm wound surface area, or part thereof (List separately in addition to code for primary procedure)
	15273	Application of skin substitute graft to trunk, arms, legs, total wound surface area greater than or equal to 100 sq cm; first 100 sq cm wound surface area, or 1% of body area of infants and children
	15274	Application of skin substitute graft to trunk, arms, legs, total wound surface area greater than or equal to 100 sq cm; each additional 100 sq cm wound surface area, or apart thereof, 1% of body area of infants and children or part thereof (List separately in addition to code for primary procedure)
	15275	Application of skin substitute graft to face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits, total wound surface area up to 100 sq cm; first 25 sq cm or less wound surface area
	15276	Application of skin substitute graft to face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits, total wound surface area up to 100 sq cm; each additional 25 sq cm wound surface area, or part thereof (List separately in addition to code for primary procedure)
	15277	Application of skin substitute graft to face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits, total wound surface area greater than or equal to 100 sq cm; first 100 sq cm wound surface area, or 1% of body area of infants and children
	15278	Application of skin substitute graft to face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits, total wound surface area greater than or equal to 100 sq cm; each additional 100 sq cm surface area, or part thereof, or each additional 1% of body area (List separately in addition to code for primary procedure)

HCPCS		Application Codes
	C1832	Autograft suspension, including cell processing and application, and all system components (ReCell)
	G0681	Application of a premarket approval (pma), 510(k), 361 human cells, tissues or cellular and tissue-based products (hct/p) non-sheet form skin substitute for a wound surface area up to 100 sq cm; first 25 sq cm or less of wound surface area
	G0682	Application of a premarket approval (pma), 510(k), 361 human cells, tissues or cellular and tissue-based products (hct/p) non-sheet form skin substitute for a wound surface area up to 100 sq cm; each additional 25 sq cm wound surface area, or part thereof (list separately in addition to code for primary procedure)
	G0683	Application of a premarket approval (pma), 510(k), 361 human cells, tissues or cellular and tissue-based products (hct/p) non-sheet form skin substitute graft for a wound surface greater than or equal to 100 sq cm; first 100 sq cm wound surface area, or 1% of body area of infants and children
	G0684	Application of a premarket approval (pma), 510(k), 361 human cells, tissues or cellular and tissue-based products (hct/p) non-sheet form skin substitute graft for a wound surface greater than or equal to 100 sq cm; each additional 100 sq cm wound surface area or part thereof, or each additional 1% of body area of infants and children, or part thereof (list separately in addition to code for primary procedure)
		Products
	A2002	Mirragen advanced wound matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	A2003	Bio-connekt wound matrix, per square centimeter
	A2004	Xcellstem, per square centimeter
	A2005	Microlyte matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	A2006	Novosorb synpath dermal matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	A2007	Restrata, per square centimeter (add-on, list separately in addition to primary procedure)
	A2008	Theragenesis, per square centimeter (add-on, list separately in addition to primary procedure)
	A2009	Symphony, per square centimeter (add-on, list separately in addition to primary procedure)

	A2010	Apis, per square centimeter (add-on, list separately in addition to primary procedure)
	A2011	Supra sdrm, per square centimeter (add-on, list separately in addition to primary procedure)
	A2012	Suprathel, per square centimeter (add-on, list separately in addition to primary procedure)
	A2013	Innovamatrix fs, per square centimeter (add-on, list separately in addition to primary procedure)
	A2014	Omeza collagen matrix or omeza complete matrix, per 100 mg
	A2015	Phoenix wound matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	A2016	Permeaderm b, per square centimeter (add-on, list separately in addition to primary procedure)
	A2017	Permeaderm glove, each
	A2018	Permeaderm c, per square centimeter (add-on, list separately in addition to primary procedure)
	A2019	Kerecis Omega3, margin shield per square centimeter (add-on, list separately in addition to primary procedure)
	A2020	Ac5 Advanced Wound System (Ac5)
	A2021	Neomatrix, per square centimeter (add-on, list separately in addition to primary procedure)
	A2022	Innovaburn or innovamatrix xl, per square centimeter (add-on, list separately in addition to primary procedure)
	A2023	Innovamatrix pd, 1 mg
	A2024	Resolve matrix or xenopatch, per square centimeter (add-on, list separately in addition to primary procedure)
	A2025	Miro3d, per cubic centimeter (add-on, list separately in addition to primary procedure)
	A2026	Restrata minimatrix, 5mg
	A2027	Matriderm, per square centimeter (add-on, list separately in addition to primary procedure)
	A2028	Micromatrix flex, per mg
	A2029	Mirotract wound matrix sheet, per cubic centimeter (add-on, list separately in addition to primary procedure)

	A2030	Micro3d fibers, per milligram
	A2031	Mirodry wound matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	A2032	Myriad matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	A2033	Myriad morcells, 4 milligrams
	A2034	Foundation drs solo, per square centimeter (add-on, list separately in addition to primary procedure)
	A2036	Cohealyx collagen dermal matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	A2037	G4derm plus/suprello, per milliliter
	A2038	Marigen pacto, per square centimeter (add-on, list separately in addition to primary procedure)
	A2039	Innovamatrix fd, per square centimeter (add-on, list separately in addition to primary procedure)
	A2040	Microlyte painguard, per square centimeter
	A2041	Foundation drs + duo, per square centimeter
	A2042	Foundation drs + solo, per square centimeter
	A2043	Biobrane, per square centimeter
	A2044	Biobrane glove, each
	A2045	Novashield or novagen wound matrix, per square centimeter
	A4100	Non-sheet form skin substitute, fda cleared as a device, not otherwise specified (list in addition to primary procedure)
	A6460	Synthetic resorbable wound dressing, sterile, pad size 16 sq. in. or less, without adhesive border, each dressing
	A6461	Synthetic resorbable wound dressing, sterile, pad size more than 16 sq. in. but less than or equal to 48 sq. in., without adhesive border, each dressing
	C9354	Acellular pericardial tissue matrix of nonhuman origin (Veritas), per square centimeter
	C9356	Tendon, porous matrix of cross-linked collagen and glycosaminoglycan matrix (TenoGlide Tendon Protector Sheet), per square centimeter
	C9358	Dermal substitute, native, non-denatured collagen, fetal bovine origin (SurgiMend Collagen Matrix), per 0.5 square centimeters

	C9360	Dermal substitute, native, nondenatured collagen, neonatal bovine origin (SurgiMend Collagen Matrix), per 0.5 square centimeters
	C9363	Skin substitute, Integra Meshed Bilayer Wound Matrix, per square centimeter
	C9364	Porcine implant, Permacol, per square centimeter
	Q4101	Apligraf, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4102	Oasis Wound Matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4103	Oasis Burn Matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4104	Integra Bilayer Matrix Wound Dressing (BMWD), per square centimeter (add-on, list separately in addition to primary procedure)
	Q4105	Integra Dermal Regeneration Template (DRT) or Integra Omnigraft dermal regeneration matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4107	Graftjacket, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4108	Integra Matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4110	PriMatrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4111	GammaGraft, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4112	Cymetra, injectable, 1 cc
	Q4113	Graftjacket Xpress, injectable, 1 cc
	Q4114	Integra Flowable Wound Matrix, injectable, 1 cc
	Q4115	AlloSkin, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4116	AlloDerm, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4117	Hyalomatrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4118	MatriStem micromatrix, 1 mg

	Q4121	TheraSkin, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4122	Dermacell, dermacell awm or dermacell awm porous, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4123	AlloSkin RT, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4124	Oasis Ultra Tri-Layer Wound Matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4125	Arthroflex, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4126	Memoderm, Deraspan, Transgraft or Integuply, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4127	Talymed, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4128	Flex hd, or allopatch hd, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4130	Strattice TM, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4134	hMatrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4135	Mediskin, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4136	EZ-derm, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4141	Alloskin AC, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4142	Xcm biologic tissue matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4143	Repriza, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4146	TenSIX, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4147	Architect, Architect PX, or Architect FX, extracellular matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4149	Excellagen, 0.1 cc

	Q4152	DermaPure per square centimeter (add-on, list separately in addition to primary procedure)
	Q4158	Kerecis Omega3, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4161	Bio-ConneKt Wound Matrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4164	Helicoll, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4165	Keramatrix, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4166	Cytal, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4167	Truskin, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4175	Miroderm, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4179	Flowerderm, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4182	Transcyte, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4193	Coll-e-derm, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4195	Puraply, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4196	Puraply am, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4197	Puraply xt, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4200	Skin te, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4202	Keroxx (2.5g/cc), 1cc
	Q4203	Derma-gide, per square centimeter (add-on, list separately in addition to primary procedure)
	Q4222	Progenamatrix, per square centimeter (add-on, list separately in addition to primary procedure)

	Q4226	MyOwn skin, includes harvesting and preparation procedures, per square centimeter
	Q4431	Pma skin substitute product, not otherwise specified (list in addition to primary procedure)
	Q4432	510(k) skin substitute, not otherwise specified (list in addition to primary procedure)
	Q4433	361 hct/p skin substitute product, not otherwise specified (list in addition to primary procedure)
Type of Service	Medicine	
Place of Service	Inpatient/Outpatient	

POLICY HISTORY

Date	Action	Action
July 2026	Interim Review	Policy Revised
May 2026	Annual Review	Policy Revised
April 2025	Annual Review	Policy Revised
April 2024	Annual Review	Policy Revised
December 2023	Interim Review	Policy Revised
April 2023	Annual Review	Policy Revised
March 2022	Annual Review	Policy Revised
March 2021	Annual Review	Policy Revised
March 2020	Annual Review	Policy Revised
March 2019	Annual Review	Policy Revised
March 2018	Annual Review	Policy Revised
March 2017	Annual Review	Policy Revised
November 2016	Interim Review	Policy Revised
September 2016	Interim Review	Policy Revised

Date	Action	Action
March 2016	Annual Review	Policy Revised
April 2015	Annual Review	Policy Revised
June 2014	Annual Review	Policy Revised
July 2013	Annual Review	Policy Revised
May 2013	Interim Review	Policy Renewed
August 2012	Annual Review	Policy Renewed
September 2011	Annual Review	Policy Renewed

New information or technology that would be relevant for Wellmark to consider when this policy is next reviewed may be submitted to:

Wellmark Blue Cross and Blue Shield
Medical Policy Analyst
PO Box 9232
Des Moines, IA 50306-9232

*CPT® is a registered trademark of the American Medical Association.