

06.01.34 Tumor Treatment Fields Therapy

Original Effective Date: November 2014

Review Date: September 2025

Revised: September 2025

DISCLAIMER/INSTRUCTIONS FOR USE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations, or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

This Medical Policy document describes the status of medical technology at the time the document was developed. Since that time, new technology may have emerged, or new medical literature may have been published. This Medical Policy will be reviewed regularly and updated as scientific and medical literature becomes available; therefore, policies are subject to change without notice.

Related Policies:

- [06.01.15 Stereotactic Radiosurgery \(SRS\) and Stereotactic Body Radiation Therapy \(SBRT\)](#)
[10.01.14 Humanitarian Use Devices](#)

Summary

Description

Tumor treating fields (TTF) therapy is a noninvasive technology intended to treat glioblastoma, malignant pleural mesothelioma and non-small cell lung cancer on an outpatient basis and at home using electrical fields. Glioblastoma multiforme (GBM) is the most common and deadly malignant brain tumor. It has a very poor prognosis and is associated with low quality of life during treatment. Malignant pleural mesothelioma is an aggressive tumor with few treatment options that is associated with significant morbidity and mortality. Non-small cell cancer is the most common type of lung cancer (85%) encompassing 3 subtypes (adenocarcinoma, squamous cell, and large cell carcinoma) and prognosis depends on various factors, including the stage of the cancer, the type of treatment received, and the patient's overall health.

Summary of Evidence

Glioblastoma

For individuals who have newly diagnosed glioblastoma multiforme (GBM) on maintenance therapy after initial treatment who receive tumor treating fields (TTF) therapy as an adjunct to standard maintenance therapy, the evidence includes a randomized controlled trial (RCT) and a systematic review. Relevant outcomes include overall survival (OS), disease-specific survival, symptoms, functional outcomes, quality of life, and treatment-related morbidity. The EF-14 trial found a significant increase of 2.7 months in progression-free survival (PFS) and an increase of 4.9 months in OS with the addition of TTF therapy to standard maintenance therapy (i.e., temozolomide) in patients with newly diagnosed GBM. Although patients were not blinded to treatment assignment, PFS was assessed by blinded evaluators, and the placebo effects on the objective measure of OS are expected to be minimal. In a systematic review that included the EF-14 trial along with other observational studies, the pooled median OS and PFS in newly diagnosed patients who received TTF therapy was 21.7 months and 7.2 months, respectively. This technology represents a clinically significant option in the treatment of patients with GBM, for whom options are limited. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have progressive or recurrent glioblastoma multiforme (GBM) who receive tumor treating fields (TTF) therapy as an adjunct or alternative to standard medical therapy, the evidence includes a randomized controlled trial (RCT), nonrandomized comparative studies, and a systematic review of these data. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related morbidity. The single RCT evaluating TTF therapy for recurrent GBM did not show superiority of TTF therapy for the primary outcome (OS) compared with physicians' choice chemotherapy. Because no serious adverse effects have been identified with TTF therapy, this raises the possibility that treatment with TTF might reduce the toxicity associated with treatment for recurrent GBM. A reduction in chemotherapy-associated toxicity without loss of efficacy would be considered a net health benefit. However, this RCT is not sufficient to permit conclusions on the efficacy of the device. Because the trial was not designed as a noninferiority trial, no inferences of noninferiority compared with chemotherapy can be made. Also, quality of life assessment was measured in an insufficient number of patients to reach firm conclusions on differences in quality of life between TTF therapy and medical treatment. The highest quality study of TTF combined with medical treatment for recurrent GBM is a post hoc analysis of the EF-14 trial. Two registry studies also evaluated real-world outcomes in patients enrolled in the PRiDe registry compared to patients in the EF-11 study. In a systematic review that included the RCT, and post hoc analysis of the EF-14 trial, along with other observational studies, the pooled median OS and PFS in individuals with recurrent GBM who received TTF therapy was 10.3 months and 5.7 months, respectively. A high-quality, prospective RCT is needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome. However, even though there is a paucity of data in the peer-reviewed scientific literature comparing TTF to other treatment options for recurrent glioblastoma multiforme (GBM), the therapy is supported by relevant professional societies (see [Practice Guidelines and Position Statements](#) below) TTF will be considered medically necessary for the treatment of recurrent glioblastoma (GBM) when the [criteria](#) below has been met.

Malignant Pleural Mesothelioma (MPM)

For individuals who have unresectable, locally advanced or metastatic, malignant pleural mesothelioma (MPM) who receive TTF therapy as an adjunct to standard maintenance therapy, the evidence includes a single-arm prospective study conducted in 80 patients and a retrospective study of 5 US patients. Relevant outcomes include OS, disease-specific survival, symptoms, functional outcomes, quality of life, and treatment-related morbidity. In patients who received TTF therapy in combination with pemetrexed and cisplatin or carboplatin, median OS was 18.2 months (95% confidence interval [CI], 12.1 to 25.8

months). Because there was no comparison group, it is not possible to make conclusions about the effectiveness of the intervention compared to medical therapy alone. The retrospective study is the first publication of real-world implementation of TTF for MPM. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Non-Small Cell Lung Cancer

For individuals who have metastatic non-small cell lung cancer (NSCLC) who receive TTF with concurrent standard care including an immune checkpoint inhibitor or docetaxel and who have progressed on or after platinum-based therapy, the evidence includes an open-label RCT conducted in 276 patients. Relevant outcomes include OS, disease-specific survival, symptoms, functional outcomes, quality of life, and treatment-related morbidity. The LUNAR trial found a significant increase of 3.3 months in OS with TTF in combination with an immune checkpoint inhibitor or docetaxel, but there was no significant improvement in PFS or overall response rate. The trial is limited by the lack of a sham comparator, a lack of baseline molecular testing, and changes in standard of care. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Additional Information

None

OBJECTIVE

The objective of this evidence review is to determine whether the use of tumor treating fields therapy improves the net health outcome for individuals with solid tumors including glioblastoma multiforme, malignant pleural mesothelioma and non-small cell lung cancer (NSCLC)

PRIOR APPROVAL

Prior approval is required.

POLICY

Diagnosed Glioblastoma (GBM)

Tumor treatment fields (TTF) therapy to treat histologically confirmed **glioblastoma multiforme (GBM)** as an adjunctive treatment to standard maintenance therapy with Temodar® (temozolomide) is considered **medically necessary** following initial treatment with **surgery, radiotherapy and/or chemotherapy** when **one of the** following are met (Choose A or B):

- A. In newly diagnosed GBM when **all the following** criteria are met
 - Individuals ≥ 18 years of age;
 - Supratentorial tumor (see [Policy Guidelines](#));
 - Karnofsky Performance Status (KPS) score ≥ 60%;
 - The individual understands device use and will comply with the criteria according to the Food and Drug Administration (FDA) label (see [Policy Guidelines](#)).

- B. In **recurrent GBM** when all of criteria have been previously met for a newly diagnosed GBM.

Continuation for Diagnosed Glioblastoma (GBM)

Tumor treatment field (TTF) therapy (e.g., Optune®) for continued use to treat a diagnosed glioblastoma (GBM) is considered **medically necessary** when **all of the** following conditions are met:

- The initial review for the applicable clinical scenario for a newly diagnosed glioblastoma (GBM) **or** recurrent glioblastoma (GBM) medical review criteria above have been met; **and**
- No evidence of disease progression within 90 days of the continuation request (see [Policy Guidelines](#)); **and**
- The individual continues to understand device use, including the requirement of a shaved head and is still willing to comply with the use criteria according to the Food and Drug Administration (FDA) label (see [Policy Guidelines](#)).

Mesothelioma

- Tumor Treating Field (TTF) may be considered medically necessary for the treatment of mesothelioma in accordance with the humanitarian device exemption [HDE] specifications of the U.S. Food and Drug Administration [FDA] See policy [10.01.14 Humanitarian Use Devices](#).

Tumor Treatment Field (TTF) Therapy: Investigational

Tumor treatment field (TTF) therapy is considered **investigational in all other conditions** including, but not limited to the following, because the evidence is insufficient in determining the technology results in improved net health outcomes:

- Not meeting the above criteria for **initial** and **recurrent** glioblastoma (GBM); **or**
- As an adjunct to standard medical therapy (e.g., bevacizumab, chemotherapy) for individuals with **progressive glioblastoma** (GBM); **or**
- As an adjunct to **maintenance therapy** other than temozolomide (Temodar®); **or**
- As an alternative to standard medical therapy for individuals with **initial, progressive or recurrent** glioblastoma (GBM) **or**
- For brain metastases; **or**
- For cancer in areas other than the brain; **or**
- As an adjunct to standard medical therapy (pemetrexed and platinum-based chemotherapy) for individuals with malignant pleural mesothelioma
- As an adjunct to standard medical therapy for individuals with non-small cell lung cancer (NSCLC).

ALL Category III codes will be considered investigational unless the code is explicitly addressed as a covered service in a Wellmark BlueCross BlueShield Medical Coverage Policy.

POLICY GUIDELINES

Age

Per the pivotal trial, individuals ≥ 18 years of age were eligible for enrollment. The median age was about 56 years with a range of 19 to 83 years; subgroup analyses for younger age groups were not provided.

U.S. Food and Drug Administration Label

The U.S. Food and Drug Administration label includes the following notices:

- Individuals will maintain a shaved head
- Individuals should use Optune® for at least 18 hours daily to get the best response to treatment.
- Individuals should finish at least 4 full weeks of therapy to get the best response to treatment. Stopping treatment before 4 weeks lowers the chances of a response to treatment.
- Tumor treating fields (TTF) therapy has been used 24 months or less.

Miscellaneous Information

- **Karnofsky Performance Status:** The Karnofsky Performance Scale Index allows individuals to be classified as to their functional impairment. This can be used to compare effectiveness of different therapies and to assess the prognosis in individual patients. The lower the Karnofsky score, the worse the survival for most serious illnesses.
 - The recommended Karnofsky Performance Status (KPS) varies from the NCCN guideline (score ≥ 60). In the pivotal trial the median KPS score at baseline was 90.0, with a range from 60 to 100. Subgroup analyses for individuals with score 60 to 70 were not provided.

Karnofsky Performance Status Scale Definitions Rating (%) Criteria

Able to carry on normal activity and to work; no special care needed.	100	Normal no complaints; no evidence of disease.
	90	Able to carry on normal activity; minor signs or symptoms of disease.
	80	Normal activity with effort; some signs or symptoms of disease.
Unable to work; able to live at home and care for most personal needs; varying amount of assistance needed.	70	Cares for self; unable to carry on normal activity or to do active work.
	60	Requires occasional assistance but is able to care for most of his personal needs.
	50	Requires considerable assistance and frequent medical care.
Unable to care for self; requires equivalent of institutional or hospital care; disease may be progressing rapidly.	40	Disabled; requires special care and assistance.
	30	Severely disabled; hospital admission is indicated although death not imminent.
	20	Very sick; hospital admission necessary; active supportive treatment necessary.
	10	Moribund; fatal processes progressing rapidly.
	0	Dead

- **Progressive Disease:** Progression was defined in the EF-14 trial (Stupp et al [2015, 2017]) according to the MacDonald criteria (tumor growth $> 25\%$ compared with the smallest tumor area measured in the individual during the trial or appearance of 1 or more new tumors in the brain that are diagnosed radiologically as glioblastoma multiforme).
- **Recurrent Disease:** Cancer that has recurred (come back), usually after a period of time during which the cancer could not be detected. The cancer may come back to the same place as the original (primary) tumor or to another place in the body.
- **The Supratentorial Region:** The supratentorial region of the brain is the area located above the tentorium cerebelli and it contains the cerebrum. The area of the brain below the tentorium cerebelli is the infratentorial region and it contains the cerebellum. Tumor treatment field therapy is only recommended for those with supratentorial disease.

Coding

See the [Codes table](#) for details.

BACKGROUND

Glioblastoma Multiforme

Glioblastomas, also known as glioblastoma multiforme (GBM), are the most common form of malignant primary brain tumor in adults. Glioblastomas are grade IV astrocytomas, a rapidly progressing and deadly type of glial cell tumor that is often resistant to standard medical therapy (e.g., bevacizumab, chemotherapy). Together, anaplastic astrocytomas and glioblastomas comprise approximately 38% of all primary malignant brain tumors. Mean age at GBM diagnosis is 65 years. Glioblastomas have the lowest survival rate of any central nervous system tumor; the 5-year survival rate and average length of survival is estimated at 6.9% and 8 months, respectively. Currently there is no known cure, but treatments are available. Tumor treatment fields (TTF) (e.g., Optune® device) is one such treatment that aims to prolong progression-free survival (PFS) and overall survival (OS).

Note: Due to its multiple forms, it is also termed Glioblastoma multiforme (GBM). The term "multiforme" is no longer part of the World Health Organization (WHO) designation, though glioblastoma is still often abbreviated "GBM."

The types of glioblastomas collectively referenced as glioblastoma are differentiated below:

- Glioblastoma, IDH-wildtype
 - Giant cell glioblastoma
 - Gliosarcoma
 - Epithelioid glioblastoma
- Glioblastoma, IDH-mutant
- Glioblastoma, NOS (not otherwise specified)

Treatment of Newly Diagnosed Glioblastoma Multiforme

The primary treatment for individuals newly diagnosed with GBM is to resect the tumor to confirm a diagnosis while debulking the tumor to relieve symptoms of increased intracranial pressure or compression. If total resection is not feasible, subtotal resection and open biopsy are options. During surgery, some patients may undergo implantation of the tumor cavity with a carmustine (bis-chloroethylnitrosourea) impregnated wafer. Due to the poor efficacy of local treatment, postsurgical treatment with adjuvant radiotherapy (RT), chemotherapy (typically temozolomide), or a combination of these 2 therapies is recommended. After adjuvant therapy, patients may undergo maintenance therapy with temozolomide. Maintenance temozolomide is given for 5 days of every 28-day cycle for 6 cycles. Response and overall survival rates with temozolomide are higher in patients who have O⁶-methylguanine-DNA methyltransferase (*MGMT*) gene promoter methylation.

Prognostic factors for therapy success are age, histology, performance status or physical condition of the patient, and extent of resection. National Comprehensive Cancer Network recommendations include patient age and Karnofsky Performance Status score as important determinants of postsurgical treatment choice (see the Supplemental Information section). For patients with good performance status, the most aggressive treatment (standard RT plus temozolomide) is recommended. For patients with poor performance status, only single treatment cycles or even palliative or supportive care are recommended. Hypofractionated RT is indicated for patients with poor performance status because it is better tolerated, and more patients are able to complete RT.

Treatment of GBM is rarely curative, and tumors will recur in essentially all individuals.

Treatment of Recurrent Glioblastoma Multiforme

When disease recurs, additional debulking surgery may be used if the recurrence is localized. Due to radiation tolerances, re-radiation options for patients with recurrent GBM who have previously received initial external-beam RT are limited. There is no standard adjunctive treatment for recurrent GBM. Treatment options for recurrent disease include various forms of systemic medications such as the antivascular endothelial growth factor drug bevacizumab, alkylating agents such as nitrosoureas (e.g., lomustine, carmustine), or retreatment with temozolomide. Medical therapy is associated with side effects that include hematologic toxicity, headache, loss of appetite, nausea, vomiting, and fatigue. Response rates in recurrent disease are less than 10%, and the progression-free survival rate at 6 months is less than 20%. There is a need for new treatments that can improve survival in patients with recurrent GBM or reduce the side effects of treatment while retaining survival benefits.

Malignant Pleural Mesothelioma

Malignant pleural mesothelioma (MPM) is an aggressive tumor that is associated with significant morbidity and mortality. It is associated with asbestos exposure and has a latency period of about 40 years after asbestos exposure. Recommendations for treatment are mainly chemotherapy as first line with pemetrexed plus platinum. Surgical cytoreduction is also recommended in selected individuals with early-stage disease. Adjuvant radiation can be offered for individuals who have resection of intervention tracts found to be histologically positive or for palliation of symptomatic individuals.

Non-small Cell Lung Cancer

Lung cancer, including non-small cell lung cancer (NSCLC), is the leading cause of cancer-related death in the United States. There are numerous treatment options for NSCLC which have improved survival rates. Patients eligible for targeted or immunotherapies now have 5-year survival rates up to 62.5%. Tumor treating fields have been studied in combination with immune checkpoint inhibitors (i.e., PD-1/PD-L1 inhibitors) or docetaxel in patients with metastatic NSCLC who progressed with platinum-based therapy

Regulatory Status

Date	Description
April 2011	The NovoTTF-100A™ System (Novocure; assigned the generic name of TTF) was approved by the U.S. Food and Drug Administration (FDA) through the premarket approval process. The FDA approved label reads as follows: "The NovoTTF-100A System is intended as a treatment for adult patients (22 years of age or older) with confirmed GBM, following confirmed recurrence in an upper region of the brain (supratentorial) after receiving chemotherapy. The device is intended to be used as a stand-alone treatment and is intended as an alternative to standard medical therapy for recurrent GBM after surgical and radiation options have been exhausted."
September 2014	The FDA approved Novocure's request for a product name change from NovoTTF-110A System to Optune®.
October 2015	The FDA expanded the indication for Optune in combination with temozolomide to include newly diagnosed GBM. ⁷ The device was granted priority review status in May 2015 because there was no legally marketed alternative device available for the treatment of newly diagnosed GBM, a life-threatening condition.
July 2016	A smaller, lighter version of the Optune® device, called the Optune® System (NovoTTF-200A System), received FDA approval. The FDA labeled indication are as follows:

	<p>Optune™ is intended as a treatment for adult patients (22 years of age or older) with histologically confirmed glioblastoma (GBM).</p> <p>Optune™ with temozolomide (TMZ) is indicated for the treatment of adult patients with newly diagnosed, supratentorial glioblastoma following maximal debulking surgery and completion of radiation therapy together with concomitant standard of care chemotherapy.</p> <p>For the treatment of recurrent GBM, Optune™ is indicated following histologically- or radiologically confirmed recurrence in the supra-tentorial region of the brain after receiving chemotherapy.</p> <p>The device is intended to be used as a monotherapy and is intended as an alternative to standard medical therapy for GBM after surgical and radiation options have been exhausted.</p>
May 2019	The FDA approved a modified version of the Optune System (NovoTTF-100A System), which is now called the Optune Lua™ System (NovoTTF™-100L System), for “treatment of adult patients with unresectable, locally advanced or metastatic, malignant pleural mesothelioma (MPM) to be used concurrently with pemetrexed and platinum-based chemotherapy.
March 2020	The manufacturer of Optune products announced a plan to include a suffix after the brand name for newly approved indications to further delineate specific indications for individual products (e.g., Tumor Treatment Fields).
September 2021	The FDA granted breakthrough designation to the NovoTTF-200T System for use together with atezolizumab and bevacizumab for the first-line treatment of patients with unresectable or metastatic liver cancer.
2023	Optune was renamed Optune Gio™.
October 2024	The Optune Lua was approved for "metastatic non-small cell lung cancer along with concurrent drug treatments in adults who have progressed on or after a platinum-based chemotherapy."

To date, all the existing tumor treating fields products fall under the brand name Optune®. In March 2020, the manufacturer of Optune®. products announced a plan to include a suffix after the brand name for newly approved indications to further delineate specific indications for individual products (e.g., Optune®. Lua).

The subjects receive multiple one-month courses of continuous Optune treatment. Optune treatment was discontinued in the case of clinical disease progression, if a device-related serious adverse event occurred, or after 24 months or second progression whichever occurred first.

FDA Labeled Contraindications:

For TTF therapy (e.g., Optune®) in individuals with a glioblastoma (GBM) the FDA has listed contraindications to include but not limited to the following:

- bullet fragment(s)
- a skull defect (i.e., missing bone with no replacement)
- an active implanted medical device to include but not limited to **one of the following:**
 - defibrillators
 - deep brain stimulators

- pacemakers
- programmable shunts
- spinal cord stimulators
- vagus nerve stimulators

RATIONALE

This evidence review was created in November 2014 and has been updated regularly with searches of the PubMed database. The most recent literature update was performed through August 2025.

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function, including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Tumor Treating Fields Therapy as an Adjunct to Standard Maintenance Care for Newly Diagnosed Glioblastoma Multiforme

Clinical Context and Therapy Purpose

The purpose of TTF therapy, also referred to as alternating electrical field therapy, is to provide a treatment option that is an alternative to or an improvement on existing therapies for individuals with newly diagnosed GBM. Tumor treating fields therapy has been investigated as an adjunct to temozolomide for the treatment of newly diagnosed GBM and as an alternative or adjunct to medical therapy for progressive or recurrent GBM.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest are individuals who have newly diagnosed GBM and good performance status. Newly diagnosed patients would have undergone initial treatment with surgery, radiation therapy, and chemotherapy and be receiving maintenance chemotherapy.

Interventions

Tumor treating fields therapy is a noninvasive technology intended to treat GBM on an outpatient basis and at home using electrical fields. Tumor treating fields therapy exposes rapidly dividing cancer cells to electric fields of low intensity and intermediate frequency (150 or 200 kHz) that alternate in perpendicular orientation. Tumor treating fields therapy is proposed to inhibit tumor growth by 2 mechanisms: the arrest

of cell proliferation by causing microtubule misalignment in the mitotic spindle of rapidly dividing tumor cells and apoptosis due to movement of macromolecules and organelles during telophase. Preclinical studies have indicated that the electric fields may also make the cells more susceptible to chemotherapy.

Optune branded products (formerly NovoTTF-100A System) are the only legally marketed TTF delivery system available in the United States. The portable, battery-powered device is carried in a backpack or shoulder pack while carrying out activities of daily living. For the treatment of glioblastoma, 4 disposable transducer arrays with insulated electrodes are applied to the patient's shaved head. The transducer array layout is typically determined using specialized software. The patient's scalp is re-shaved and the transducer arrays replaced twice a week by the patient, caregiver, or device technician. The device is worn for up to 24 hours a day for the duration of treatment, except for brief periods for personal hygiene and 2 to 3 days at the end of each month. The minimum daily treatment is 18 hours. The minimum duration of treatment is 1 month, with the continuation of treatment available until recurrence.

Comparators

The following practice is currently being used to make decisions about newly diagnosed GBM: maintenance chemotherapy with temozolomide alone.

Tumor treating fields therapy might also be compared with palliative or supportive care, where survival rarely exceeds 3 to 5 months.

Outcomes

The general outcomes of interest are whether TTF improves survival or quality of life during treatment and the time to tumor recurrence because most GBMs recur. Measures of cognitive status and quality of life measures are also of interest to determine whether TTF alters the decline in cognition and quality of life that occur with GBM. Also, adverse events of treatment, such as side effects of chemotherapy and the possibility of seizures, need to be assessed.

Due to the rapid progression of GBM, the time of interest for both progression-free survival (PFS) and overall survival (OS) is months.

Study Selection Criteria

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

A Health Technology Assessment last reviewed by Hayes Inc. on January 5, 2023, reported a Hayes C rating for adults who are 22 years of age and older with a newly diagnosed GBM following surgery and radiation therapy with concomitant chemotherapy and uses TTF. Hayes indicated the C rating, "reflects positive but low-quality evidence from 1 randomized controlled trial and 1 very-poor-quality cohort study suggesting that TTF plus TMZ may prolong survival (OS and PFS) with similar QOL relative

to TMZ alone, with no serious safety issues. The Rating also reflects the small body of evidence available for this indication.”

Regev et al (2021) conducted a systematic review of studies describing the use of TTF therapy for the treatment of GBM. The authors included a total of 20 studies of patients with newly diagnosed GBM and recurrent GBM. For newly diagnosed GBM (n=542), only 1 RCT was identified (Stupp et al, 2017), which is described in further detail in the section below. The remainder of the data for newly diagnosed GBM was observational. The pooled median OS and PFS in newly diagnosed patients was 21.7 months (95% confidence interval [CI], 19.6 to 23.8) and 7.2 months (95% CI, 6.1 to 8.2) months, respectively. The pooled rate of OS at 1, 2, and 3 years was 73.5%, 45.1%, and 29.3%, respectively. The pooled rate of PFS at 6, 12, and 18 months was 55.9%, 32.4%, and 21.7%, respectively. Statistical comparisons to other treatment modalities were not provided.

Randomized Controlled Trials

Stupp et al (2017) published results of the EF-14 multicenter, open-label phase 3 RCT that evaluated maintenance therapy with TTF for newly diagnosed GBM. The trial included 695 patients from 83 sites who had supratentorial GBM and had completed standard treatment consisting of biopsy or surgical resection followed by RT and chemotherapy (see the table below). A Karnofsky Performance Status (KPS) score of 70 or higher was an additional inclusion criterion to ensure independence in activities of daily living, and patients with rapidly progressing GBM following radiochemotherapy were excluded from the trial. Patients were randomized in a 2:1 fashion to TTF plus maintenance temozolomide or maintenance temozolomide alone.

All individuals were seen monthly for follow-up. Quality of life was assessed every 3 months, and magnetic resonance imaging (MRI) was performed every 2 months until tumor progression. Tumor progression on MRI was adjudicated by a central review committee blinded to treatment group. The primary outcome was PFS, and the secondary outcome was OS. The analysis was by intention-to-treat, including 26 patients from the control arm who crossed over to TTF following the planned interim analysis.

In 2014, an independent data and safety monitoring board concluded from the planned interim analysis that the trial met its predefined boundaries for success (improvement in PFS and OS) and recommended trial termination. The U.S. Food and Drug Administration (FDA) approved the trial termination, and the trial was closed to recruitment with 695 of the planned 700 participants randomized. Control arm participants were allowed to cross over to the experimental treatment at this time. The interim analysis, which the U.S. FDA considered for the 2015 expanded approval of Optune, was published by Stupp et al (2015). At the time of the interim analysis, data were available for 210 patients randomized to TTF plus temozolomide and 105 patients to temozolomide alone. Follow-up of the remainder of the 695 enrolled patients continued after enrollment was closed.

Table 1: Key Randomized Controlled Trial Characteristics for Newly Diagnosed Glioblastoma

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
Stupp et al (2017); EF-14	U.S., E.U., South Korea, Israel	83	2009-2016	<ul style="list-style-type: none"> • 695 newly diagnosed with GBM and treated by radiochemotherapy • KPS score ≥ 70 	TTF >18 h/d plus maintenance temozolomide (n=466)	Maintenance temozolomide alone (5 d every 28 d for 6 cycles) (n=229)

E.U.: European Union; GBM: glioblastoma multiforme; h/d: hours per day; KPS: Karnofsky Performance Status; TTF: tumor treatment fields.

Results of the final analysis of the EF-14 trial were similar to the interim analysis and are shown in Table 2. Both PFS and OS improved with the addition of TTF therapy to standard maintenance chemotherapy (i.e., temozolomide). PFS increased by 2.7 months ($p < .001$) and OS increased by 4.9 months ($p < .001$) in the TTF group. The time to a decrease in mental function was 2.5 months longer with TTF therapy ($p < .01$).

There was a similar percentage of dropouts at the final analysis with 49 (11%) patients in the TTF group and 27 (12%) patients in the temozolomide alone group. More treatment cycles with temozolomide were administered in the TTF group (median, 6 for TTF group vs 5 for controls), a finding that is consistent with the longer PFS. Rates of adverse events were similar between the groups, including rates of seizures. In a secondary analysis of patients who had not progressed, there was no reduction in health-related quality of life with TTF compared with temozolomide alone aside from "itchy skin".¹⁴ Interpretation of this result is limited by the low percentage of patients who completed the health-related quality of life assessments at follow-up (65.8% of the 655 patients alive at 3 months and 41.7% of the 473 patients alive at 12 months). A mixed-model analysis, which accounts for missing data, confirmed the results of the mean change from baseline analysis.

Table 2: Key Randomized Controlled Trial Results for Newly Diagnosed Glioblastoma

Study	Final N (%)	Median PFS (95% CI), months	Median OS (95% CI), months	Systemic Adverse Events, n (%)	Seizures, n (%)	Time to 6-Point Decline in MMSE Score (95% CI), months
Stupp et al (2017)						
TTF + temozolomide	417 (89)	6.7 (6.1 to 8.1)	20.9 (19.3 to 22.7)	218 (48)	26 (6)	16.7 (14.7 to 19.0)
Temozolomide alone	202 (88)	4.0 (3.8 to 4.4)	16.0 (14.0 to 18.4)	94 (44)	13 (6)	14.2 (12.7 to 17.0)
HR (95% CI)		0.63 (0.52 to 0.76)	0.63 (0.53 to 0.76)			0.79 (0.66 to 0.95)
p-value		<.001	<.001	.58		.01

CI: confidence interval; HR: hazard ratio; MMSE: Mini-Mental State Examination; OS: overall survival; PFS: progression-free survival; TTF: tumor treatment fields.

The tables below display notable limitations identified in this trial; a major limitation is the lack of patient blinding to treatment assignment. However, PFS was assessed by investigators who were blinded to treatment, and placebo effects on OS measurement were expected to be minimal. Investigators considered it practically unfeasible (due to the heat and current of the TTF therapy) and ethically unacceptable to submit the control patients to repeated shaving of the head and continuous wear of a sham device over many months.

Table 3: Study Relevance Limitations

Study; Trial	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
Stupp et al (2017); EF-14			3. Possible differences in post-progression treatment affecting OS		

OS: overall survival. The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. ^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other. ^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other. ^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other. ^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other. ^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 4: Study Design and Conduct Limitations

Study; Trial	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Stupp et al (2017); EF-14		1. No sham control and not blinded to treatment assignment				

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. ^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other. ^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other. ^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other. ^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other. ^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other. ^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

Section Summary: Tumor Treating Fields Therapy as an Adjunct to Standard Maintenance Care for Newly Diagnosed Glioblastoma Multiforme

The final analysis of the EF-14 trial, which included 695 patients from 83 sites, found a statistically and clinically significant increase of 2.7 months in PFS and an increase of 4.9 months in OS with the addition of TTF therapy to standard maintenance therapy (i.e., temozolomide) in patients with newly diagnosed GBM. There was no sham control, and patients were not blinded to treatment assignment. However, PFS

was assessed by blinded evaluators, and placebo effects on the objective measure of OS were likely to be minimal. There was no evidence of a negative impact of TTF therapy on health-related quality of life, except for itchy skin from the transducers. In a systematic review that included the EF-14 trial along with other observational studies, the pooled median OS and PFS in newly diagnosed patients who received TTF therapy was 21.7 months and 7.2 months, respectively.

Tumor Treating Fields Therapy as an Adjunct or Alternative to Medical Therapy for Progressive or Recurrent Glioblastoma Multiforme

Clinical Context and Therapy Purpose

The purpose of TTF therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies for individuals with progressive or recurrent GBM.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals who have recurrent GBM with good performance status.

Interventions

The therapy being considered is TTF therapy as an adjunct or alternative to standard medical therapy.

Comparators

The following practice is currently being used to make decisions about progressive or recurrent GBM: standard medical therapy (e.g., bevacizumab, nitrosoureas, temozolomide rechallenge).

Outcomes

The general outcomes of interest are whether TTF improves survival or quality of life during treatment and the time to tumor recurrence because most GBMs recur. Measures of cognitive status and quality of life measures are also of interest to determine whether TTF alters the decline in cognition and quality of life that occur with GBM. Also, adverse events of treatment, such as side effects of chemotherapy and the possibility of seizures, need to be assessed.

Due to the rapid progression of GBM, the time of interest for both PFS and OS is months.

Study Selection Criteria

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

A Health Technology Assessment last reviewed by Hayes Inc. on January 5, 2023, reported a Hayes C rating for adults who are 22 years of age and older with a recurrent GBM following surgery and radiation therapy who uses TTF as monotherapy. Hayes indicated the C rating, “reflects some positive but low-quality evidence suggesting that TTF therapy is at least as effective as chemotherapy for prolonging survival (overall survival [OS], progression-free survival [PFS]) and improving quality of life (QOL), with no serious safety issues. The Rating also reflects the small body of evidence available, small sample sizes, and lack of concurrent control or comparator groups in most studies.”

A systematic review by Regev et al (2021) is introduced above. For patients with recurrent GBM (n=1094), only 2 RCTs were identified (Stupp et al [2012] and post hoc analysis of Kesari et al [2017]), which are described in further detail in the section below. The remainder of the data for recurrent GBM was observational. For patients with recurrent GBM, the pooled median OS and PFS were 10.3 months (95% CI, 8.3 to 12.8) and 5.7 (95% CI, 2.8 to 10) months, respectively. The pooled rate of OS at 1, 2, and 3 years was 43.7%, 21.3%, and 14%, respectively. The pooled rate of PFS at 6, 12, and 18 months was 47.8%, 29.3%, and 19.7%, respectively. As previously noted, statistical comparisons to other treatment modalities were not provided.

Randomized Controlled Trials

The 2011 U.S. FDA approval of the NovoTTF-100A System (now called Optune) was based on a phase 3 multinational RCT (EF-11), results of which were published by Stupp et al (2012). This trial compared TTF therapy alone with physician's choice medical therapy in 237 adults who had relapsed or progressive glioblastoma (see the Table below). Patients had failed conventional treatment with RT, chemotherapy, and/or surgery, and more than 80% of participants had failed 2 or more prior chemotherapy regimens. In this trial, the term chemotherapy also applied to targeted agents such as bevacizumab. Patient characteristics and performance of additional post-recurrence debulking surgery were similar in the 2 groups.

Table 5: Summary of Key Randomized Controlled Trial Characteristics for Progressive or Recurrent Glioblastoma

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Stupp et al (2012); EF-11	U.S., E.U., Israel	28	1987-2013	<ul style="list-style-type: none"> • 237 adults with relapsed or progressive supratentorial glioblastoma • KPS score ≥70% 	120 patients treated with TTF alone, 93 (78%) completed 1 cycle	117 patients treated with physician's choice of medical therapy

E.U.: European Union; KPS: Karnofsky Performance Status; TTF: tumor treating fields. Medical therapy included bevacizumab, irinotecan, nitrosoureas, platinum-based chemotherapy (i.e., carboplatin); temozolomide; or a combination of procarbazine, chloroethyl ether, and vincristine.

Participants were followed monthly, which included laboratory tests. Magnetic resonance images were evaluated at 2, 4, and 6 months from initiation of treatment, with subsequent MRIs performed according to local practice until disease progression. Quality of life questionnaires were completed every 3 months. Medical follow-up continued for 2 months after disease progression. Monthly telephone interviews with participants' caregivers were used to assess mortality rates. The primary end point was OS. Secondary end points included PFS, the percentage of patients with PFS at 6 months, time to progression, 1-year survival rate, quality of life, and radiologic response. All end points were evaluated using intention-to-treat analysis.

The trial did not reach its primary end point of improved survival compared with active medical therapy (see the Table below). With a median follow-up of 39 months, 93% of patients had died. There was not a statistically significant difference in survival rates at 1, 2, and 3 years between groups. Patients in the TTF group did not, however, suffer the typical systemic side effects of chemotherapy. The most common adverse event in the TTF group was grade 1 and 2 contact dermatitis on the scalp, which resolved with topical corticosteroids and did not require treatment breaks. Control participants experienced grade 2, 3, or 4 events by organ system related to the pharmacologic activity of chemotherapy agents used. Hematologic events of grade 2 or greater were observed in 17% of chemotherapy patients compared with 3% of TTF patients. Gastrointestinal disorders of grade 2 or greater were identified in 17% of chemotherapy patients compared with 4% of TTF patients. Severe (grades 3 to 4) hematologic and gastrointestinal toxicity was observed in 7% of chemotherapy controls compared with 1% of the TTF group.

Longitudinal quality of life data, available in 63 (27%) participants, showed no meaningful differences between groups for the domains of global health and social functioning. However, cognitive and emotional functioning domains favored TTF therapy. Symptom scale analysis was by treatment-associated toxicity; appetite loss, diarrhea, constipation, nausea, and vomiting were directly related to the chemotherapy administration.

The trial had a number of limitations (see the Tables below), which included lack of blinding and high loss to follow-up. Discontinuation of TTF therapy occurred in 22% of patients due to noncompliance or inability to handle the device, usually within the first few days. In the control group, 21 (18%) patients did not return to the treatment site, and details on disease progression and toxicity were not available. Longitudinal quality of life could be analyzed only for 27% of patients who remained on study therapy for 3 months. The trial was designed as a superiority trial and did not provide adequate evidence of noninferiority.

Table 6: Summary of Key Randomized Controlled Trial Results for Recurrent or Progressive Glioblastoma

Study; Trial	LTFU, n (%)	Median OS, mo	PFS		OS (95% CI), %		
			Median, mo	Rate at 6 Months (95% CI), %	1 Year	2 Years	3 Years
Stupp et al (2012); EF-11							
TTF	23 (22)	6.6	2.2	21.4 (13.5 to 29.3)	20	8 (4 to 13)	4 (1 to 8)
PCC	12 (18)	6.0	2.1	15.1 (7.8 to 22.3)	20	5 (3 to 10)	1 (0 to 3)
HR (95% CI)		0.86 (0.66 to 1.12)	0.81 (0.60 to 1.09)				
P value		.27	.16	.13			

CI: confidence interval; HR: hazard ratio; LTFU: loss to follow-up; OS: overall survival; PCC: physician's choice chemotherapy; PFS: progression-free survival; TTF: tumor treating fields.

Table 7: Study Relevance Limitations

Study	Population	Intervention	Comparator	Outcomes	Follow-Up ^e
Stupp et al (2012); EF-11			2. Physician's choice chemotherapy		

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use. ^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest. ^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively. ^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported. Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 8: Study Design and Conduct Limitations

Study; Trial	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Stupp et al (2012); EF-11		1. Not blinded to treatment assignment		1. 78% of TTF group completed only 1 cycle of therapy, 18% of control group lost to follow-up; longitudinal QOL data were available for 27% of patients		1. Not designed as a noninferiority trial

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

QOL: quality of life; TTF: tumor treating fields. ^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias. ^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician. ^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication. ^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials). ^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference. ^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Nonrandomized Comparative Studies

Zhu et al (2022) conducted a prospective, post-marketing registry study (the EF-19 study) to evaluate the safety and efficacy of TTF versus physician's choice standard of care in patients from the EF-11 study with recurrent glioblastoma. The patient population was comprised of patients already enrolled in the PRiDe registry and included a total of 309 patients. Primary and secondary endpoints assessed included OS in the intention-to-treat (ITT) and per-protocol (PP) populations. In the ITT population, median OS in patients treated with TTF was comparable to physician's choice of standard of care (7.4 vs 6.4 months, respectively; log-rank test p=.053). The Cox test HR was 0.66 (95% CI, 0.47 to 0.92; p=.016). In the PP population, median OS in patients treated with TTF was significantly longer than patients treated with standard of care (8.1 vs 6.4 months; log-rank test p=.017). The Cox test HR was 0.60 (95% CI, 0.42 to 0.85; p=.004). Tumor treating fields therapy showed a favorable safety profile as well.

Kesari et al (2017) conducted a post hoc analysis of the EF-14 trial (see Stupp et al [2017] above) to evaluate the efficacy of TTF in patients who had the first recurrence. Some patients in the temozolomide

alone group crossed over to receive TTF plus chemotherapy after the first recurrence, resulting in 144 patients who received TTF fields plus chemotherapy and 60 patients who received chemotherapy alone for recurrent GBM (see the Table below). Patient characteristics and second-line treatments were well-balanced between the groups, with bevacizumab the most common second-line therapy. The median OS in patients treated with systemic therapy alone was 9.2 months (see Table 10). In comparison, the group of patients who received TTF therapy in addition to systemic therapy had a median OS of 11.8 months ($p=.043$).

A registry study published Mrugala et al (2014) assessed OS data from patients who received NovoTTF therapy in a real-world, clinical practice setting (see Table 9). Concurrent treatment was not captured in the registry, and it is possible that some patients received combination therapy. Median OS in the PRiDe clinical practice dataset (9.6 months) was reported as superior to that attained in the EF-11 pivotal trial (6.6 months, $p<.001$) (see the Table below). More patients in the PRiDe registry were treated for first recurrence (33% vs 9%), and more had received bevacizumab as prior therapy (55% vs 19%). The PRiDe investigators reported no novel or unexpected treatment-related adverse events compared with the EF-11 trial.

Table 9: Characteristics of Key Nonrandomized Trial Results

Study	Study Type	Country	Dates	Participants	TTF	Controls	FU
Zhu et al (2022)	Registry	U.S	2016 - 2018	309 patients with recurrent GBM	192 patients treated with TTF already enrolled in the PRiDe registry	117 patients in the SOC cohort from the EF-11 study	12 months
Kesari et al (2017)	EF-14 post hoc analysis	U.S., E.U., South Korea, Israel	2009-2016	204 patients with first recurrence in the EF-14 trial	144 patients treated with TTF plus second-line chemotherapy	60 patients treated with second-line chemotherapy	12.6 months
Mrugala et al (2014)	Registry	U.S. (91 centers)	2011-2013	457 patients with recurrent GBM	Patient Registry Dataset (PRiDe)	EF-11	NR

E.U.: European Union; FU: follow-up; GBM: glioblastoma; NR: not reported; TTF: tumor treating fields.

Table 10: Summary of Key Nonrandomized Trial Results

Study	Median OS, months	Additional OS outcomes	
Zhu et al (2022)	Median OS with TTF (ITT population), months	Median OS with TTF (PP population), months	
TTF monotherapy	7.4	8.1	
Physician's choice SOC	6.4	6.4	
HR (95%, CI)	0.66 (0.47 to 0.92)	0.60 (0.42 to 0.85)	

Study	Median OS, months	Additional OS outcomes	
p-value	.016	.004	
Kesari et al (2017); EF-14	Median OS without bevacizumab, months	Median OS with bevacizumab, months	
TTF plus chemotherapy	11.8	11.8	
Chemotherapy alone	9.2	9.0	
HR (95% CI)	0.70 (0.48 to 1.00)	0.61 (0.37 to 0.99)	
P value	.049	.043	
Mrugala et al (2014)	Median OS with TTF	1-Year OS, %	2-Year OS, %
PRiDe Registry	9.6	44	30
EF-11	6.6	20	9
HR (95% CI)	0.66 (0.05 to 0.86)	NR	NR
P value	<.001	NR	NR

CI: confidence interval; HR: hazard ratio; NR: not reported; OS: overall survival, TTF: tumor treating fields.

Post hoc analyses of the EF-11 pivotal trial have been reported. Wong et al (2014) published a subgroup analysis to determine characteristics of responders and nonresponders in the active treatment and active treatment control. They found that responders had a lower grade of histology and lower daily dexamethasone use than nonresponders. A second post hoc analysis by Kanner et al. (2014) of the EF-11 pivotal trial data was performed to evaluate OS among patients who finished at least 1 complete course of TTF or chemotherapy. The investigators reported that median OS was 7.7 months in the TTF group compared with 5.9 months in the chemotherapy group ($p=.009$). These post hoc analyses are considered to be hypothesis-generating.

Section Summary: Tumor Treating Fields Therapy as an Adjunct or Alternative to Chemotherapy for Progressive or Recurrent Glioblastoma Multiforme

The single RCT for TTF as an alternative to chemotherapy reported that outcomes following TTF therapy were similar to outcomes following standard chemotherapy. However, this RCT is not sufficient to permit conclusions on the efficacy of the device. The noninferiority of TTF compared with chemotherapy might be considered a sufficient health benefit, if TTF reduced treatment toxicity. However, because the trial was not designed as a noninferiority trial no inferences of noninferiority compared with chemotherapy can be made. Physician's choice therapy during the trial was heterogeneous, although analysis indicated that survival was not affected by choice of chemotherapy. More patients in the TTF group than in the control group did not complete the treatment course. The number of patients who contributed quality of life data was approximately one-quarter of total enrollment, and the self-reported quality of life indicators might have been subject to bias due to the lack of blinding. A nonrandomized post hoc evaluation of the EF-14 trial suggests that TTF may improve survival when combined with chemotherapy for recurrent GBM. This analysis should be considered hypothesis-generating, and further study in high-quality RCTs is needed. In a systematic review that included the RCT and post hoc analysis of the EF-14 trial, along with other observational studies, the pooled median OS and PFS in patients with recurrent GBM who received TTF therapy was 10.3 months and 5.7 months, respectively.

Tumor Treating Fields Therapy as an Adjunct or Alternative to Standard Medical Therapy for Unresectable, Locally Advanced, or Metastatic Malignant Pleural Mesothelioma

Clinical Context and Therapy Purpose

The purpose of TTF therapy as an adjunct or alternative to standard medical therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies for individuals with malignant pleural mesothelioma. Tumor treating fields has been investigated as an adjunct to pemetrexed and platinum-based chemotherapy for the treatment of unresectable, locally advanced or metastatic, malignant pleural mesothelioma (MPM).

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with unresectable, locally advanced or metastatic, MPM.

Interventions

The therapy being considered is TTF as an adjunct or alternative to standard medical therapy.

Optune branded products (formerly NovoTTF-100A System) are the only legally marketed TTF delivery system available in the United States. For the treatment of malignant pleural mesothelioma, the Optune Lua system is used in the same way as the Optune system is used for glioblastoma; however, the 4 disposable transducer arrays with insulated electrodes are applied to the patient's shaved chest and back.

Comparators

The following practice is currently being used to make decisions about unresectable, locally advanced or metastatic, MPM: standard medical therapy with pemetrexed and platinum-based chemotherapy.

Outcomes

The general outcomes of interest are whether TTF improves survival or quality of life during treatment.

The time of interest for both PFS and OS is months to years.

Study Selection Criteria

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Tumor treatment fields therapy for patients with metastatic, MPM has been evaluated in 1 prospective, single-arm study (STELLAR). and a much smaller single-arm retrospective study of 5 patients at a single US center.

Prospective Single-Arm Study

The STELLAR study enrolled 80 patients with inoperable, previously untreated MPM. Study characteristics and results are summarized in the tables below. Patients were treated with cisplatin or carboplatin in combination with TTF therapy delivered by the NovoTTF-100L System at 12 sites outside the U.S. The primary outcome was OS as measured from start of study treatment until date of death. Secondary outcomes were PFS based on investigator assessment of computed tomography (CT) scan imaging, radiological response rate, 1- and 2-year survival rates, and safety.

In STELLAR the median OS was 18.2 months and median PFS was 7.6 months. Seventy-two of the 80 patients enrolled had at least 1 follow-up CT scan. Of those, 40% had a partial response, 57% had stable disease, and 3% progressed. The only adverse event associated with TTF treatment was skin reaction; this adverse event was mild to moderate for the majority of patients who experienced it (66%). The limitations of the STELLAR study are summarized in the tables below. Because there was no control group, it is not possible to draw conclusions about the effectiveness of TTF therapy compared to standard medical care alone. Additional limitations include the small sample size and no reporting of symptoms or quality of life outcomes.

Table 11: Summary of The STELLAR Single Arm Study

Study	Study Type	Country	Dates	Participants	Treatment	Follow-Up
STELLAR (2019) NCT02397928	Prospective, single-arm, multicenter (12 sites)	E.U.	2015-2017	Age 18 years or older, with mesothelioma, not candidate for curative treatment (surgery or RT), ≥1 evaluable lesion, ECOG Performance Status of 0 to 1, at least 4 weeks since last surgery, life expectancy at least 3 months, and able to operate the device independently or with help of a caregiver	TTF (delivered by the NovoTTF-100L System) for ≥18 hours per day in combination with pemetrexed and cisplatin or carboplatin N=80	Protocol specified minimum follow-up of at least 12 months

ECOG: Eastern Cooperative Oncology Group; E.U.: European Union; RT: radiotherapy; TTF: tumor treating fields

Table 12: Summary of The STELLAR Single Arm Study Results

Study	Median OS (95% CI), months	Median PFS (95% CI), months	One-year Survival (95% CI)	2-year survival (95% CI)	Response
STELLAR (2019) NCT02397928	18.2 (12.1 to 25.8)	7.6 (6.7 to 8.6)	62.2% (50.3% to 72.0%)	41.9% (28.0% to 55.2%)	Of 72 who had a follow-up CT scan: 29/70 (40%) partial response 41 /70 (57%) stable disease 2/70 (3%) progressed

CI: confidence interval; CT: computed tomography; OS: overall survival; PFS: progression free survival

Table 13: Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
STELLAR (2019) NCT02397928			2. No comparator	1. Quality of life not assessed	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other. ^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other. ^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other. ^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other. ^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 14: Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
STELLAR (2019) NCT02397928	1. Not randomized	1. Not blinded		1. 8 patients lost to follow-up (10%)		

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias. ^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician. ^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication. ^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials). ^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference. ^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Retrospective Studies

Kutuk et al (2022) published a single-arm retrospective study of 5 patients with unresectable MPM who received TTF therapy from 2019 to 2021 at a single center in the US. The median follow-up was 5.4 months (range, 1.1 to 20.9). All patients were also treated with pemetrexed plus platinum-based chemotherapy. The median number of 4-week TTF cycles was 5 (range, 2 to 7) and the median TTF device usage in the first 3 months was 12.5 hours per day (range, 5 to 16.8). Treatment-related dermatitis was the only side effect associated with TTF and was reported as grade 1 to 2 in all patients; no patient had grade 3+ device-related toxicities. The authors note that this was the first publication of real-world implementation of TTF for MPM.

Section Summary: Tumor Treating Fields Therapy as an Adjunct or Alternative to Standard Medical Therapy for Unresectable, Locally Advanced, or Metastatic Malignant Pleural Mesothelioma

For individuals with metastatic MPM, TTF therapy has been evaluated in a prospective, single-arm study conducted in 80 patients (STELLAR) and a retrospective study of 5 US patients. The STELLAR study enrolled 80 patients with inoperable, previously untreated MPM who were treated with cisplatin or carboplatin in combination with TTF therapy at 12 sites outside the U.S. Median OS was 18.2 months and median PFS was 7.6 months. Seventy-two of the 80 individuals enrolled had at least 1 follow-up CT scan. Of those, 40% had a partial response, 57% had stable disease, and 3% progressed. Because there was no control group, it is not possible to draw conclusions about the effectiveness of TTF therapy compared to standard medical care alone. Additional limitations include the small sample size and no reporting of symptoms or quality of life outcomes. The retrospective study is the first publication of real-world implementation of TTF for MPM.

Tumor Treating Fields Therapy with Concurrent Standard Care for Metastatic Non-small Cell Lung Cancer

Clinical Context and Therapy Purpose

The purpose of TTF therapy, also referred to as alternating electrical field therapy, is to provide a treatment option that is an alternative to or an improvement on existing therapies for individuals with metastatic NSCLC. Tumor treating fields therapy has been investigated with a concurrent immune checkpoint inhibitor (i.e., a PD-1/PD-L1 inhibitor) or docetaxel after progression on or after a platinum-based regimen.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals who have metastatic NSCLC and are concurrently treated with an immune checkpoint inhibitor or docetaxel and have progressed on or after a platinum-based regimen.

Interventions

Optune branded products (formerly NovoTTF-100A System) are the only legally marketed TTF delivery system available in the United States. The portable, battery-powered device is carried in a bag while carrying out activities of daily living. For the treatment of NSCLC, transducer arrays with insulated electrodes are applied to the patient's chest cavity. The transducer array layout is typically determined using specialized software. The minimum daily treatment for NSCLC is 12 hours.

Comparators

The following practice is currently being used to make decisions about metastatic NSCLC in individuals who have progressed on or after a platinum-based regimen: PD-1/PD-L1 inhibitors or docetaxel alone.

Outcomes

The general outcomes of interest are whether TTF improves survival or quality of life. The time of interest for both PFS and OS is months to years.

Study Selection Criteria

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Randomized Controlled Trials

Leal et al (2023) published results of LUNAR (NCT02973789), a randomized, open-label phase 3 RCT that evaluated TTF for metastatic NSCLC.²⁶ The trial included 276 patients from 130 sites who had metastatic NSCLC and were receiving an immune checkpoint inhibitor (nivolumab, pembrolizumab, or atezolizumab) or docetaxel following progression on or after platinum-based therapy (see Table 15). Patients were at least 22 years of age and had an Eastern Cooperative Oncology Group (ECOG) performance status of 0 to 2. The primary endpoint was OS with key secondary endpoints of PFS and overall response rate. Median follow-up was 10.6 months (interquartile range [IQR, 6.1-33.7) for patients receiving TTF therapy with standard therapy, and 9.5 months (IQR, 0.1-32.1) for patients receiving standard therapy. Results are summarized in Table 16.

Table 15. Key Randomized Controlled Trial Characteristics for Metastatic NSCLC

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					<i>Active</i>	<i>Comparator</i>
Leal et al (2023); LUNAR	Countries in North America, Europe, and Asia	130	2017-2021	276 patients with metastatic NSCLC who had progression on or after platinum-based therapy	TTF to achieve an average of at least 18 h/d plus immune checkpoint inhibitor or docetaxel (n=137)	Immune checkpoint inhibitor or docetaxel alone (n=139)

NSCLC: non-small cell lung cancer; TTF: tumor treatment fields.

Table 16. Key Randomized Controlled Trial Results for Metastatic NSCLC

Study	Median OS (95% CI), months	Median PFS (95% CI), months	Overall response rate (95% CI), %	Overall AEs, %	Serious AEs, %
Leal et al (2023); LUNAR					
TTF + standard therapy	13.2 (10.3 to 15.5)	4.8 (4.1 to 5.7)	20.4 (14.0 to 28.2)	97	19
Standard therapy alone	9.9 (8.1 to 11.5)	4.1 (3.1 to 4.6)	17.3 (11.4 to 24.6)	91	15
HR (95% CI)	0.74 (0.56 to 0.98)	0.85 (0.67 to 1.11)	NR	NR	NR
p-value	.035	.23	.5	NR	NR

AEs: adverse events; CI: confidence interval; HR: hazard ratio; NR: not reported; OS: overall survival; PFS: progression-free survival; TTF: tumor treatment fields.

Tables 17 and 18 display notable limitations identified in this trial; a major limitation is the lack of patient blinding to treatment assignment. In addition, due to the rapidly changing landscape in the treatment of NSCLC, many of the patients did not receive current standard of care immunotherapy.

Table 17. Study Relevance Limitations

Study; Trial	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
Leal et al (2023); LUNAR	3. Enrolled population did not fully reflect treatment with current baseline testing and medical therapy standard of care				

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 18. Study Design and Conduct Limitations

Study; Trial	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Leal et al (2023); LUNAR		1. No sham control and not blinded to treatment assignment			4. Original sample size of 534 patients was reduced to 276 at an unplanned interim analysis	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

Section Summary: Tumor Treating Fields Therapy with Concurrent Standard Care for Metastatic Non-small Cell Lung Cancer

Analysis of the LUNAR trial, a phase 3, open-label RCT in 267 patients, found a significant 3.3-month OS improvement when TTF was added to standard care (immune checkpoint inhibitor or docetaxel) compared to standard care alone. Overall response rates and PFS were not significantly different between groups, and serious AEs were more common in the group treated with TTF. The trial is limited by the lack of a sham comparator, a lack of baseline molecular testing, and changes in standard of care.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Clinical Input from Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2016 Input

In response to requests, input was received from 3 physician specialty societies (1 of which provided 6 responses and 2 of which provided 1 response each) and 1 academic medical center (total of 9 individual responses) while this policy was under review in 2016. There was majority support, but not consensus, for the use of tumor treatment fields therapy as an adjunct to maintenance treatment following initial therapy for glioblastoma multiforme. There was mixed support for the use of tumor treatment fields as an alternative to chemotherapy in advanced or recurrent glioblastoma multiforme.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

Congress of Neurological Surgeons

In 2022, the Congress of Neurological Surgeons released guidelines on role of cytotoxic chemotherapy and other cytotoxic therapies in the management of progressive glioblastoma. In regard to TTF use in adult patients with progressive glioblastoma, the Congress states that "the use of TTF with other chemotherapy may be considered when treating adult patients with progressive glioblastoma [pGBM]. There is insufficient evidence to recommend TTF to increase overall survival in adult patients with pGBM".

The National Comprehensive Cancer Network (NCCN): Glioblastoma

National Comprehensive Cancer Network guidelines on central nervous system cancers (v.5.2024) include recommendations for the treatment of glioblastoma (see the Table 19).

Initial Glioblastoma: For the initial treatment of individuals with glioblastoma with good performance status and either methylated or unmethylated or indeterminate O⁶-methylguanine-DNA methyltransferase promoter status, treatment with standard brain radiotherapy plus concurrent temozolomide and adjuvant temozolomide plus alternating electric field therapy is a category 1 recommendation. Alternating electric currents therapy is only an option for patients with supratentorial disease.

Recurrent Glioblastoma: Consideration of alternating electric field therapy for recurrent glioblastoma is a category 2B recommendation.

Table 19: Guidelines for Adjuvant Treatment of Glioblastoma, by Age and Performance Status

Age, y	KPS Score, %	Treatment Options	Category
≤ 70	≥ 60	<ul style="list-style-type: none"> Standard RT plus concurrent and adjuvant temozolomide plus TTF (preferred) Standard RT plus concurrent and adjuvant temozolomide 	1
≤ 70	≥ 60	<ul style="list-style-type: none"> Standard RT alone (for unmethylated MGMT promoter status only) 	2A
≤ 70	≥ 60	<ul style="list-style-type: none"> Standard RT plus concurrent and adjuvant lomustine and temozolomide (for methylated or indeterminate MGMT promoter status only) 	2B
≤ 70	< 60	<ul style="list-style-type: none"> Hypofractionated RT with/without concurrent or adjuvant temozolomide Temozolomide alone Palliative/best supportive care 	2A
> 70	≥ 60	<ul style="list-style-type: none"> Hypofractionated RT plus concurrent and adjuvant temozolomide (for methylated or indeterminate MGMT promoter status only) 	1

Age, y	KPS Score, %	Treatment Options	Category
		<ul style="list-style-type: none"> Standard RT plus concurrent and adjuvant temozolomide plus TTF 	
> 70	≥ 60	<ul style="list-style-type: none"> Standard RT plus concurrent and adjuvant temozolomide Temozolomide alone (for methylated or indeterminate MGMT promoter status only) Hypofractionated RT alone (for unmethylated MGMT promoter status only) 	2A
> 70	≥ 60	<ul style="list-style-type: none"> Hypofractionated RT alone (for methylated or indeterminate MGMT promoter status only) 	2B
> 70	< 60	<ul style="list-style-type: none"> Hypofractionated brain RT alone Temozolomide alone Palliative/best supportive care 	2A

KPS: Karnofsky Performance Status; MGMT: O⁶-methylguanine-DNA-methyltransferase; RT: radiotherapy; TTF: tumor treating fields.

Malignant Pleural Mesothelioma

The National Comprehensive Cancer Network guidelines on malignant pleural mesothelioma (v.1.2024) do not address tumor treating fields (TTF) as a treatment option for malignant pleural mesothelioma.

Non-Small Cell Lung Cancer (NSCLC)

The National Comprehensive Cancer Network guidelines on non-small cell lung cancer (NSCLC)(v.3.2025) do not address TTF as a treatment option for NSCLC.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review can be located at clinicaltrials.gov.

REFERENCES

1. National Cancer Institute (NCI). Adult Central Nervous System Tumors Treatment (PDQ)Health Professional Version. Updated March 28, 2025; https://www.cancer.gov/types/brain/hp/adult-brain-treatment-pdq#cit/section_1.1 Accessed September 2, 2025.
2. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Central Nervous System Cancers. Version 2.2025. https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf Accessed September 2, 2025.
3. National Cancer Institute (NIH). Adult Central Nervous System Tumors Treatment (PDQ)-Health Professional Version; Updated March 28, 2025 https://www.cancer.gov/types/brain/hp/adult-brain-treatment-pdq#cit/section_1.1 Accessed September 2, 2025
4. Stupp R, Wong ET, Kanner AA, et al. NovoTTF-100A versus physician's choice chemotherapy in recurrent glioblastoma: a randomized phase III trial of a novel treatment modality. Eur J Cancer. Sep 2012;48(14):2192- 2202. PMID 22608262
5. U.S. Food and Drug Administration (FDA). Tumor treatment fields. NovoTTF-10A System. Summary of safety and effectiveness data (SSED). Premarket Approval Application (PMA)

- No. P100034. 2011; http://www.accessdata.fda.gov/cdrh_docs/pdf10/P100034b.pdf
Accessed September 2, 2025
6. U.S. Food and Drug Administration (FDA). Instructions for Use. Optune. NovoTTF-100A System. www.accessdata.fda.gov/cdrh_docs/pdf10/p100034s013c.pdf Accessed September 2, 2025
 7. Wong ET, Lok E, Swanson KD, et al. Response assessment of NovoTTF-100A versus best physician's choice chemotherapy in recurrent glioblastoma. *Cancer Med.* Jun 2014;3(3):592-602. PMID 24574359
 8. Ram Z, Gutin PH. Subgroup and quality of life analyses of the phase III clinical trial of NovoTTF-100A versus best standard chemotherapy for recurrent glioblastoma. *Neuro-Oncology.* 2010;12:iv48-iv49. PMID
 9. Ram Z, Wong ET, Gutin PH. Comparing the effect of novottf to bevacizumab in recurrent GBM: A post-HOC sub-analysis of the phase III trial data. *Neuro-Oncology.* 2011;13
 10. Salzberg M, Kirson E, Palti Y, et al. A pilot study with very low-intensity, intermediate-frequency electric fields in patients with locally advanced and/or metastatic solid tumors. *Onkologie.* Jul 2008;31(7):362-365. PMID 18596382
 11. Rulseh AM, Keller J, Klener J, et al. Long-term survival of patients suffering from glioblastoma multiforme treated with tumor-treating fields. *World J Surg Oncol.* 2012;10:220. PMID 23095807
 12. Turner SG, Gergel T, Wu H, et al. The effect of field strength on glioblastoma multiforme response in patients treated with the NovoTTF-100A system. *World J Surg Oncol.* 2014;12(1):162. PMID 24884522
 13. Stupp R, Kanner A, Engelhard H, Heidecke V, Taillibert S, Lieberman FS, Dbaly V, Kirson ED, Palti Y, Gutin PH. A prospective, randomized, open-label, phase III clinical trial of NovoTTF-100A versus best standard of care chemotherapy in patients with recurrent glioblastoma. *J Clin Oncol.* 2010 Jun 20;28(18 Suppl 1):LBA2007. Also available: https://ascopubs.org/doi/10.1200/jco.2010.28.18_suppl.lba2007 Accessed September 2, 2025.
 14. Novocure NovoCure Ltd. Sponsor executive summary: P100034. Novocure Ltd. NOVOTTF-100A system for the treatment of recurrent glioblastoma multiforme (GBM). Rockville (MD): U.S. Food and Drug Administration (FDA); 2011 Mar 17. 122 p. Also available: <https://hemonc.org/w/images/0/0c/Novottf.pdf> Accessed September 2, 2025
 15. Kanner AA, Wong ET, Villano JL, et al. Post Hoc analyses of intention-to-treat population in phase III comparison of NovoTTF-100A system versus best physician's choice chemotherapy. *Semin Oncol.* Oct 2014;41 Suppl 6:S25-34. PMID 25213871
 16. Mrugala MM, Engelhard HH, Dinh Tran D, et al. Clinical practice experience with NovoTTF-100A system for glioblastoma: The Patient Registry Dataset (PriDe). *Semin Oncol.* Oct 2014;41 Suppl 6:S4-S13. PMID 25213869
 17. FDA approves drug for treatment of aggressive brain cancer [FDA news release]. [internet]. Silver Spring (MD): U.S. Food and Drug Administration (FDA); 2009 May 8 [accessed 2015 May 22]. [2 p]. Available: <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm152295.htm>
 18. Stupp R, Brada M, van den Bent MJ, Tonn JC, Pentheroudakis G, ESMO Guidelines Working Group. High-grade glioma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2014 Sep;25 Suppl 3:iii93-101. Also available: <http://dx.doi.org/10.1093/annonc/mdu050>. PMID: 24782454 Accessed September 2, 2025
 19. Weller M, van den Bent M, Hopkins K, Tonn JC, Stupp R, Falini A, Cohen-Jonathan-Moyal E, Frappaz D, Henriksson R, Balana C, Chinot O, Ram Z, Reifenberger G, Soffietti R, Wick W, European Association for Neuro-Oncology (EANO) Task Force on Malignant Glioma. EANO guideline for the diagnosis and treatment of anaplastic gliomas and glioblastoma.

- Lancet Oncol. 2014 Aug;15(9):e395-403. Also available: [http://dx.doi.org/10.1016/S1470-2045\(14\)70011-7](http://dx.doi.org/10.1016/S1470-2045(14)70011-7). PMID: 25079102 Accessed September 2, 2025.
20. Pharma FW. FDA Grants Priority Review Status for Novocure's PMA Supplement Application of Optune in Newly Diagnosed Glioblastoma. 2015; <http://www.firstwordpharma.com/node/1282764#axzz3bpYYLTNx>
 21. Stupp R, Taillibert S, Kanner AA, et al. Maintenance therapy with tumor-treating fields plus temozolomide vs temozolomide alone for glioblastoma: a randomized clinical trial. JAMA. Dec 15 2015;314(23):2535-2543. PMID 26670971
 22. Wong ET, Lok E, Swanson KD. Clinical benefit in recurrent glioblastoma from adjuvant NovoTTF-100A and TCCC after temozolomide and bevacizumab failure: A preliminary observation. Cancer Med. 2015;4(3):383-391.
 23. Food and Drug Administration. FDA approves expanded indication for medical device to treat a form of brain cancer. October 5, 2015. FDA: Silver Spring, MD.
 24. McNamara MG, Sahebjam S, Mason WP. Emerging biomarkers in glioblastoma. Cancers. 2013;5(3):1103-19. Also available: <https://www.mdpi.com/2072-6694/5/3/1103>. PMID: 24202336. Accessed September 2, 2025
 25. NovoCure Ltd. Effect of novoTTF-100A in non-small cell lung cancer (NSCLC) patients with 1-3 brain metastases following stereotactic radio-surgery (SRS). In: ClinicalTrials.gov [database online]. Bethesda (MD): National Library of Medicine (U.S.); 2000- [accessed 2013 Apr 03]. [4 p]. Available: <http://www.clinicaltrials.gov/ct2/show/NCT01755624> NLM Identifier: NCT01755624 Accessed September 2, 2025.
 26. Novocure Ltd. Novocure announces the completion of enrollment of its PANOVA trial of tumor treating fields plus gemcitabine for treatment of pancreatic cancer. St. Helier (Jersey): Novocure Ltd.; 2015 Jan 12. 2 p. Also available: http://www.novocure.com/~media/Files/N/Novocure/press-release/2015/201502_PANOVA_Last_Patient.pdf.
 27. Novocure Ltd. Novocure enrolls first patient in a pilot study of tumor treating fields (TTFIELDS) together with paclitaxel for the treatment of recurrent ovarian cancer. St. Helier (Jersey): Novocure Ltd.; 2014 Oct 30. 2 p. Also available: <http://www.novocure.com/~media/Files/N/Novocure/press-release/2014/201405-ef22-fpi-press-release.pdf>.
 28. Taphoorn MJB, Dirven L, Kanner AA, et al.(2018) Influence of treatment with tumor-treating fields on health-related quality of life of patients with newly diagnosed glioblastoma: a secondary analysis of a randomized clinical trial. JAMA Oncol. Apr 1 2018;4(4):495-504. PMID 29392280
 29. Trusheim J, Dunbar E, Battiste J, et al. A state-of-the-art review and guidelines for tumor treating fields treatment planning and patient follow-up in glioblastoma. CNS Oncol. 2017 Jan;6(1):29-43. Epub 2016 Sep 15
 30. National Cancer Institute (NCI) – Adult Brain Tumors Treatment (PDQ®). Last modified March 28,2025. Available from: <http://www.cancer.gov/cancertopics/pdq/treatment/adultbrain/HealthProfessional> Accessed on September 2, 2025.
 31. Zhu JJ, Demireva P, Kanner AA, et al. Health-related quality of life, cognitive National Cancer Institute (NCI). Adult screening, and functional status in a randomized phase III trial (EF-14) of tumor treating fields with temozolomide compared to temozolomide alone in newly diagnosed glioblastoma. J Neurooncol 2017; 135:545.
 32. Stupp, R, Taillibert S, Kanner A, et al. Effect of Tumor-Treating Fields Plus Maintenance Temozolomide vs Maintenance Temozolomide Alone on Survival in Patients With Glioblastoma. A Randomized Clinical Trial.. *Jama*. 2017;318(23):2306-2316. PMID 29260225

33. Magouliotis, DE, Asprodini, EK, Svokos, KA, Tasiopoulou, VS, Svokos, AA, and Toms, SA. Tumor-treating fields as a fourth treating modality for glioblastoma: a meta-analysis. *Acta Neurochirurgica*. 2018;160(6):1167-1174.
34. X Armoiry, P Auguste, C Dussart, J Guyotat, M Connock, P14.12 The cost-effectiveness of tumor-treating fields in patients with newly diagnosed glioblastoma, *Neuro-Oncology*, Volume 21, Issue Supplement_3, August 2019, Pages iii68–iii69, <https://doi.org/10.1093/neuonc/noz126.247> Accessed September 2, 2025
35. Mittal S, Klinger NV, Michelhaugh SK, et al. Alternating electric tumor treating fields for treatment of glioblastoma: Rationale, preclinical, and clinical studies. *J Neurosurg*. 2018;128(2):414-421.
36. Vergote I, von Moos R, Manso L, et al. Tumor treating fields in combination with paclitaxel in recurrent ovarian carcinoma: Results of the INNOVATE pilot study. *Gynecol Oncol*. 2018 Jul 27 [Epub ahead of print].
37. Riley, M. M., San, P., Lok, E., Swanson, K. D., Wong, E. T. The Clinical Application of Tumor Treating Fields Therapy in Glioblastoma. *J. Vis. Exp.* (146), e58937, doi:10.3791/58937 (2019).
38. Novocure. Novocure announces Optune Lua as the brand name for the NovoTTF-100L system. 2020; <https://www.novocure.com/novocure-announces-optune-lua-as-the-brand-name-for-the-novottf-100l-system/>. Accessed September 2, 2025
39. Ceresoli GL, Aerts JG, Dziadziuszko R, et al. Tumour Treating Fields in combination with pemetrexed and cisplatin or carboplatin as first-line treatment for unresectable malignant pleural mesothelioma (STELLAR): a multicentre, single-arm phase 2 trial. *Lancet Oncol*. Dec 2019; 20(12): 1702-1709. PMID 31628016
40. Davies AM, Weinberg U, Palti Y. Tumor treating fields: a new frontier in cancer therapy. *Ann N Y Acad Sci*. Jul 2013; 1291: 86-95. PMID 23659608
41. Pless M, Weinberg U. Tumor treating fields: concept, evidence and future. *Expert Opin Investig Drugs*. Aug 2011; 20(8): 1099-106. PMID 21548832
42. Stupp R, Taillibert S, Kanner A, et al. Effect of Tumor-Treating Fields Plus Maintenance Temozolomide vs Maintenance Temozolomide Alone on Survival in Patients With Glioblastoma: A Randomized Clinical Trial. *JAMA*. Dec 19 2017; 318(23): 2306-2316. PMID 29260225
43. Kesari S, Ram Z. Tumor-treating fields plus chemotherapy versus chemotherapy alone for glioblastoma at first recurrence: a post hoc analysis of the EF-14 trial. *CNS Oncol*. Jul 2017; 6(3): 185-193. PMID 28399638
44. Mrugala MM, Engelhard HH, Dinh Tran D, et al. Clinical practice experience with NovoTTF-100A system for glioblastoma: The Patient Registry Dataset (PriDe). *Semin Oncol*. Oct 2014; 41 Suppl 6: S4-S13. PMID 25213869
45. National Institute for Health and Care Excellence (NICE) Guideline Brain Tumors (primary) and Brain Metastases in Adults (2018; Updated 2021) ISBN: 978-1-4731-3001-2 <https://www.nice.org.uk/guidance/ng99/resources/brain-tumours-primary-and-brain-metastases-in-adults-pdf-1837763558341> Accessed September 2, 2025
46. Batchelor T., Loeffler J.S., Wen PY, et al. UpToDate. Initial treatment and prognosis of IDH-wildtype glioblastoma in adults. Aug. 2021. https://www.uptodate.com/contents/initial-treatment-and-prognosis-of-newly-diagnosed-glioblastoma-in-adults?search=tumor%20treating%20fields&source=search_result&selectedTitle=1~6&usage_type=default&display_rank=1 Accessed September 2, 2025
47. Batchelor T., Shih H.A., Carter B.S., et al. UpToDate. Management of recurrent high-grade gliomas. May 2021 https://www.uptodate.com/contents/management-of-recurrent-high-grade-gliomas?search=tumor%20treating%20fields&source=search_result&selectedTitle=2~6&usage_type=default&display_rank=2 Accessed September 2, 2025

48. Weller, M., van den Bent, M., Preusser, M. et al. EANO guidelines on the diagnosis and treatment of diffuse gliomas of adulthood. *Nat Rev Clin Oncol* 18, 170–186 (2021). <https://doi.org/10.1038/s41571-020-00447-z> Accessed September 2, 2025
49. National Cancer Institute (NIH). Recurrent Cancer definition. <https://www.cancer.gov/publications/dictionaries/cancer-terms/def/recurrent-cancer> Accessed September 2, 2025
50. Jiang, H., Yu, K., Li, M., et al. 2020. 10.3389/fonc.2020.590648. 2408. Classification of Progression Patterns in Glioblastoma: Analysis of Predictive Factors and Clinical Implications. *Frontiers in Oncology*. <https://www.frontiersin.org/journals/oncology/articles/10.3389/fonc.2020.590648/full> Accessed September 2, 2025
51. U.S. Food and Drug Administration (FDA). Humanitarian Device Exemption (HDE). NovoTTF™-100L System. Summary of Safety and Probable Benefit. H180002. May 23, 2019. Available at: https://www.accessdata.fda.gov/cdrh_docs/pdf18/H180002B.pdf. Accessed September 2, 2025.
52. U.S. Food and Drug Administration (FDA). Neurological Devices Panel. Available at: <https://www.fda.gov/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/NeurologicalDevicesPanel/> Accessed September 2, 2025
53. U.S. Food and Drug Administration (FDA) Premarket Notification Database. Tumor Treatment Fields. NovoTTF-10A System. Summary of Safety and Effectiveness Data (SSED). No. P100034. Rockville, MD: FDA. April 8, 2011b. Available at: http://www.accessdata.fda.gov/cdrh_docs/pdf10/P100034b.pdf. Accessed September 2, 2025
54. U.S. Food and Drug Administration (FDA) Premarket Approval (PMA) Database. Premarket Notification Database. Tumor Treatment Fields. Optune™. Summary of Safety and Effectiveness Data (SSED). No. P100034/S013. Rockville, MD: FDA. October 5, 2015. Available at: http://www.accessdata.fda.gov/cdrh_docs/pdf10/p100034s013b.pdf. Accessed September 2, 2025.
55. PDQ® Adult Treatment Editorial Board. PDQ Adult Central Nervous System Tumors Treatment. Bethesda, MD: National Cancer Institute. Updated 01/18/2022. Available at: <https://www.cancer.gov/types/brain/hp/adult-brain-treatment-pdq> Accessed 09/22/2022. [PMID: 26389419]
56. National Brain Tumor Society. Glioblastoma Facts & Figures. <https://braintumor.org/take-action/about-gbm/>. Accessed September 2, 2025
57. U.S. Food and Drug Administration (FDA). Supplemental application for device name change. 2014; http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma_template.cfm?id=p100034s010 Accessed September 2, 2025
58. U.S. Food and Drug Administration (FDA). Summary of Safety and Effectiveness Data (SSED): Optune™ (formerly NovoTTF-100ATM System) 2015; https://www.accessdata.fda.gov/cdrh_docs/pdf10/P100034S013B.pdf Accessed September 2, 2025.
59. U.S. Food and Drug Administration (FDA). NovoTTF 100L System: Summary of Safety and Probable Benefit. May 23, 2019. Available at: https://www.accessdata.fda.gov/cdrh_docs/pdf18/H180002B.pdf Accessed September 2, 2025.
60. FDA Grants Breakthrough Device Designation to the NovoTTF-200T System for Advanced Liver Cancer. September 9, 2021. <https://www.novocure.com/fda-grants-breakthrough-device-designation-to-the-novottf-200t-system-for-advanced-liver-cancer> Accessed September 2, 2025.

61. Novocure. Novocure announces Optune Lua as the brand name for the NovoTTF-100L system. March 19, 2020; <https://www.novocure.com/novocure-announces-optune-lua-as-the-brand-name-for-the-novottf-100l-system>
62. Novocure Announces Presentations on Tumor Treating Fields Therapy, Including New Clinical Data and Real-World Evidence, at 2023 Society for Neuro-Oncology Annual Meeting. Novocure. Published November 10, 2023. <https://www.novocure.com/novocure-announces-presentations-on-tumor-treating-fields-therapy-including-new-clinical-data-and-real-world-evidence-at-2023-society-for-neuro-oncology-annual-meeting/>. Accessed September 2, 2025.
63. Regev O, Merkin V, Blumenthal DT, et al. Tumor-Treating Fields for the treatment of glioblastoma: a systematic review and meta-analysis. *Neurooncol Pract*. Aug 2021; 8(4): 426-440. PMID 34277021
64. Stupp R, Taillibert S, Kanner A, et al. Effect of Tumor-Treating Fields Plus Maintenance Temozolomide vs Maintenance Temozolomide Alone on Survival in Patients With Glioblastoma: A Randomized Clinical Trial. *JAMA*. Dec 19 2017; 318(23): 2306-2316. PMID 29260225
65. Stupp R, Taillibert S, Kanner AA, et al. Maintenance Therapy With Tumor-Treating Fields Plus Temozolomide vs Temozolomide Alone for Glioblastoma: A Randomized Clinical Trial. *JAMA*. Dec 15 2015; 314(23): 2535-43. PMID 26670971
66. Kutuk T, Appel H, Avendano MC, et al. Feasibility of Tumor Treating Fields with Pemetrexed and Platinum-Based Chemotherapy for Unresectable Malignant Pleural Mesothelioma: Single-Center, Real-World Data. *Cancers (Basel)*. Apr 16 2022; 14(8). PMID 35454925
67. Zhu JJ, Goldlust SA, Kleinberg LR, et al. Tumor Treating Fields (TTFields) therapy vs physicians' choice standard-of-care treatment in patients with recurrent glioblastoma: a post-approval registry study (EF-19). *Discov Oncol*. Oct 14 2022; 13(1): 105. PMID 36239858
68. Germano IM, Ziu M, Wen P, et al. Congress of Neurological Surgeons systematic review and evidence-based guidelines update on the role of cytotoxic chemotherapy and other cytotoxic therapies in the management of progressive glioblastoma in adults. *J Neurooncol*. Jun 2022; 158(2): 225-253. PMID 35195819
69. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Mesothelioma: Pleural. Version 2.2025. https://www.nccn.org/professionals/physician_gls/pdf/meso_pleural.pdf Accessed September 2, 2025.
70. National Palliative Care Research Center (NPCRC). Karnofsky Performance Scale. Available at: http://www.npcrc.org/files/news/karnofsky_performance_scale.pdf Accessed September 2, 2025
71. Hayes, a symplr company.. Health Technology Assessment. Tumor Treating Fields (Optune) for Treatment of Glioblastoma. Dec 27, 2019. Annual review: January 5 2023. Hayes,a symplr company. Available at: www.hayesinc.com. Accessed September 2, 2025.
72. Leal T, Kotecha R, Ramlau R, et al. Tumor Treating Fields therapy with standard systemic therapy versus standard systemic therapy alone in metastatic non-small-cell lung cancer following progression on or after platinum-based therapy (LUNAR): a randomised, open-label, pivotal phase 3 study. *Lancet Oncol*. Sep 2023; 24(9): 1002-1017. PMID 37657460
73. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Non-Small Cell Lung Cancer. Version 3.2025. https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed May 14, 2025.
74. U.S. FDA Premarket Approval. Optune Lua. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P230042>. Accessed May 12, 2025.

75. Tsao, Anne S. UpToDate. Systemic treatment for unresectable malignant pleural mesothelioma. February 2025 https://www.uptodate.com/contents/systemic-treatment-for-unresectable-malignant-pleural-mesothelioma?search=Role%20for%20tumor-treating%20fields%3F%20&source=search_result&selectedTitle=2~150&usage_type=default&display_rank=2#H1 Accessed September 8, 2025.

CODES

To report provider services, use appropriate CPT codes, HCPCS codes, Revenue codes, and/or ICD diagnosis codes.

Codes	Number	Description
CPT		
	1025T	Diagnostic imaging for computed tomography and magnetic resonance imaging simulation imported into the therapy treatment-planning system may be reported in addition to the alternating electric fields dosimetry.
HCPCS		
	E0766	Electrical stimulation device, used for cancer treatment, includes all accessories, any type
	A4555	Electrode/transducer for use with electrical stimulation device, used for cancer treatment, replacement only
Type of Service	Durable Medical Equipment	
Place of Service	Outpatient/Physician's Office	

POLICY HISTORY

Date	Reason	Action
September 2025	Annual Review	Policy Revised
September 2024	Annual Review	Policy Revised

Date	Reason	Action
September 2023	Annual Review	Policy Revised
October 2022	Annual Review	Policy Revised
March 2022	Interim Review	Policy Revised
October 2021	Annual Review	Policy Revised
November 2020	Annual Review	Policy Revised
March 2020	Interim Review	Policy Revised
November 2019	Annual Review	Policy Revised
June 2019	Interim Review	Policy Revised
October 2018	Annual Review	Policy Revised
October 2017	Annual Review	Policy Revised
October 2016	Annual Review	Policy Revised
October 2015	Annual Review	Policy Revised
November 2014		New Policy

New information or technology that would be relevant for Wellmark to consider when this policy is next reviewed may be submitted to:

Wellmark Blue Cross and Blue Shield
 Medical Policy Analyst
 PO Box 9232
 Des Moines, IA 50306-9232

*CPT® is a registered trademark of the American Medical Association.