



Wellmark Blue Cross and Blue Shield is an Independent Licensee of the Blue Cross and Blue Shield Association.

## QUANTITY LIMIT POLICY

# Oral Vancomycin Quantity Limit Policy

### NOTICE

This policy contains information which is clinical in nature. The policy is not medical advice. The information in this policy is used by Wellmark to make determinations whether medical treatment is covered under the terms of a Wellmark member's health benefit plan. Physicians and other health care providers are responsible for medical advice and treatment. If you have specific health care needs, you should consult an appropriate health care professional. If you would like to request an accessible version of this document, please contact customer service at 800-524-9242.

### BENEFIT APPLICATION

Benefit determinations are based on the applicable contract language in effect at the time the services were rendered. Exclusions, limitations or exceptions may apply. Benefits may vary based on contract, and individual member benefits must be verified. Wellmark determines medical necessity only if the benefit exists and no contract exclusions are applicable. This medical policy may not apply to FEP. Benefits are determined by the Federal Employee Program.

### DESCRIPTION

The intent of the Oral Vancomycin Quantity Limit policy is to ensure safe, appropriate use of Vancocin (vancomycin oral capsules) and Firvanq (vancomycin oral solution) based on product labeling, clinical guidelines, and clinical studies.

#### FDA-Approved Indications

##### **Vancocin**

Vancocin is indicated for the treatment of *Clostridioides difficile*-associated diarrhea. Vancocin is also used for the treatment of enterocolitis caused by *Staphylococcus aureus* (including methicillin-resistant strains) in adult and pediatric patients less than 18 years of age.

#### Limitations of Use

- Parenteral administration of vancomycin is not effective for the above infections; therefore, Vancocin capsules must be given orally for these infections.
- Orally administered Vancocin is not effective for other types of infections.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Vancocin capsules and other antibacterial drugs, Vancocin capsules should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

##### **Firvanq**

Firvanq is indicated for the treatment of *Clostridium difficile*-associated diarrhea in adults and pediatric patients less than 18 years of age.

Firvanq is also indicated for the treatment of enterocolitis caused by *Staphylococcus aureus* (including methicillin-resistant strains) in adults and pediatric patients less than 18 years of age.

#### Limitations of Use

- Parenteral administration of vancomycin is not effective for the above infections; therefore, vancomycin must be given orally for these infections.
- Orally administered vancomycin hydrochloride is not effective for treatment of other types of infections.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Firvanq and other antibacterial drugs, Firvanq should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

### **POLICY**

#### Quantity Limits Apply

| <u>Drug</u>   | <u>Limit*</u>         |
|---|-----------------------|
| Vancocin (vancomycin oral capsules 125mg, 250mg)    | 80 capsules / 10 days |
| Firvanq (vancomycin oral solution 25mg/mL, 50mg/mL) | 450 mL / 10 days      |

\* This drug is indicated for short-term acute use.

### **CLINICAL RATIONALE**

Vancocin capsules and Firvanq powder for oral solution are both indicated for the treatment of *C. difficile*-associated diarrhea. The American College of Gastroenterology last published guidelines on the diagnosis, treatment and prevention of *Clostridium difficile* infection (CDI) in 2013. Since that publication, there was a change in the taxonomic classification in 2016, with the organism assigned to a new genus and now called *Clostridioides difficile*. The US Centers for Disease Control and Prevention has adopted the new nomenclature, which has become standard throughout the scientific literature. Vancocin capsules and Firvanq powder for oral solution are also used for the treatment of enterocolitis caused by *Staphylococcus aureus* (including methicillin-resistant strains). Parenteral administration of vancomycin is not effective for the above infections; therefore, vancomycin must be given orally for these infections. Orally administered vancomycin is not effective for other types of infections. To reduce the development of drug-resistant bacteria and maintain the effectiveness of Firvanq, Vancocin capsules, and other antibacterial drugs, Firvanq and Vancocin capsules should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

The recommended dose of Vancocin and Firvanq for adults with *C. difficile*-associated diarrhea is 125 mg given four times daily for ten days. The recommended total daily dosage of Vancocin and Firvanq for adults with staphylococcal enterocolitis is 500 mg to 2 grams given in three or four divided doses for seven to ten days. For both *C. difficile*-associated diarrhea and staphylococcal enterocolitis in pediatric patients, the usual daily dosage of Firvanq and Vancocin is 40 mg/kg given in three or four divided doses for seven to ten days. The total daily dosage should not exceed 2 grams. Vancocin 125 mg and 250 mg capsules are available as two blister packs with ten capsules each, for a total of 20 capsules per carton. Firvanq is available as 25 mg/mL reconstituted to 150 mL or 300 mL and 50 mg/mL reconstituted to 150 mL or 300 mL.

The Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA) 2021 Focused Update Guideline of the Management of *Clostridioides difficile* infection in adults recommends a dosage for vancomycin of 125 mg taken orally four times a day for 10 days for an initial CDI episode and first recurrent episode of CDI. For a second or subsequent recurrence, a vancomycin dosing of 125 mg taken orally four times a day for 10 days, followed by rifaximin 400 mg three times daily for 20 days is recommended. Alternatively, a tapered/pulsed vancomycin regimen (e.g., 125 mg four times daily for 10-14 days, 2 times daily for 7 days, once daily for 7 days, and then every 2 to 3 days for 2 to 8 weeks) can be given for both a first CDI recurrence and a second or subsequent CDI recurrence. For fulminant CDI, vancomycin 500 mg four times daily by mouth or nasogastric tube is recommended. The American College of Gastroenterology (ACG) Clinical Guideline: Prevention, Diagnosis and Treatment of *Clostridioides difficile* Infections recommends oral vancomycin 125 mg four times daily for 10 days for the treatment of initial episodes of nonsevere CDI and initial episodes of severe CDI in adult patients. The ACG guideline suggests tapering/pulsed-dose vancomycin for adult patients experiencing a first recurrence of CDI and 500 mg of oral vancomycin every 6 hours for 48-72 hours in adult patients with fulminant CDI.

The Infectious Disease Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA) 2017 Clinical Practice Guideline for *Clostridium difficile* Infection in Adults and Children recommends an oral vancomycin dose of 10 mg/kg/dose four times daily for 10 days (maximum of 125 mg four times daily) for an initial episode of non-severe CDI and a first recurrence of non-severe CDI in children. For a second or subsequent recurrence of CDI in children, the guideline recommends oral vancomycin 10 mg/kg/dose four times daily for 10 days (maximum of 500 mg four times daily) followed by rifaximin for 20 days. Alternatively, oral vancomycin in a tapered and pulsed regimen (10 mg/kg/dose with max of 125 mg 4 times per day for 10 - 14 days, then 10 mg/kg with max of 125 mg 2 times per day for a week, then 10 mg/kg with max of 125 mg once per day for a week, and then 10 mg/kg with max of 125 mg every 2 or 3 days for 2 - 8 weeks) can be given for a second or subsequent recurrence of CDI in children. For an initial episode of severe/fulminant CDI in children, an oral vancomycin dose of 10 mg/kg/dose four times daily for 10 days (up to a maximum of 500 mg four times daily) is given with or without metronidazole.

The quantity limit will be 80 Vancocin capsules OR 450 mL Firvanq solution to accommodate a 10 day standard course of vancomycin for CDI or 10 days of treatment for staphylococcal enterocolitis. The maximum dose for staphylococcal enterocolitis is 2 grams given in three or four divided doses for ten days, which equates to 80 units of the 250 mg oral capsules and 400 mL of the 50 mg/mL oral solution. The initial limit of 450 mL of Firvanq is reflective of the available package sizes for the 50 mg/mL oral solution (150 mL or 300 mL). The maximum pulse dose for CDI is 125 mg four times per day for 14 days, two times per day for a week, once per day for one week, and then every two days for eight weeks, which is equal to a total of 13,125 mg. The maximum pulse dose equates to 105 units of the 125 mg oral capsules, 262.5 mL of the 50 mg/mL oral solution or 525 mL of the 25 mg/mL oral solution. All pulse treatment courses using the 125 mg capsules, or the 25 mg/mL solution can be fulfilled in two or less fills based on the initial quantity limits. A full pulse regimen can be met with one fill of the 50 mg/mL oral solution. For a maximum pulse dose using the 25 mg/mL solution or the 125 mg capsules, the quantity limit will cover at least 4 weeks of therapy before a refill is required. Since quantity limits are set at every 10 days, patients will have 2-3 weeks to obtain a refill to finish the pulse treatment.

Recurrent *Clostridium difficile* infection (rCDI) is generally defined as the recurrence of diarrhea and a confirmatory positive test (NAAT or EIA) within 8 weeks after treatment of an initial episode of CDI. Approximately 20% of patients will experience an initial recurrence, and rates of further recurrences continue to go up significantly after each one. Another course of antibiotics is generally required for the treatment of a first recurrence of CDI. Therefore, no limit has been placed on the number of fills per year.

If the patient is requesting more than the initial quantity limit, then the claim will reject with a message indicating that quantity limits are exceeded.

## PROCEDURES AND BILLING CODES

To report provider services, use appropriate CPT\* codes, Alpha Numeric (HCPCS level 2) codes, Revenue codes, and/or ICD diagnostic codes.

## REFERENCES

- Firvanq [package insert]. Wilmington, MA: Azurity Pharmaceuticals; February 2022.
- Vancocin [package insert]. Baudette, MN: ANI Pharmaceuticals, Inc.; December 2021.
- Lexicomp Online, AHFS DI (Adult and Pediatric) Online, Waltham, MA: UpToDate, Inc.; 2024; Accessed November 4, 2024.
- Micromedex (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: <https://www.micromedexsolutions.com>. Accessed November 4, 2024.
- McDonald L, Gerding D, Johnson S, et al. Clinical Practice Guidelines for *Clostridium difficile* Infection in Adults and Children: 2017 Update by the Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA). *Clinical Infectious Diseases* 2018;66 (7): e1-e48. <https://doi.org/10.1093/cid/cix1085>. Accessed September 26, 2022.
- Johnson S, Lavergne V, Skinner A et al. Clinical Practice Guideline by the Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA): 2021 Focused Update Guidelines on Management of Clostridioides difficile Infection in Adults, *Clinical Infectious Diseases* 2021;73 (5): e1029–e1044. <https://doi.org/10.1093/cid/ciab549> . Accessed September 26, 2022.
- Kelly CR, Fischer M, Allegretti JR, LaPlante K, et al. ACG Clinical Guidelines: Prevention, Diagnosis, and Treatment of Clostridioides difficile Infections. *Am J Gastroenterol*. 2021 Jun 1;116(6):1124-1147.

## POLICY HISTORY

**Policy #:** 05.02.71

**Original Effective Date:** September 14, 2017

**Reviewed:** August 2025

**Revised:** April 2019

**Current Effective Date:** May 16, 2019